

3RD INTERNATIONAL CONFERENCE: CONSTRUCTIVE MATHEMATICAL ANALYSIS

2-5 JULY 2025

ABSTRACT BOOK



The Third International Conference: Constructive Mathematical Analysis

Abstract Book

July 2-5, 2025
KONYA, TÜRKİYE



**This conference is supported by the Scientific and Technological Research Council of Türkiye (TÜBİTAK, 2223-B, 1929B022400901).*

CMA
CONSTRUCTIVE MATHEMATICAL ANALYSIS

FOREWORDS

Dear Participants,

It's my pleasure to chair The International Conference on "3rd International Conference: Constructive Mathematical Analysis". 3rd International Conference: Constructive Mathematical Analysis" is an activity of the journal Constructive Mathematical Analysis. The first two series of the event were organized as workshops. Based on the requests from international researchers studying on constructive mathematical analysis, this series was organized as a conference. Our 2025 conference was supported by Scientific Research Projects Coordinatorship of Selçuk University, The Scientific and Technological Research Council of Türkiye, Republic of Turkey Ministry of Youth and Sports Logo and BEYSU.

The purpose of this conference is to bring together experts and young analysts from all over the world working in different fields of mathematics and its applications to present their researches, exchange new ideas, discuss challenging issues, foster future collaborations and interact with each other.

The main goal of this conference is to promote, encourage, and provide a forum for the academic exchange of ideas and recent research works on any field of Analysis and Function Theory. The conference presented new results and future challenges, in a series of keynote lectures and contributed short talks.

We thank invited speakers distinguished **Prof. Francesco Altomare**, distinguished **Prof. Erdal Karapınar**, distinguished **Prof. Harun Karlı**, distinguished **Prof. Ioan Raşa**, distinguished **Prof. Gianluca Vinti**, distinguished **Prof. Xiaoming Wang** and distinguished **Prof. Ferenc Weisz** for contribution to the our conference.

The papers presented in this conference will be considered in the special issues of journals *ALTAY Conference Proceedings in Mathematics*, *Carpathian Mathematical Publications*, *Constructive Mathematical Analysis*, *Demonstratio Mathematica*, *Dolomites Research Notes on Approximation*, *Modern Mathematical Methods* and *Symmetry*.

This booklet contains the titles and abstracts of all invited and contributed talks at the *The Third International Conference: Constructive Mathematical Analysis* and is available on the conference website.

Prof. Dr. Tuncer ACAR
On Behalf of Organizing Committee - Chairman

to the memory of my uncle Hüseyin Acar



01.01.1966 - 02.01.2025

Chairman of Conference

Tuncer Acar
Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
tunceracar@ymail.com

Organizing Committee

Özlem Acar
Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
acarozlem@ymail.com

Orhan Boztaş
BSc, Yakutiye Anatolian High School, Türkiye

Borislav R. Draganov
Department of Mathematics and Informatics, Sofia University "St. Kliment Ohridski", Bulgaria
bdraganov@fmi.uni-sofia.bg

Mohamed A. Khamsi
University of Texas at El Paso, USA
mkhamsi@kfupm.edu.sa

Peter Massopust
Department of Mathematics, School of Computation, Information and Technology, Technical University of
Munich, Germany
massopust@ma.tum.de

Donal O'Regan
School of Mathematics, Statistics and Applied Mathematics, National University of Ireland, Ireland
donal.oregan@nuigalway.ie

Mahpeyker Öztürk
Department of Mathematics, Faculty of Science, Sakarya University, Türkiye
mahpeykero@sakarya.edu.tr

Ioan Raşa
Department of Mathematics, The Faculty of Automation and Computer Sciences, Technical University of
Cluj-Napoca, Romania
Ioan.Rasa@math.utcluj.ro

Yoshihiro Sawano
Department of Mathematics, Chuo University, Japan
yoshihiro-sawano@celery.ocn.ne.jp

Calogero Vetro
Department of Mathematics, Faculty of Science, University of Palermo, Italy
calogero.vetro@unipa.it

Dariusz Wardowski
Faculty of Mathematics and Computer Science, Department of Nonlinear Analysis, University of Lodz, Poland
dariusz.wardowski@wmii.uni.lodz.pl

Kehe Zhu
Department of Mathematics and Statistics State University of New York, USA
kzhu@math.albany.edu

Scientific Committee

Francesco Altomare
University of Bari Aldo Moro, Italy

Ali Aral
Kırıkkale University, Türkiye

Raul Curto
University of Iowa, USA

Barkat Bhayo
Lappeenranta University, Finland

Feng Dai
University of Alberta, Canada

Fuad Kittenah
University of Jordan, Jordan

Francisco Marcellan
Universidad Carlos III de Madrid, Spain

Manuel Lopez-Pellicer
University of Valencia, Spain

Guozhen Lu
University of Connecticut, USA

Adrian Petruşel
Babeş-Bolyai University, Cluj-Napoca

Jie Xiao
Memorial University, Canada

Local Organizing Committee

Osman Alagöz
Department of Mathematics, Bilecik Seyh Edebali University, Türkiye
osman.alagoz@bilecik.edu.tr

Murat Bodur
Department of Engineering Basic Sciences, Türkiye
mbodur@ktun.edu.tr

Fırat Özşarac
Department of Mathematics, Kirikkale University, Türkiye
frat_ozsarac@hotmail.com

Metin Turgay
Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
metinturgay@yahoo.com

Secretary

Samet Değle
Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
smt.degle@gmail.com

Tuğçe Delen
Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
tugcedelen@yahoo.com

Eşref Erdoğan
Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
esreferdogan@yahoo.com

Şerife Nur Karaman
Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
serifenurkaraman@yahoo.com

Sadettin Kurşun
National Defence University, Turkish Military Academy, Türkiye
sadettinkursun@yahoo.com

Dilek Özer

Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
dilekozer@yahoo.com

Aybala Sevde Özkapu

Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
aybalasevdeozkapu@yahoo.com

Contents

Local Convergence Problems and Positive Approximation Processes on Function Spaces on Convex Sets and Convex Cones	1
FRANCESCO ALTOMARE	
Primary Indispensable Remarks on Some Recent Results in Metric Fixed Point Theory	2
ERDAL KARAPINAR	
Wavelet-Based Approximation Operators: Applications to Bivariate Functions and Digital Image Processing	3
HARUN KARSLI	
Tracial Functionals on Operator Algebras	4
MOHAMMAD SAL MOSLEHIAN	
Sequences of Positive Linear Operators and Their Limits	5
IOAN RAŞA	
Approximation Results for a General Family of Nonlinear Operators and Applications: A Unifying Approach	6
GIANLUCA VINTI	
Construction of Transition Layers Associated with a Few Fluid Models	7
XIAOMING WANG	
Lebesgue Points and Summability of Higher Dimensional Fourier Series	8
FERENC WEISZ	
A restart derivative-free algorithm for solving pseudomonotone equations	9
AUWAL BALA ABUBAKAR	
Neimark-Sacker Bifurcation and Control of Chaos in Prey-Predator System Subject to Harvesting Effect on Prey	10
OZLEM AK GUMUS	
Integration of Modified Durrmeyer-Type Sampling Operators into Orlicz Spaces	11
OSMAN ALAGÖZ	
Generalized Split Equality Feasibility Problem in a Hadamard Space	12
BASHIR ALI	
Approximation with Nonlinear Lototsky-Durrmeyer Type Operators	13
HUSEYIN ERHAN ALTIN	
On Some Families of Exponential Type Polynomials	14

LAURA ANGELONI

Recent Results on the Weighted Approximation 15

ALI ARAL

Approximation Results on Neural Network Operators of Convolution Type 16

ASIYE ARIF

On Uniform Exhaustiveness and Korovkin-Type Theorem 17

ALI ARPACIOGLU

A Note on Modified Landau Type Operators 18

ALI HAN AVCI

On Some Direct Estimates Results of the Rate Approximation by the Kantorovich Operator in Weighted Grand Lebesgue Spaces 19

ISMAIL AYDIN

On Some Properties and Compactness Criteria of Variable Exponent Fofana's Space 20

ISMAIL AYDIN

Calculation of Two-Center Coulomb Integrals over STOs in Nonlined-up Coordinate System 21

ARIF AYKIN

Application of Wavelet Transformation in Analyzing Railway Track Profile Irregularities 22

ERDEM BALCI

Spectral Analysis of The Impulsive Sturm-Liouville Equation with Quadratic Spectral Parameters and Matrix Coefficients 23

GULER BASAK OZNUR

Quantitative Estimates for Wavelet Type Extension of Generalized Kantorovich Operators 24

GULEN BASCANBAZ-TUNCA

Robust Estimation of Parameters in Simple Linear Regression Model 25

OMER BAYRAKTAR

Minimum Hellinger Distance Estimate for a Periodic Fractional Autoregressive Model of First-Order 26

KARIMA BELAIDE

Solvability of a System Arising in Nonlinear Ultrasonic Heating 27

IMEN BENABBAS

On Modular Equation and Related Distortion Functions	28
BARKAT ALI BHAYO	
Estimation of a Finite Population Mean under Random Non-Response Using Improved Nadaraya-Watson Kernel Weights	29
NELSON KIPRONO BII	
Spectrum of the Operator $H(r, s, t, u, v)$ Defined on the Sequence Spaces ℓ_p and bv_p	30
MUSTAFA CEMIL BISGIN	
Recent Results on Sampling Kantorovich Operators of the Max-Product Type in Functional Spaces	31
LORENZO BOCCALI	
Approximation Properties of a New Generalization of Brass-Stancu Operators in L^p -Spaces	32
MURAT BODUR	
Cross-Approximation of Functions of 2 Variables	33
PETR BORODIN	
Fixed Point Results for Generalized Cyclic $\alpha - f\pi\varpi$ -Contractions	34
ABDERRAHMANE BOUDRAA	
Compactness Result in the Set of Measures	35
GHEORGHE BUCUR	
A Converse of Triangle Inequality: Applications	36
ILEANA BUCUR	
On Von Neumann Approximation Theorem	37
ILEANA BUCUR	
Some Novel Fixed Point Results in the Sense of Interpolative Metric Space	38
ABDURRAHMAN BUYUKKAYA	
Convergence in the Variation Seminorm by Generalized Kantorovich-Type Szász-Mirakyan Operators Constructed via Appell Polynomials	39
GÜL CANBOLAT	
A Novel Inflated Mixture Distribution with Exponential and Unit Lindley Components	40
ERDEM CANKUT	
Disjoint $\mathcal{F}$ -transitive Composition Operators on Orlicz Spaces	41
CHUNG-CHUAN CHEN	
Best Proximity Points for Cyclic Operators with Applications in Image Processing	42

MIHAELA ANCUTA CHIRA

Completeness Theorems on the Boundary 43

ALBERTO CIALDEA

On Three-Layer Conjugation Problem for the Heat Conduction Equation in a Rectangle 44

YUE CONGCONG

Hermite-Type Sampling Operator 45

ROSARIO CORSO

Nonlinear Reduced Modeling for FitzHugh-Nagumo Equation: A Comparison of Kernel Functions 46

YUSUF ÇAKIR

Commutator of Sharp Maximal Operator on Total Morrey Spaces 47

HAMIT C. CALISGAN

Commutator of Anisotropic Maximal Operator on Total Anisotropic Morrey Spaces 48

SULEYMAN CELIK

Approximation Theorem via Statistical ψ -convergence on $H_\omega(K)$ 49

SELIN CINAR

Generalized k -Lauricella Functions 50

YAHYA ÇIN

A New Formula of Delta Laplace Transform for the Anti-causal System Defined on Isolated Time Scales 51

HURIYE DAG

Construction of New Curves on n^{th} Degree Polynomial-Type Curves Using the Flc Frame 52

ESRA DAMAR

Local Laplacian Equienergetic Graphs 53

CAHIT DEDE

Approximation by Bivariate Modified Generalized Sampling Operators 54

SAMET DEGLE

Result for Multivalued Contractions on Ultrametric Spaces 55

TUGCE DELEN

Stability to Perturbations of S -fractions with Applications 56

MARTA DMYTRYSHYN

On the Approximation of the Lauricella-Saran's Hypergeometric Functions F_K and Their Ratios by Branched Continued Fractions 57

ROMAN DMYTRYSHYN

Approximation by Kantorovich Sampling Operators in Weighted Variable Exponent Lebesgue Spaces 58

BORISLAV R. DRAGANOV

On λ_f —Statistical Convergence of Sequences of Sets 59

BASIL ELSIBLI

On the Least Squares and Weighted Least Squares Estimation Based on Ranked Set Sampling 60

TENZILE ERBAYRAM

Trees Families with Algebraic Connectivity Greather than or Equal to $2(1 - \cos(\pi/9))$ 61

EMRECAN ERDEM

Fuzzy Soft Element and Its Application to Decision-making 62

BUSE ERDINC

New Fixed Point Theorem for Rational F-Contractions 63

ESREF ERDOGAN

A Theoretical Approach to Superfluous Parameters in Constrained Multivariate Linear Models 64

MELEK ERIS BUYUKKAYA

$d(d-1)$ -Continuous Multi-valued Mappings and Some Fixed Point Results 65

IREM EROGLU

Non-parametric Estimation for Doubly Geometric Process 66

GULTAC EROGLU INAN

Parameter Estimation for Doubly Geometric Process with the Inverse Gaussian Distribution 67

GULTAC EROGLU INAN

Quasi-metric Spaces and Related Fixed Point Results 68

ANDREEA FULGA

On the Two-layer Adjoint Problem for the Heat Conduction Equation in a Rectangle 69

AJIAHEER GEMINGNUER

On Controlled Interpolative Metric Spaces 70

EKBER GIRGIN

Commutator of Sharp Maximal Operator on Lorentz Spaces	71
SEZGIN GIRGINOL	
Comparison of Various Types of Inverse Soft Covering Upper Approximations	72
ZEHRA GÜZEL ERGÜL	
Generalizations of Certain Classes of Single-Valued Mappings to Multivalued Cases	73
EMIRHAN HACIOGLU	
On Groups and Vector Spaces with Easy Direct Limits	74
EMÍLIA HALUŠKOVÁ	
Solution of Split Equality Fixed Point Problem for Finite Family of Demigeneralized Mappings	75
MURTALA HARUNA HARBAU	
Preserving Non-Negativity in Rational Approximation of Fractional Diffusion Problems	76
STANISLAV HARIZANOV	
Anti-Periodic Boundary Value Problems for Singular Fractional p-Laplacian Equations	77
MAHIR HASANSOY	
Squared Basis Operators Related to Modified Bessel Functions	78
MONIKA HERZOG	
Non-Instantaneous Impulsive Differential Equations with a Caputo Fractional Derivative with respect to Other Functions and Strict Stability	79
SNEZHANA HRISTOVA	
Analysis of the mKdV Equation in Magnetohydrodynamics for Exploring Solitons and Nonlinear Dynamics	80
MUHAMMAD ABUBAKAR ISAH	
Fixed Points in Δ -symmetric Quasi Modular Metric Spaces of Non-Archimedean Type	81
DILEK KARABACAK KESIK	
A Comparative Stability Analysis of Fixed Point Iterations	82
RUMEYSA KARABOGA	
Benchmarking Iterative Processes on Contractive Mappings with Numerical Applications	83
HALIL IBRAHIM KARABOGA	
Remark on Neumann Problem in the Ring Domain	84
BAHRIYE KARACA	

Continued Fractions in Solving Congruences and Diophantine Equations	85
BUSRA KARACA	
Approximation by a Generalization of the Szász-Baskakov Operators	86
VERDA KARADAS	
Surface Pencil along a Common Spatial Geodezic According to Positional Adapted Frame in Euclidean 3-space	87
BILGE KARAHAN	
Convergence of Steklov Type Neural Network Operators	88
SERIFE NUR KARAMAN	
Quantitative Convergence and Multivariate Extensions of Activated Convolution Operators via Positive Linear Methods	89
SEDA KARATEKE	
Stability and Solution of Triple i - f -Hom-Der on Triple Banach Algebras	90
VAHID KESHAVARZ	
An Inverse Source Problem for Time-fractional Pseudoparabolic Equations with Memory Term	91
KHOMPYSH KHONATBEK	
Analysis of Approximation Processes for Fuzzy Numbers via Truncated Max-Product Operators	92
SEVILAY KIRCI SERENBAY	
Some Basic Results of Eccentricity Sombor Index	93
HAVVA KIRGIZ	
On Optimal BV-control Problem in Coefficients for Elliptic Equation with Variable Order of Nonlinearity	94
PETER KOGUT	
Korovkin Type Approximation Theorem via Statistical Relative Quasi Uniform Convergence	95
BURÇAK KOLAY	
Convergence of Remote Projections onto Convex Sets	96
EVA KOPECKÁ	
Modified Exponential Sampling Series in Logarithmic Weighted Spaces of Functions	97
SADETTIN KURSUN	
On the Generic Configurations of Two Dimensional Strongly Competing Systems	98
FLAVIA LANZARA	

Resolvent Approaches to Elliptic Regularity in Stationary Fokker-Planck Equations	99
HAESUNG LEE	
Boundary Integral Equations of the First Kind: Solvability and Applications	100
VITA LEONESSA	
Existence and Monotonicity of Principal Eigenvalues for a Class of Fractional Boundary Value Problems	101
YULONG LI	
Design Construction and Model Selection for Small Mixture-Process Variable Experiments with High-dimensional Model Terms	102
CHANG-YUN LIN	
Sensitivity Approach for the Evaluation of Statistical Test Batteries	103
ELENA ALMARAZ LUENGO	
Fake Nodes Approach and Some Applications	104
STEFANO DE MARCHI	
Rational Approximations of Fractional Elliptic Operators	105
SVETOZAR MARGENOV	
On the Complete Controllability of Fractional-Order Integrodifferential System Involving Generalized Conformable Derivatives	106
ROLA MASRI	
On Some Results Related to Positive Definite Matrices by Means of Cauchy-Schwarz Inequality	107
MAHMUT FURKAN MENIZ	
A New Generalization of the Frank Matrix	108
EFRUZ OZLEM MERSIN	
Highly Efficient Computational Technique for Equal Width Equation Modelling Dispersive Waves	109
MEHMET ALI MERSIN	
On Close to Convexity of a Class of Bounded Harmonic Mappings	110
OYA MERT COSKUN	
Solving Sequential Linear Fractional Differential Equations Using α -Trigonometric Functions	111
MUSSA YOUSEF MOHAMED	
Generalized Gamma-Type Operators	112
MIRELLA CAPPELLETTI MONTANO	

The Probabilistic Characteristics of Ion-Acoustic Waves in a Randomized Plasma Model with Non-extensive Electrons	113
ABDALLA HUSSEIN ZAKY MOUSTAFA	
Propagation of Solitons and Nonlinear Behavior in Nonlinear Power Law Fibers	114
AHMAD MUHAMMAD	
A Constructive Definition of the Riemann Integral on a Separable Banach Space	115
TIMOTHY MYERS	
Geometry of Curves as a way Establish Connections between Integrable Nonlinear Equations	116
GULGASSYL NUGMANOVA	
Zero Divisors of a Pythagorean Ring	117
RAYMOND CALVIN OCHIENG	
An Alternative Method for Associated Curves in Modified Orthogonal Frame	118
NURDAN OGRAS	
Fixed Point Theorems in Extended G-Menger Spaces and Application	119
LAHCEN OUMERTOU	
Fixed Point Theory in Graphical Modular Metric Spaces: A Unified Approach via Banach, Kannan and Reich-Type Contractions	120
ELIF OZBAY AZAP	
Convergence of a Family of Mellin-Steklov Type Exponential Sampling Operators	121
DILEK OZER	
A New Generalization of the Digital Contraction Principle and Image Compression Applications	122
AYBALA SEVDE OZKAPU	
Convergence Properties for a New Version of the Mellin-Gauss-Weierstrass Operators	123
FIRAT OZSARAC	
Modular Fixed Point Framework for Seljuk Knot Motifs: A Topological Study of Karatay Madrasa and Kubadabad Palace	124
MAHPEYKER ÖZTÜRK	
On Parabolic Class Number of a Special Subgroup of the Modular Group	125
SEDA ÖZTÜRK	
A New Proof of the Girard-Fermat Theorem	126
SEDA ÖZTÜRK	

Mappings Contracting Triangles	127
CRISTINA MARIA PăCURAR	
Properties of Hyperbolic Convolution on the Unit Disc	128
MARGIT PAP	
Optimal Recovery of Mappings Based on Linear Information with the help of T -splines in Banach Spaces	129
NATALIIA PARFINOVYCH	
Some Fixed Point Theorems in Modular b-Metric-Like Spaces	130
BANU PAZAR VAROL	
Recent Advancements on Semi-Discrete Sampling Operators in Sobolev-Orlicz Spaces	131
MICHELE PICONI	
A Study of Fixed Points in Soft S-Metric Spaces via Hybrid-Interpolative Reich-Istrecescu Type Mappings	132
HANDE POŞUL	
Normal Structure and Contractions which Diminish Radius in Metric and Banach Spaces	133
NAJAH REDJEL	
Sampling Type Operators Versus AI in Medical Image Processing	134
LUCREZIA RINELLI	
Respecting Cauchy Equivalence Classes	135
ANTHONY J. RIPA	
S_{pmk}^* Index for the Burr X-Perks Distribution with Applications	136
ŞULE SAĞLAM	
Variable-Order Hadamard-Type Fractional Calculus with respect to Function	137
IBRAHIM ABDULLAHI SALEH	
Adjoint Curve Using Conformable Derivative	138
BURCIN SALTİK BAEK	
Structural Modeling of Covariate-Driven Misclassification Risk: A Likelihood-Based Approach	139
SUMEYRA SERT	
An Inverse Problem for Pseudoparabolic Equation with Memory Term and Damping	140
SHAKIR AIDOS	

An Insight into Effects of Geography and Topography on the Urban Roadway Network Development of the Istanbul through Fractal Analysis	141
RECEP BILAL SIKAR	
Controlling Periodic Limit Cycles of Math Anxiety	142
SARA D. SONY	
Julia Sets-Geometric and Potential-Theoretic Insights	143
PRZEMYSŁAW SPRUS	
Evaluation of Lebesgue Constant with Admissible Meshes over Balls and Simplices	144
MATEUSZ SUDER	
Exploring State-Measure Dynamics in Algebraic Logic Structures	145
IBRAHIM SENTURK	
Mappings Contracting Perimeter of Triangles in Perturbed Metric Spaces	146
MIRELA ADRIANA TARNOVEANU	
Robust numerical method for the singularly perturbed problem with a nonlocal condition	147
ZELAL TEMEL	
An Approximation Study with Mellin Integral Operators	148
CEM TOPUZ	
Approximation Properties of Neural Network and Sampling Kantorovich Operators in Terms of Image Reconstruction	149
ARIANNA TRAVAGLINI	
New Generalization of Sampling Type Series and Applications to Image Processing	150
METIN TURGAY	
Geometric Evolution of Space Curves According to Positional Adapted Frame in Euclidean 3-Space	151
GÖKHAN TÜRK	
Approximation of Continuous Functions by Goodman-Sharma Type Modifications of the Bernstein Operator and the Baskakov Operator	152
RUMEN ULUCHEV	
A Generalized Approach to the Definition of Baskakov-Durrmeyer Operators	153
GÜLSÜM ULUSOY ADA	
Approximation Properties of Szász Mirakyan Durrmeyer Operators Preserving Exponential Functions	154
GÜLSÜM ULUSOY ADA	

Development of a New Efficient Algorithm for Multiplication of Large Integers MELTEM UZUN	155
On Some New Perspectives on Hardy-Amalgam Spaces with Variable Exponents CIHAN UNAL	156
Commutator of Anisotropic Sharp Maximal Operator on Lorentz Spaces NECATI UYAR	157
Characteristically Near Groups in Complex Plane Vector Fields TANE VERGILI	158
Local Fractional Hilbert-Type Inequalities PREDRAG VUKOVIĆ	159
Convergence Types on Interpolative Metric Spaces CEYLAN YALCIN	160
Extended Semi-Parallel Tensor Product Surfaces in $\mathbb{E}_2^4$ AZIZ YAZLA	161
Power Series Statistical Relative Uniform Convergence of a Double Sequence of Functions at a Point and Approximation Results SEVDA YILDIZ	162
On the Convergence of Generalized Szász-Mirakjan Operators in the Variation Seminorm SARE YUMRUÇALI	163
A Modification of Bernstein Operators on the Interval $(-1, 1)$ SULE YUKSEL GUNGOR	164
On Approximation by Algebraic Version of the Trigonometric Jackson Integrals $G_{s,n}$ in Generalized Weighted Integral Metric TEODORA ZAPRYANOVA	165
Estimating Covariance in Generalized Least Squares for Frechet Distribution KHEMISSA ZAHIA	166
Numerical Study and Convergence of a Finite Element Method for the Kazhikhov-Smagulov-Korteweg Pollution System GAFAROU ZONGO	167
On a Two-layer Conjugation Problem for the Heat-Conduction Equation in a Cylinder JIAYUE WANG	168

A Numerical Approach Based on a Dimensionality Reduction Procedure for Approximating Multiple Integrals	169
MOHAMED RAHAL	
Approximation Properties of Modified Jakimovski-Leviatan Operators	170
KADIR KANAT	

3rd International Conference: Constructive Mathematical Analysis

Local Convergence Problems and Positive Approximation Processes on Function Spaces on Convex Sets and Convex Cones

Francesco Altomare

Department of Mathematics, University of Bari Aldo Moro, Bari, Italy

francesco.altomare@uniba.it

key-words: Positive linear operator, local approximation, Bernstein-Chlodovsky operator.

Abstract:

Very recently, we developed some investigations ([1], [2], [3]) whose main aim was to deepen and to extend a general local convergence criterion for sequences of positive linear operators which was due to P. P. Korovkin ([6]).

The obtained new criteria guarantee whether a given linear function subspace, named here as local Korovkin subspace, satisfies the property that any sequence of positive linear operators locally converges on bounded locally continuous functions towards a given positive linear operator or towards the identity operator provided that the same property is checked on them.

The talk will be concerned with some new results in this direction which we have obtained in [1] and [2] where, among other things, it has been shown that for every given admissible subset S of continuous functions on a completely regular Hausdorff space, setting $S^2 := \{u^2 \mid u \in S\}$, then the linear subspace generated by $\{\mathbf{1}\} \cup S \cup S^2$ is a local Korovkin subspace for every weighted composition operator associated with a homeomorphic mapping and, in particular, for the identity operator.

As a consequence, we shall discuss some applications which concern the sequences of Bernstein-Schnabl operators acting on function spaces defined on convex Borel subsets as well as a new sequence of positive linear operators acting on function spaces defined on convex Borel cones.

Among other things such operators generalize a well-investigated modification of *Chlodovsky operators* on $[0, +\infty[$, also named *Bernstein-Chlodovsky operators*. Our approach describes a possibly unifying way on how to extend these operators to more general (both multidimensional and infinite-dimensional) settings as well.

The study is mainly addressed to highlight their pointwise convergence as well as their uniform convergence on compact subsets for functions which are bounded, Borel measurable and locally continuous.

In some particular cases, by means of such operators, a Weierstrass-type density result for the above mentioned class of functions in terms of polynomials is obtained as well.

References

- [1] Altomare, F. (2022). Korovkin-type Theorems and Local Approximation Problems. *Expo. Math.*, 40(4), 1229-1243.
- [2] Altomare, F. (2024). On the Convergence of Sequences of Positive Linear Operators towards Composition Operators. *Banach J. Math. Anal.*, 18(3).
- [3] Altomare, F. (2024). Local Korovkin-type Approximation Problems for Bounded Function Spaces. *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat.*, 118:88.
- [4] Altomare, F. (2025). Local Convergence Problems for Sequences of Positive Linear Operators. Preprint.
- [5] Altomare, F. (2025). On a generalization of Bernstein-Chlodovsky operators on convex cones. Preprint, 2025.
- [6] Korovkin, P. P. (1960). *Linear Operators and Approximation Theory* (translated from the Russian ed. (1959), Russian Monographs and Texts on Advances Mathematics and Physics, Vol III. Gordon and Breach Publishers, Inc. New York, Hindustan Publ. Corp. (India), Delhi).

3rd International Conference: Constructive Mathematical Analysis

Primary Indispensable Remarks on Some Recent Results in Metric Fixed Point Theory

Erdal Karapinar

Department of Mathematics, Atılım University, Ankara, Türkiye

erdalkarapinar@atilim.edu.tr

key-words: Metric fixed point, contractions, abstract metric structures.

Abstract:

The aim of this talk is to reveal the dilemmas of the fixed point theorems that were started to be constructed by Banach in 1922. It is possible to transform many of the real-world problems into fixed point theorems. This shows how the fixed point theory is useful and how huge application potential it has. On the other hand, a lot of results are repeated in this theory, which is of great interest. A substantial part of the new results suggested overlaps with the old results. To support this observation, I shall share a few examples from the current literature.

References

- [1] Karapinar, E. (2023). Recent Advances on Metric Fixed Point Theory: A Review. Appl. Math. Comput., 22(1), 3-30.
- [2] Karapinar, E. (2025). On the novelty of “Contracting Perimeters of Triangles in Metric Space”. Results Nonlinear Anal., 8(1), 115–123.

3rd International Conference: Constructive Mathematical Analysis

Wavelet-Based Approximation Operators: Applications to Bivariate Functions and Digital Image Processing

Harun Karsli

Department of Mathematics, Bolu Abant İzzet Baysal University, Bolu, Türkiye

karsli_h@ibu.edu.tr

key-words: Wavelet analysis, positive linear operators, approximation, bivariate operators, image reconstruction.

Abstract:

The aim of this talk is to demonstrate that the theory of wavelet-based operators plays a fundamental role in signal analysis and has significant applications in digital image processing. In classical Fourier analysis, signals are examined in the frequency domain by transforming them from the time domain. However, this transformation often results in the loss of time-related information, making the analysis of non-stationary signals particularly challenging. To address this limitation, various methods have been developed—one of the most prominent being wavelet analysis, originally introduced by Alfred Haar. Wavelets provide a powerful tool for decomposing signals into components that are localized in both time and frequency, allowing for more precise analysis of complex signals. This talk will focus especially on recent applications of wavelet methods to two-dimensional functions and image reconstruction. We will also explore how operators defined using specific types of wavelets serve as natural extensions of classical operators and their Kantorovich-type modifications. Notably, the wavelet-based operators and the associated algorithms presented here can be seamlessly applied to real-world problems traditionally handled by Kantorovich operators, thereby enhancing both the theoretical framework and practical effectiveness of signal and image processing techniques.

Acknowledgements: The author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Acar, T., Alagoz, O., Aral, A., Costarelli, D., Turgay, M. and Vinti, G. (2022). Approximation by Sampling Kantorovich Series in Weighted Spaces of Functions. *Turkish J. Math.*, 46(7), 2663-2676.
- [2] Acar, T., Costarelli, D. and Vinti, G. (2020). Linear Prediction and Simultaneous Approximation by m -th Order Kantorovich-type Sampling Series. *Banach J. Math. Anal.*, 14(4), 1481-1508.
- [3] Altomare, F., Campiti, M. (1994). Korovkin Type Approximation Theory and its Applications, De Gruyter Studies in Mathematics, 17, Walter de Gruyter and Co., Berlin.
- [4] Butzer, P. L. (1953). On Two Dimensional Bernstein Polynomials. *Canad. J. Math.*, 5, 107-113.
- [5] Butzer, P. L., Nessel, R. J. (1971). *Fourier Analysis and Approximation*, V.1, Academic Press, New York, London.
- [6] Costarelli, D., Vinti, G. (2013). Approximation by Nonlinear Multivariate Sampling Kantorovich Type Operators and Applications to Image Processing. *Numer. Funct. Anal. Optim.*, 34(8), 819-844.
- [7] Daubechies, I. (1992). *Ten Lectures on Wavelets*. CBMS-NSF Series in Appl. Math. 61, SIAM Publ. Philadelphia.
- [8] Karsli, H. (2023). On Wavelet Type Generalized Bézier Operators. *Math. Found. of Comp.*, 6(3), 439-452.
- [9] Karsli, H. (2024). Asymptotic Properties and Quantitative Results of the Wavelet Type Bernstein Operators. *Dolomites Res. Notes Approx.*, 17(1), 50-62.
- [10] Meyer, Y. (1993). *Ondelettes: Wavelets and Operators*. Yves Meyer. Cambridge University Press, New York. xvi, 223 pp.

3rd International Conference: Constructive Mathematical Analysis

Tracial Functionals on Operator Algebras

Mohammad Sal Moslehian

Department of Pure Mathematics, Ferdowsi University of Mashhad, Mashhad, Iran

moslehian@um.ac.ir

key-words: Trace, positive linear functional, matrix algebra, C^* -algebra, von Neumann algebra.

Abstract:

In this talk, we investigate tracial functionals on C^* -algebras and von Neumann algebras. We explore various characterizations of the standard trace on the full matrix algebra M_n and present several conditions under which states satisfy the tracial property. As applications, we establish new criteria for determining the commutativity of C^* -algebras.

References

- [1] Bikchentaev, A. M., Kittaneh, F., Moslehian, M. S. and Seo, Y. (2024). Trace Inequalities, for Matrices and Hilbert Space Operators. Forum for Interdisciplinary Mathematics, Springer, ISBN 978-981-97-6519-5.
- [2] Bikchentaev, A. M., Moslehian, M. S. (2025). Characterizations of Tracial Functionals on C^* -algebras. Infin. Dimens. Anal. Quantum Probab. Relat. Top., to appear.
- [3] Dadkhah, A., Kian, M. and Moslehian, M. S. (2024). Decomposition of Tracial Positive Maps and Applications in Quantum Information. Anal. Math. Phys., 14(3), Paper No. 48.

3rd International Conference: Constructive Mathematical Analysis

Sequences of Positive Linear Operators and Their Limits

Ioan Raşa

Department of Mathematics, Technical University of Cluj-Napoca, Cluj-Napoca, Romania

ioan.rasa@math.utcluj.ro

key-words: Positive linear operators, convergence, limit operator.

Abstract:

Usually, In Approximation Theory one investigates sequences of positive linear operators converging in some sense to the identity operator Id . However, some modifications of classical operators, inspired by the famous "Poisson approximation to the binomial distribution", lead to sequences of positive linear operators with limits different from Id . See, e.g., [2], [1] and the references therein. This talk is devoted to such sequences and to the characterization of their limits.

References

- [1] Acu, A.M., Heilmann, M., Raşa, I. and Seserman, A. (2023). Poisson Approximation to the Binomial Distribution: Extensions to the Convergence of Positive Operators. *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.*, 117:162.
- [2] de la Cal, J., Luquin, F. (1992). A Note on Limiting Properties of Some Bernstein-type Operators. *J. Approx. Theory*, 68(3), 322-329.

3rd International Conference: Constructive Mathematical Analysis

Approximation Results for a General Family of Nonlinear Operators and Applications: A Unifying Approach

Gianluca Vinti

Department of Mathematics and Computer Science, University of Perugia, Perugia, Italy

gianluca.vinti@unipg.it

key-words: Approximation, nonlinear operators, discrete operators, Orlicz spaces.

Abstract:

I will discuss some approximation results for a general family of nonlinear operators, proposing a general unifying approach. I will consider both pointwise and uniform convergence, and modular convergence in Orlicz spaces [4, 5]. Furthermore, I will also address the problem of qualitative estimates in suitable Lipschitz classes [6]. The theory presented allows to simultaneously treat families of discrete ([2, 1] and semi-discrete operators [3] (such as sampling or Durrmeyer-type sampling operators) and integral operators. The first ones (of sampling type) have also been studied in relation to concrete applications in the real world.

References

- [1] Angeloni, L., Vinti, G. (2017). Discrete Operators of Sampling Type and Approximation in φ -variation. Math. Nachr., 291(4), 546-555.
- [2] Bardaro, C., Butzer, P. L., Stens, R.L. and Vinti, G. (2010). Prediction by Samples from the Past with Error Estimates Covering Discontinuous Signals. IEEE, Transaction on Information Theory, 56, 614-633.
- [3] Piconi, M., Vinti, G. (2025). Semi-discrete Sampling in Sobolev-Orlicz Spaces. to be submitted.
- [4] Vinti, G., Zampogni, L. (2018). A General Approximation Approach for the Simultaneous Treatment of Integral and discrete operators. Adv. Nonlinear Stud., 18, 705-724.
- [5] Vinti, G., Zampogni, L. (2022). A General Method to Study the Convergence of Nonlinear Operators in Orlicz Spaces, Adv. Nonlinear Stud., 22, 594-618.
- [6] Vinti, G., Zampogni, L. (2025). An Abstract Setting for the Study of Quantitative and Qualitative Estimates of Nonlinear Operators, in preparation.

3rd International Conference: Constructive Mathematical Analysis

Construction of Transition Layers Associated with a Few Fluid Models

Xiaoming Wang

School of Mathematical Sciences, Eastern Institute of Technology, Ningbo, China
Department of Mathematics and Statistics, Missouri University of Science and Technology, Rolla, USA

wxm.nse@gmail.com, wxm.math@outlook.com

key-words: Transition layer, fluid model, stability, construction.

Abstract:

Most physical models contain parameters that are either very small or very large. The incompressible Navier-Stokes system at small viscosity (large Reynolds number) is a prime example. When the small parameter is set to zero, we often obtain a qualitatively different model whose behavior deviates significantly from that of the nonzero case. The zero-parameter model is usually simpler; therefore, we are interested in approximating the behavior of the nonzero model using the zero-parameter counterpart. However, such an approximation is not valid unless supported by a transition layer that bridges the two qualitatively different behaviors. The construction of such transition layers, as well as their stability and validity, will be the focus of this talk. Applications to the Navier-Stokes equations and the vanishing material interface limit of the convection in layered porous media, among others, will also be discussed. The results reported here are joint with H.J. Dong and R.M. Temam.

References

- [1] Dong, H. J., Wang, X., (2025). Sharp Material Interface Limit of the Darcy-Boussinesq System. to appear in SIAM J. Appl. Math., DOI:10.1137/25M1740310.
- [2] Wang, X., (2010). Examples of boundary layers associated with the incompressible Navier-Stokes flows. Chin. Ann. Math. ser. B, 31(5), 781-792.
- [3] Temam, R. M., Wang, X., (2002). Boundary Layer Associated with the Incompressible Navier-Stokes Equations: the Non-characteristic Boundary Case. J. Diff. Eqs., 179, 647-686.

3rd International Conference: Constructive Mathematical Analysis

Lebesgue Points and Summability of Higher Dimensional Fourier Series

Ferenc Weisz

Department of Numerical Analysis, Eötvös Loránd University, Budapest, Hungary

weisz@inf.elte.hu

key-words: Higher dimensional Fourier series, Hardy-Littlewood maximal functions, Fejér means, Cesaro means, Lebesgue point.

Abstract:

We consider the convergence of partial sums and of summability methods for higher dimensional Fourier series. We focus on the Cesàro and Riesz methods and the ℓ_q - and rectangular summability. We study the norm and almost everywhere convergence. We investigate five different types of Hardy-Littlewood maximal operators and their weak and strong boundedness on L_p spaces. We characterize the set of the almost everywhere convergence as different types of Lebesgue points.

3rd International Conference: Constructive Mathematical Analysis

A restart derivative-free algorithm for solving pseudomonotone equations

Auwal Bala abubakar¹, Hassan Mohammad² and Yunho Kim³

¹ Department of Art and Science, George Mason University, Songdomunhwa-ro 119-4 Yeonsu-gu, Incheon 21985, South Korea

² Numerical Optimization Research Group, Department of Mathematical Sciences, Faculty of Physical Sciences, Bayero University, Kano, Kano, Nigeria

³ Ulsan National Institute of Science and Technology, UNIST-gil 50, Ulsan, South Korea

aabubak@gmu.edu, hmuhd.mth@buk.edu.ng, yunhokim@unist.ac.kr

key-words: Iterative methods, Pseudo-monotone nonlinear equations, conjugate gradient method, Derivative-free method, Logistic regression.

Abstract: In this work, we propose an efficient derivative-free algorithm strengthened by a restart strategy for solving nonlinear pseudomonotone equations. We have shown that the algorithm's direction of search is descent and bounded and that the algorithm generates globally convergent sequences to the solutions under the assumptions of pseudomonotonicity and continuity. Moreover, the behavior of the convergent sequence near its limit is revealed for which we have shown that the sequence approaches its limit along the direction in the normal cone at the limit when the solution is not unique. Numerical simulations on benchmark test problems reveal the computational efficiency of our proposed algorithm in comparison with some existing recent ones. The suitability of the suggested algorithm for logistic regression problems is demonstrated, as well.

3rd International Conference: Constructive Mathematical Analysis

Neimark-Sacker Bifurcation and Control of Chaos in Prey-Predator System Subject to Harvesting Effect on Prey

Ozlem Ak Gumus

Department of Mathematics, Adiyaman University, Adiyaman, Türkiye

akgumus@adiyaman.edu.tr

key-words: Prey-Predator system, Neimark-Sacker bifurcation, stability, bifurcation theory, harvesting effect, chaos control.

Abstract:

This article aims to examine the dynamics of the positive coexistence fixed point of the discrete-time model given in [3] by taking the conversion efficiency of prey into predator as the bifurcation point. We show that the system is exposed to Neimark-Sacker bifurcation. We also control the Neimark-Sacker bifurcation with a feedback control strategy. Additionally, computer simulations are presented to verify the theoretical results obtained with the stability, bifurcation and chaos control strategy.

References

- [1] Brauer, F. Castillo-Chavez, C. (2001). Mathematical Models in Population Biology and Epidemiology. Springer-Verlag, New York.
- [2] Elaydi, S. N. (1996). An Introduction to Difference Equations. Springer-Verlag, New York.
- [3] Gumus, O. A., Elsadany, A. A., Yousef, A. M. and Agiza H.N. (2024). Theoretical and Numerical Bifurcation Analysis in A Prey-predator Model with Prey Harvesting Effort. Int. J. of Modeling, Simul., and Sci. Comput., 15(6), 22-39.
- [4] Gumus, O. A., Feckan, M. (2021). Stability, Neimark-Sacker Bifurcation and Chaos Control for a Prey-predator System with Harvesting Effect on Predator. Miskolc Math. Notes, 22(2), 663-679.
- [5] Kuznetsov, Y. A. (1998). Elements of Applied Bifurcation Theory. Springer-Verlag, New York.
- [6] Lotka, A. J. (1925). Elements of Physical Biology. Williams and Wilkins Company, Baltimore.
- [7] Volterra, V. (1927). Variations and Fluctuations in the Numbers of Coexisting Animal Species. Lect. notes in Biomath., Springer-Verlag, Berlin, Heidelberg, New York, 22(1), 65-273.

3rd International Conference: Constructive Mathematical Analysis

Integration of Modified Durrmeyer-Type Sampling Operators into Orlicz Spaces

Osman Alagöz

Department of Mathematics, Bilecik Seyh Edebali University, Bilecik, Türkiye

osman.alagoz@bilecik.edu.tr

key-words: Durrmeyer type sampling operator, Orlicz space, modular convergence.

Abstract:

This document details the integration of modified Durrmeyer-type sampling operators into Orlicz spaces. The primary focus is on proving boundedness, establishing modular convergence, and deriving error estimates, ensuring that these operators can approximate functions within the Orlicz space framework effectively.

Acknowledgements: The author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The authors thank to TUBITAK for their supports.

References

- [1] Alagöz, O. (2024). Weighted approximations by sampling type operators: recent and new results. *Constructive Mathematical Analysis*, 7(3), 114–125. <http://dergipark.org.tr/en/pub/cma>
- [2] Costarelli, D., Piconi, M., and Vinti, G. (2023). Quantitative estimates for Durrmeyer-sampling series in Orlicz spaces. *Sampling Theory, Signal Processing, and Data Analysis*, 21(3). <https://doi.org/10.1007/s43670-022-00042-6>

3rd International Conference: Constructive Mathematical Analysis

Generalized Split Equality Feasibility Problem in a Hadamard Space

Bashir Ali¹, Abdulnasir Bala Nuhu^{1,2}

¹ Department of Mathematical Sciences, Bayero University, Kano, Nigeria

² Department of Mathematics, Aliko Dangote, niversity of Science and Technology, Kano, Nigeris

bashiralik@yahoo.com, abdulnasirnuhu@gmail.com

key-words: Mixed equilibrium problem, convex joint Δ –lower semicontinuous function, generalized split equality feasibility problem.

Abstract:

In this paper, we introduce and study a Generalized Split Equality Feasibility Problem in the setting of Hadamard space. We construct a perturbed Halpern-viscosity type iterative algorithm and proved its strong convergence to solution of the above mentioned problem. Our technique of proof is new and of independent interest.

References

- [1] Alizadeh, S., Dehghan, H. and Moradlou, F. (2018) Δ -convergence theorem for inverse strongly monotone mappings in $CAT(0)$ spaces, Fixed Point Theory 19(1) , 45-56.
- [2] Byrne, C. (2004) A unified treatment of some iterative algorithms in signal processing and image reconstruction., Inverse Problems 20, no.1, 103-120.
- [3] Chidume, C. E. Bello, A.U. and Ndambonwe, P. (2014) Strong and Δ -convergence theorems for common fixed points of a finite family of multivalued demicontractive mappings in $CAT(0)$ spaces, Abstr. Appl. Anal.
- [4] Censor, Y. Segal, A. (2009) The split common fixed point problem for directed operators., J. Convex Anal. 16, no.2, 587-600.
- [5] Dehghan, H. Izuchukwu, C. Mewomo, O.T. Taba D.A. and Ugwunnadi, G.C. (2019) Iterative algorithm for a family of monotone inclusion problems in $CAT(0)$ spaces, Quest. Mat. 24, 1-24.
- [6] Dhompongsa, S. and Panyanak, B. (2008) On Δ -convergence theorems in $CAT(0)$ spaces, Comput.Math. Appl. 56 pp. 2572-2579.
- [7] Dhompongsa, S. Kirk W.A. and Panyanak, B. (2007) Nonexpansive set-valued mappings in metric and Banach spaces, J. Nonlinear Convex Anal. 8, 35-45.
- [8] Izuchukwu, C. Aremu, K.O. Oyewole, O.K. Mewomo, O.T. and Khan, S.H. (2019) On Mixed Equilibrium Problems in Hadamard Spaces, J.Maths 3210649, 13 pages.
- [9] Moudafi, A. (2014) Alternating CQ-algorithm for convex feasibility and split fixed point problems., J.Nonlinear Convex Anal., 15 (14), 809-818.
- [10] Osisiogu, U. A. Adum, F.L. and Efor, T.E. (2020) Strong convergence results for variational inequality problem in $CAT(0)$ spaces, Advances in Nonlinear Variational Inequalities 23(1), 84-101.
- [11] Stampacchia, G. (1968) Variational Inequalities in: Theory and Application of Monotone Operators, Proceedings of the NATO Advanced Study Institute, Venice, Edizioni Odersi, Gubbio, Italy, 1968, 102-192.
- [12] Tang, J. (2014) Viscosity approximation methods for a family of nonexpansive mappings in $CAT(0)$ spaces, Abstr. Appl. Anal., Article ID 389804, 9 pages.
- [13] Xu, H. K. (2002) Iterative algorithms for nonlinear operators., J. London. Math. Soc. 66, 240-256.

3rd International Conference: Constructive Mathematical Analysis

Approximation with Nonlinear Lototsky-Durrmeyer Type Operators

Huseyin Erhan Altin¹, Harun Karsli¹

¹ Department of Mathematics, Bolu Abant Izzet Baysal University, Bolu, Türkiye

erhanaltin@ibu.edu.tr, karsli_h@ibu.edu.tr

key-words: Lototsky operator, nonlinear Durrmeyer operator, rate of convergence.

Abstract:

In 1966, J. P. King [5] introduced Lototsky-Bernstein operators $L_n : C[0, 1] \rightarrow C[0, 1]$ for $n \in \mathbb{N}$ as follows:

$$L_n(f; x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) a_{n,k}(x), x \in [0, 1],$$

using the basis function $a_{n,k}(x)$ obtained from the following relation

$$\prod_{i=1}^n (h_i(x)y + 1 - h_i(x)) = \sum_{k=0}^n a_{n,k}(x) y^k, y \in \mathbb{R},$$
$$a_{n,k}(x) = \sum_{\substack{J \cup \bar{J} = \mathbb{N}_n \\ \text{Card}(J) = k}} \prod_{i \in J} (1 - h_i(x)) \prod_{i \in \bar{J}} h_i(x),$$

where $h_i : [0, 1] \rightarrow [0, 1]$ is a sequence of continuous functions and $a_{0,0}(x) = 1$, $a_{0,k}(x) = 0$ for $k > 0$. Recently, these operators have been studied by several mathematicians. Positive linear and nonlinear operators, convolution, moment and sampling operators have an important role in several branches of Mathematics. For instance, in reconstruction of signals and images, in Fourier analysis, operator theory, probability theory and approximation theory. In this way, the main objective of this work is to define a new sequence of nonlinear operators of type Lototsky-Durrmeyer and to give some convergence results for these type operators.

References

- [1] Abel, U., Agratini, O. (2021). On the Durrmeyer-Type Variant and Generalizations of Lototsky-Bernstein Operators. Symmetry, 13, 1841.
- [2] Karsli, H. (2014). On Convergence of a Sequence of Nonlinear Durrmeyer Operators. Pan American Math. J., 24(2), 1-24.
- [3] Karsli, H., Tiryaki, I. U. and Altin, H. E. (2014). Some Approximation Properties of a Certain Nonlinear Bernstein Operators. Filomat, 28(6), 1295-130524.
- [4] King, J. P. (1966). The Lototsky Transform and Bernstein Polynomials. Canad. J. Math., 18, 89-91.

3rd International Conference: Constructive Mathematical Analysis

On Some Families of Exponential Type Polynomials

Laura Angeloni¹, Danilo Costarelli² and Chiara Darielli¹

¹ Department of Mathematics and Computer Science, University of Perugia, Perugia, Italy

² Department of Mathematics and Computer Science, University of Firenze, Firenze, Italy

`laura.angeloni@unipg.it`, `danilo.costarelli@unipg.it`, `chiara.darielli@unifi.it`

key-words: Exponential type polynomials, modulus of continuity, Korovkin theorems, constructive approximation, Lipschitz classes.

Abstract:

We will present approximation results by means of some exponential type polynomials that generalize, in different directions, the classical Bernstein polynomials.

In particular, we will introduce a multidimensional version of the Bernstein-type exponential polynomials on the hypercube $[0, 1]^d$. For such operators, that fix product-type exponential functions and their square, we prove uniform convergence, by means of both the tools of Korovkin theory and a constructive approach, as well as a quantitative estimate of the order of approximation in terms of the modulus of continuity and, as a consequence, a qualitative result about the rate of convergence in some Lipschitz classes.

We will also discuss a new Kantorovich version of the exponential polynomials for which uniform convergence, convergence in weighted L^p -spaces, a Voronovskaja formula and results about the rate of approximation are established.

Acknowledgements This research was partially supported within the PRIN 2022 PNRR Project “RETINA: REmote sensing daTaNversion with multivariate functional modeling for essential climAte variables characterization”, funded by the European Union under the Italian National Recovery and Resilience Plan (NRRP) of NextGenerationEU, under the Italian Ministry of University and Research (Project Code: P20229SH29, CUP: J53D23015950001) and within the 2025 GNAMPA-INdAM Project “MultiPolExp: Polinomi di tipo esponenziale in assetto multidimensionale e multivoco”(CUP: E5324001950001).

References

- [1] Altomare, F., Campiti, M. (1994). Korovkin-type Approximation Theory and its applications. Walter de Gruyter & Co.
- [2] Angeloni, L., Costarelli, D. (2024). Approximation by Exponential-type Polynomials. *J. Math. Anal. Appl.*, 532, 127927.
- [3] Angeloni, L., Costarelli, D., Darielli, C. (2025). Approximation Processes by Multidimensional Bernstein-type Exponential Polynomials on the Hypercube. *R. Acad. Cienc. Exactas Fis. Nat. Serie A. Matemáticas*, 119:28.
- [4] Aral, A., Cárdenas-Morales, D., Garrancho, P. (2018). Bernstein-type Operators that Reproduce Exponential Functions. *J. Math. Inequal.*, 12, 861-872.
- [5] Morigi, S., Neamtu, M. (2000). Some Results for a Class of Generalized Polynomials. *Adv. Comput. Math.*, 12, 133-149.

3rd International Conference: Constructive Mathematical Analysis

Recent Results on the Weighted Approximation

Ali Aral

Department of Mathematics, Kirikkale University, Kirikkale, Türkiye

aliaral73@yahoo.com

key-words: Weighted approximation, weighted modulus of continuity, Korovkin and quantitative type theorems.

Abstract:

In this presentation, we express Korovkin and quantitative theorems under the weighted approximation. After giving the basic definitions and some of well-known spaces, we give the main theorems and their applications to linear positive operators. Moreover, we state a general operator containing well-known operators as an application of some theorems.

Acknowledgements: The author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Altomare, F., Campiti. M. (1994). Korovkin-type approximation theory and its applications. Walter de Gruyter, Berlin.
- [2] Aral, A. (2003). Approximation by Ibragimov-Gadjiev Operators in Polynomial Weighted Space. Proc. of IMM of NAS of Azerbaijan, XIX, 35-44.
- [3] Coskun, T. (1998). Some properties of Linear Positive Operators on the Spaces of Weight Functions. Commun. Fac. Sci. Univ. Ank. Series A1, (47), 175-181.
- [4] Coskun, T. (2000). Weighted Approximation of Continuous Functions by Sequences of Linear Positive Operators. Proc. Indian Acad.Sci. (Math. Sci.), 110(4), 357-362.
- [5] Gadjiev, A. D. (1974). The Convergence Problem for a Sequence of Positive Linear Operators on Unbounded Sets and Theorems Analogous to that of P. P. Korovkin. Sov. Math. Dokl., 15(5), 1433-1436.
- [6] Gadjiev, A. D. (1976). On P. P. Korovkin Type Theorems. Mathem. Zametki, 20, 781-786.
- [7] Gadjiev, A. D., Ispir, N. (1999). On a Sequence of Linear Positive Operators in Weighted Spaces. Proc. of IMM of Azerbaijan, XI(XIX), 45-56.
- [8] Gadjiev, A. D., Ibikli, E. (2002). On Non-existence of Korovkin's Theorem in the Space of L_p -locally Integrable Functions. Turkish J.Math., 26, 207-214.
- [9] Gadjiev, A. D., Aral, A. (2007). Weighted L_p -approximation with Positive Linear Operators on Unbounded Sets. Appl. Math. Lett., 20(10), 1046-1051.
- [10] Gadjiev, A. D., Aral, A. (2007). The Estimates of Approximation by Using a New Type of Weighted Modulus of Continuity. Comput. Math. with Appl., 54, 127-135.
- [11] Holhos, A. (2008). Quantitative Estimates for Positive Linear Operators in Weighted Space. Gen. Math., 16(4), 99-110.
- [12] Holhos, A. (2011). Uniform Weighted Approximation by Positive Linear Operators. Stud. Univ. Babes-Bolyai Math., 56(3), 135-146.

3rd International Conference: Constructive Mathematical Analysis

Approximation Results on Neural Network Operators of Convolution Type

Asiye Arif¹, Tuğba Yurdakadim¹

¹ Department of Mathematics, Bilecik Seyh Edebali University, Bilecik, Türkiye

arif_asiya@mail.ru, tugba.yurdakadim@bilecik.edu.tr

key-words: B -generalized logistic function, convolution type, rate of convergence, iterated approximation.

Abstract:

In the present paper, we introduce three neural network operators of convolution type activated by symmetrized, deformed and parametrized B -generalized logistic function. We deal with the approximation properties of these operators to the identity by using modulus of continuity. Furthermore, we show that our operators preserve global smoothness and consider the iterated versions of them. Here, we find it is worthy to mention that these operators play important roles in neural network approximation since most of the basic network models are activated by logistic functions.

References

- [1] Anastassiou, G. A. (1997). Rate of Convergence of Some Neural Network Operators to the Unit-univariate Case. *J. Math. Anal. Appl.* 212, 237–262.
- [2] Anastassiou, G. A. (2023). *Intelligent Computations: Parametrized, Deformed and General Neural Networks*, Springer, Heidelberg, New York.
- [3] Anastassiou, G. A. (2024). q -Deformed and λ -parametrized A -generalized Logistic Function Induced Banach Space Valued Multivariate Multi Layer Neural Network Approximations. *Stud. Univ. Babeş-Bolyai Math.*, 69(3), 587-612.
- [4] Anastassiou, G. A. (2024). Approximation by Parametrized Logistic Activated Convolution Type Operators, *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-mat.*, 118:138.
- [5] Cardaliaguet, P., Euvrard, G. (1992). Approximation of a Function and its Derivative with A Neural Network. *Neural Networks*, 5(2), 207-220.
- [6] Cybenko, G. (1989). Approximation by Superpositions of Sigmoidal Functions. *Math. Control Signals Syst.*, 2(4), 303-314.
- [7] Fausett, L. (1994). *Fundamentals of Neural Networks: Architectures, Algorithms, and Applications*, New Jersey: Prentice Hall.
- [8] Rosenblatt, F. (1957). *The Perceptron: A Perceiving and Recognizing Automaton*, Report 85-460-1, Cornell Aeronautical Laboratory, Buffalo, New York.

3rd International Conference: Constructive Mathematical Analysis

On Uniform Exhaustiveness and Korovkin-Type Theorem

Ali Arpacioğlu^{2,4}, Alper Erdem^{1,2,3} and Tuncay Tunc¹

¹ Department of Mathematics, Mersin University, Mersin, Türkiye

² Science Faculty, Mersin University, Mersin, Türkiye

³ Department of Mathematical and Statistical Sciences, University of Alberta, Edmonton, Canada

⁴ Human Resources Management Program, Tarsus University, Mersin, Türkiye

aliarpacioglu@tarsus.edu.tr, alpererdem@mersin.edu.tr, ttunc@mersin.edu.tr

key-words: Function sequences, exhaustiveness, Korovkin-type theorems.

Abstract:

This study considers a uniform version of exhaustiveness for function sequences and examined its connections with alpha convergence, uniform alpha convergence, and classical uniform convergence. A Korovkin-type approximation theorem is also established, showing that the classical conditions can be loosened under this new concept.

Let (X, d) and (Y, ρ) be metric spaces, (f_n) be a sequence of functions and f be a function from X to Y .

Definition 1. [2] The function sequence (f_n) is called *exhaustive* at $x_0 \in X$, if for every $\varepsilon > 0$ there exists $\delta = \delta(\varepsilon, x_0) > 0$ and $n_0 = n_0(\varepsilon, x_0) \in \mathbb{N}$ such that for all $x \in B_d(x_0, \delta)$ and all $n \geq n_0$ we have that $\rho(f_n(x), f_n(x_0)) < \varepsilon$.

Definition 2. [3] The function sequence (f_n) is called *uniform exhaustive* on X if for every $\varepsilon > 0$ there exists $n_0 = n_0(\varepsilon) \in \mathbb{N}$ and $\delta = \delta(\varepsilon) > 0$ such that for all $n \geq n_0$ and for all point $x, y \in X$ with $d(x, y) < \delta$ we have that $\rho(f_n(x), f_n(y)) < \varepsilon$.

Definition 3. [1] The function sequence (f_n) is called *uniform alpha convergent* to the function f on X if for every sequence (x_n) and (y_n) of points of X with the property $d(x_n, y_n) \rightarrow 0$, the sequence $\rho(f_n(x_n), f(y_n)) \rightarrow 0$.

Relationships with previously defined notions and classical uniform convergence are examined. Furthermore, a Korovkin-type approximation theorem is given within this new notion of exhaustiveness, emphasizing its significance in approximation theory. It is observed that the conditions in the classical Korovkin theorem can be loosened under this new notion.

References

- [1] Erdem, A., Tunc, T. (2025). On Cauchy Alpha and Uniform Alpha Convergence. Filomat, accepted.
- [2] Gregoriades, V., Papanastassiou, N. (2008). The Notion of Exhaustiveness and Ascoli-type Theorems. Topol. Appl., 155(10), 1111-1128.
- [3] Tunc, T., Erdem, A. (2024). Korovkin-type Theorems via Some Modes of Convergence. Filomat, 38(2), 523-530.

3rd International Conference: Constructive Mathematical Analysis

A Note on Modified Landau Type Operators

Didem Aydın Arı¹, Başar Yılmaz¹ and Ali Han Avcı¹

¹ Department of Mathematics, Kirikkale University, Kirikkale, Türkiye

didem@kku.edu.tr, basaryilmaz77@yahoo.com, avcialihan135@gmail.com

key-words: Modulus of continuity, Landau operator, weighted approximation.

Abstract:

In this study, we define modified Landau type operators. We investigate several convergence results concerning these operators. Specifically a quantitative approximation theorem and Voronovskaya type theorem are given using the modulus of continuity. Similar conclusions are also given in weighted spaces.

References

- [1] Altomare, F., Campiti, M. (1994). Korovkin-type Approximation Theory and Its Applications, Walter de Gruyter, New York.
- [2] Agratini, O., Aral, A. (2021) Approximation of Some Classes of Functions by Landau Type Operators, Results Math., 76:12.
- [3] Agratini, O., Gal, S. (2021). On Landau-Type Approximation Operators. Mediterr. J. Math., 18:64.
- [4] Gao, J. B. (1984). Approximation Properties of a Kind of Generalized Discrete Landau Operator (Chinese). J. Huazhong Univ. Sci. Technol., 12(5), 1-4.
- [5] Landau, E. (1908). Über die Approximation Einer Stetigen Funktion Durch Eine Ganze Rationale Funktion. Rend. Circ.Mat.Palermo, 25, 337-345.
- [6] Pendina, T.P. (1987). Iterations of Positive Linear Operators of Exponential Type and of Landau Polynomials (Russian). In: Geometric Problems of the Theory of Functions and Sets (Russian), pp. 105-111.

3rd International Conference: Constructive Mathematical Analysis

On Some Direct Estimates Results of the Rate Approximation by the Kantorovich Operator in Weighted Grand Lebesgue Spaces

Ismail Aydın¹, Ramazan Akgün²

¹ Department of Mathematics, Sinop University, Sinop, Türkiye

² Department of Mathematics, Balıkesir University, Balıkesir, Türkiye

iaaydin@sinop.edu.tr, rakgun@balikesir.edu.tr

key-words: Weighted grand Lebesgue spaces, Hardy-Littlewood maximal operator, Kantorovich operator.

Abstract:

In this study, we prove the boundedness of the Kantorovich operator K_n in weighted grand Lebesgue spaces $L_{\omega}^{p,\delta}(\Omega)$, where $\Omega \subset \mathbb{R}^N$ is a open bounded subset. Moreover, we establish two direct estimates by K -functionals of the rate of approximation in $L_{\omega}^{p,\delta}(\Omega)$. Finally, we extend the direct estimate inequality in classical Lebesgue spaces $L_p[0, 1]$ to $L_{\omega}^{p,\delta}$.

References

- [1] Aydın, I., Akgün, R. (2021). Weighted Variable Exponent Grand Lebesgue Spaces and Inequalities of Approximation. Hacet. J. Math. Stat., 50(1), 199-215.
- [2] Berens, H., Xu, Y. (1991). On Bernstein-Durmeier Polynomials with Jacobi Weights. In: C. K. Chui (ed.) Approximation Theory and Functional Analysis (College Station, TX, 1990), pp. 25-46. Academic Press, Boston.
- [3] Berens, H., Xu, Y. (1991). On Bernstein-Durmeier Polynomials with Jacobi Weights: the Case $p = 1$ and $p = \infty$. In: S. Baron, D. Leviatan (eds.) Approximation, Interpolation, and Summation, Jerusalem, Israel Mathematical Conference Proceedings, vol. 4, pp. 51-62, Weizmann Science Press of Israel.
- [4] Capone, C., Formica, M. R. and Giova, R. (2013). Grand Lebesgue Spaces with Respect to Measurable Functions. Nonlinear Anal., 85, 125–131.
- [5] Capone, C. (2022). Boundedness of Some Singular Integrals Operators in Weighted Generalized Grand Lebesgue Spaces, Ricerche Mat., 71, 109–120.
- [6] Draganov, B. R., Gadjev, I. (2024). Direct Estimates of the Rate of Approximation by the Kantorovich Operator in Variable Exponent Lebesgue spaces. Mediterr. J. Math., 21(112), 1-17.
- [7] Gonska, H., Zhou, X. L. (1995). The Strong Converse Inequality for Bernstein-Kantorovich Operators. Comput. Math. Appl., 30, 103-128.
- [8] Kokilashvili, V. (2010). Boundedness Criteria for Singular Integrals in Weighted Grand Lebesgue Spaces. J. Math. Sci., 170, 20-33.

3rd International Conference: Constructive Mathematical Analysis

On Some Properties and Compactness Criteria of Variable Exponent Fofana's Space

Ismail Aydın¹, Oznur Kulak²

¹Department of Mathematics, Sinop University, Sinop, Türkiye

²Department of Mathematics, Amasya University, Amasya, Türkiye

iaaydin@sinop.edu.tr, oznur.kulak@amasya.edu.tr

key-words: Amalgam spaces, Hardy-Littlewood maximal operator, variable exponent Fofana's spaces.

Abstract:

In functional analysis, the Kolmogorov-Riesz (or Fréchet-Kolmogorov) theorem gives a necessary and sufficient condition for a set of functions to be relatively compact in an L^p $[0, 1]$ space ($1 < p < \infty$) by Kolmogorov [4]. Compactness results in the spaces $L^p(\mathbb{R}^d)$ are quite important in existence of solutions and proofs for nonlinear partial differential equations. This theorem has been generalized to some function spaces, such as Tamarkin [7], Olsen and Holden[3], Rafeiro [6], Gorka and Macios [2], Pandey [5], Aydın and Unal [1]. In this regard, we recommend some new theorems for compactness criteria of bounded subsets in Variable Exponent Fofana's Spaces. Finally, we obtain also a generalization of the corresponding above studies.

References

- [1] Aydın, I., Unal, C. (2020). The Kolmogorov-Riesz Theorem and Some Compactness Criteria of Bounded Subsets in Weighted Variable Exponent Amalgam and Sobolev spaces. *Collect. Math.*, 71(3), 349-367.
- [2] Gorka, P., Macios, A. (2015). Almost Everything You Need to Know about Relatively Compact Sets in Variable Lebesgue Spaces. *J. Funct. Anal.*, 269(7), 1925-1949.
- [3] Hanche-Olsen, H., Holden, H. (2010). The Kolmogorov-Riesz Compactness Theorem. *Expo. Math.*, 28(4), 385-394.
- [4] Kolmogorov, A. N. (1931). Über kompakte der funktionenmengen bei der konvergenz im mittel. *Nach. Gess. Wiss. Göttingen Math.-Phys. Kl.*, 9, 60-63..
- [5] Pandey, S. S. (2003). Compactness in Wiener Amalgams on Locally Compact Groups. *Int. J. Math. Math. Sci.*, 2003(55), 3503-3517.
- [6] Rafeiro, H. (2009). Kolmogorov Compactness Criterion in Variable Exponent Lebesgue Spaces. *Proc. A. Razmadze Math. Inst.*, 150(2009), 105-113.
- [7] Tamarkin, J. D. (1933). On the Compactness of the Space L_p . *Bull. Amer. Math. Soc.*, 38(2)(1932), 79-84. *ach. Gess. Wiss. Göttingen Math.-Phys. Kl.*, 1933(1933), 167-170.

3rd International Conference: Constructive Mathematical Analysis

Calculation of Two-Center Coulomb Integrals over STOs in Nonlined-up Coordinate System

Erhan Akin^{1,2}, Arif Aykin¹

¹ Faculty of Science, Department of Physics, Selcuk University, Konya, Türkiye

² Institute of Science, Department of Physics, Selcuk University, Konya, Türkiye

eakin@selcuk.edu.tr

key-words: Slater-type orbitals, Coulomb integrals.

Abstract:

One of the most commonly used approaches in electronic structure calculations of many-electron systems is the Hartree-Fock-Roothaan (HFR) approach. According to this approach, the HFR equation must be solved to obtain the wave function (Slater determinant) of the molecule. One of the terms of the HFR equation is the Fock matrix, one is the kinetic energy of the electron and the interaction potential energy of the electron with the nucleus, and the other is the electron repulsion potential energy term resulting from the interaction of the electron with other electrons. The most important of the electron repulsion potential energy is the Coulomb potential energy term. In addition to these, the Coulomb potential energy must be calculated accurately in the solution of the Dirac equation (Bagci and Hoggan, 2015).

As can be understood from what we have said, it is very important to calculate the Coulomb potential energy accurately in order to obtain the wave function (Slater Determinant) of the quantum system using the HFR approach. In addition, in the calculation of the orbital energies of atoms or molecules, the most important term after the kinetic energy of the electron and the interaction potential energy with the nucleus is again the Coulomb potential energy term between electrons. In addition to what has been said, the Coulomb potential energy must also be calculated sensitively in the solution of the Dirac equation.

References

- [1] Akin, E. (2008). Evaluation of two-center electric field gradient integrals over STOs using elliptical coordinates. The European Physical Journal D, 49, 305-310.
- [2] Guseinov, I. I., Özmen, A., Atav, Ü., d Yüksel, H. (1995). Computation of Clebsch-Gordan and Gaunt coefficients using binomial coefficients. Journal of Computational Physics, 122(2), 343-347.
- [3] Guseinov, I. I., Atav, Ü., Özmen, A., Yüksel, H., Aliyeva, T. H. (1997). Calculation of rotation coefficients for overlap integrals over arbitrary atomic orbitals. Turkish Journal of Physics, 21(10), 1087-1092.
- [4] Guseinov, I., Mamedov, B., Rzaeva, A. (2002). Calculation of molecular integrals over Slater-type orbitals using recurrence relations for overlap integrals and basic one-center Coulomb integrals. Molecular modeling annual, 8, 145-149.
- [5] Özmen, A., Karakaş, A., Atav, Ü., Yakar, Y. (2003). Computation of two-center Coulomb integrals over Slater-type orbitals using elliptical coordinates. International journal of quantum chemistry, 91(1), 13-19.

3rd International Conference: Constructive Mathematical Analysis

Application of Wavelet Transformation in Analyzing Railway Track Profile Irregularities

Erdem Balci¹, Niyazi Özgür Bezgin², Mohamed Wehbi³

¹ Construction Department, Alapli Vocational School, Zonguldak Bülent Ecevit University, Zonguldak, Türkiye

² Civil Engineering Department, Engineering Faculty, Istanbul University-Cerrahpasa, Istanbul, Türkiye

³ Geobear Global, England, UK

erdembalci@outlook.com, ozgur.bezgin@iuc.edu.tr, mohamed.wehbi@geobear.com

key-words: Railway track analysis, wavelet transformation, rail profile irregularities.

Abstract:

Wavelet transformation provides a time-frequency analysis of non-stationary signals. It enables the detection of local events occurring within short time intervals. This capability is particularly critical for the analysis of railway track profile geometry [1], which often exhibits rapid and abrupt vertical irregularities over specific distances due to various mechanical factors. These factors include local voids in the supporting soil, rapid variations in track layer thickness, ballast wear and contamination, existence of rail-ends and turnouts, and the discontinuous support provided by sleepers. Inspection vehicles equipped with special measurement sensors can convert these geometric changes into a continuous signal.

Through multi-level wavelet decomposition, the dominant characteristics of the track profile signal can be extracted, along with detailed sub-signals of different frequencies and wavelengths [2]. These sub-signals represent the contribution of each underlying cause to the observed rail profile irregularities [3]. This study presents a case study where wavelet decomposition is applied to a track profile data collected from UK. The track profile is isolated into sub-parts and possible causes of the overall deformation is investigated independently. The findings can contribute to the developing effective maintenance strategies that enhance safety and extend the service life of railway tracks.

References

- [1] Zhiping, Z., Fei, L. and Yong, Z. (2010). Wavelet Analysis of Track Profile Irregularity for Beijing-Tianjin Intercity High Speed Railway on Bridge. In 2010 International Conference on Intelligent Computation Technology and Automation, 3, 1155-1158. IEEE.
- [2] Costa, M. A., Costa, J. N., Andrade, A. R. and Ambrósio, J. (2023). Combining Wavelet Analysis of Track Irregularities and Vehicle Dynamics Simulations to Assess Derailment Risks. Vehicle Syst. Dyn., 61(1), 150-176.
- [3] Balci, E. (2024). Demiryolu Hatları Üzerine, Hat ve Tren Tekerleklerindeki Düzensizlikler Nedeniyle Etki Eden Dinamik Darbe Kuvvetlerinin Analitik Olarak Çözümlemesi. PhD Thesis, Istanbul University-Cerrahpaşa.

3rd International Conference: Constructive Mathematical Analysis

Spectral Analysis of The Impulsive Sturm-Liouville Equation with Quadratic Spectral Parameters and Matrix Coefficients

Guler Basak Oznur

Department of Mathematics, Gazi University, Ankara, Türkiye

basakoznur@gazi.edu.tr

key-words: Scattering matrix, impulsive condition, eigenvalues, spectral singularities.

Abstract:

The aim of this study is to investigate some spectral and scattering properties of an impulsive Sturm-Liouville equation with matrix coefficient dependent on the spectral parameter in quadratic form. Firstly, we obtain the Jost solution and the Jost matrix. Then, we find the sets of eigenvalues and spectral singularities of our problem by using the resolvent operator and the Green matrix. Furthermore, we demonstrate that the impulsive differential equation has finite number of eigenvalues and spectral singularities with finite multiplicities under certain conditions. We also obtain the scattering matrix of our problem and examine the properties of the scattering matrix.

References

- [1] Agranovich, Z. S., Marchenko, V. A. (1963). The Inverse Problem of Scattering Theory. Gordon and Breach Science Publishers, New York-London.
- [2] Aktosun, T., Weder, R. (2020). Direct and Inverse Scattering For The Matrix Schrodinger Equation. Springer, Berlin.
- [3] Bairamov, E., Aygar, Y. and Cebesoy, S. (2019). Investigation of Spectrum and Scattering Function of Impulsive Matrix Difference Operators. Filomat, (33)5, 1301-1312.
- [4] Cebesoy, S., Bairamov, E. and Aygar, Y. (2023). Scattering Problems of Impulsive Schrodinger Equations with Matrix Coefficients. Ricerche di Matematica, 72(1), 399-415.
- [5] Dolzhenko, E. P. (1979). Boundary Value Uniqueness Theorems for Analytic Functions. Math. Notes, 26, 437-442.

3rd International Conference: Constructive Mathematical Analysis

Quantitative Estimates for Wavelet Type Extension of Generalized Kantorovich Operators

Gulen Bascanbaz-Tunca¹, Aysegül Erencin² and A. Feza Güvenilir¹

¹Department of Mathematics, Ankara University, Ankara, Türkiye

²Department of Mathematics, Bolu Abant İzzet Baysal University, Bolu, Türkiye

tunca@science.ankara.edu.tr, erencina@hotmail.com, guvenili@science.ankara.edu.tr

key-words: Generalized Kantorovich operators, wavelets, K -functional of Peetre, Bernstein-Markov type inequalities.

Abstract:

We consider a sequence of wavelet type modifications of univariate generalized Kantorovich operators depending on a positive real parameter given in [2].

In our talk; we present quantitative estimates for the rate of convergence of the operators in the spaces of continuous and p th power integrable functions defined on a suitable compact interval. The related estimates in both cases are expressed in terms of the modulus of continuity and the Peetre K -functional of the approximated function. Furthermore, we provide some inequalities such as Bernstein-Markov type for continuous functions and variation preservation property of the operators when the involved function is of bounded variation.

References

- [1] Agratini, O. (2005). Quantitative Approximations by Using Scaling Type Functions. Stud. Univ. Babes-Bolyai Math., 50(2), 3-13.
- [2] Altomare, F., Cappelletti Montano, M., Leonessa, V. and Raşa, I. (2017). A Generalization of Kantorovich Operators for Convex Compact Subsets. Banach J. Math. Anal., 11(3), 591-614.
- [3] Anastassiou, G. (2001). Quantitative Approximations. Chapman & Hall/CRC, Boca Raton.
- [4] Gonska, H. H., Zhou, D. X. (1995). Using Wavelets for Szász-type Operators. Rev. Anal. Numer. Theor. Approx., 24(1-2), 131-145.
- [5] Karsli, H. (2024). Asymptotic Properties and Quantitative Results of the Wavelet Type Bernstein Operators. Dolomites Res. Notes Approx., (17), 50-62.

3rd International Conference: Constructive Mathematical Analysis

Robust Estimation of Parameters in Simple Linear Regression Model

Demet Aydın¹, Omer Bayraktar²

¹ Department of Data Science and Analytics , Sinop University, Sinop, Türkiye

² Institute of Graduate Studies, Sinop University, Sinop, Türkiye

daydin@sinop.edu.tr, omerbayraktar55@gmail.com

key-words: Linear regression model, robust estimation methods, outliers, Monte-Carlo simulation.

Abstract:

This study investigates parameter estimation within a simple linear regression model, assuming that the error terms have a non-normal distribution. Robust estimation techniques, such as least absolute deviation, weighted least absolute deviation, and least median of squares, were employed to estimate the unknown model parameters. The performance of these estimators was evaluated using various comparison criteria. A Monte-Carlo simulation study was conducted on data sets both with and without outliers (in the y-direction). The results reveal that the least median of squares estimator outperforms the others in numerous scenarios, irrespective of the presence of outliers. Furthermore, the effectiveness of the proposed estimators is illustrated through an analysis of a data-set, which was intentionally altered with varying percentages of outliers.

References

- [1] Zeckhauser, R., Thompson, M. (1970). Linear Regression with Non-Normal Error Terms. Rev. Econ. Stat., 280-286.
- [2] Tiku, M. L., Islam, M. Q. and Selcuk, S. A. (2001). Non-normal Regression, Part II: Symmetric Distributions. Commun. Stat. - Theory Methods, 30, 1021-1045.
- [3] Tiku, M. L., Akkaya, A. D. (2004). Robust Estimation and Hypothesis Testing, New Age International (P) Limited Publishers, New Delhi, India.

3rd International Conference: Constructive Mathematical Analysis

Minimum Hellinger Distance Estimate for a Periodic Fractional Autoregressive Model of First-Order

Nesrine Benaklef¹, Karima Belaïde¹ and Djillali Seba¹

¹ Department of Mathematics, University of Bejaia/Applied Mathematics Laboratory, Bejaia, Algeria

nesrine.benaklef@univ-bejaia.dz, karima.belaide@univ-bejaia.dz, d.seba@esi-sba.dz.

key-words: Hellinger distance, estimation, periodic fractional autoregressive model.

Abstract:

In this paper, we develop an estimator for the parameter of a first-order periodic fractional autoregressive model following the minimum Hellinger distance approach. Then, we investigate two of its key properties, which are its consistency and limit distribution. A simulation study will demonstrate the most important findings of the study.

Model and notations:

A stochastic process $\{X_t, t \in \mathbb{Z}\}$ is said to be the fractional process of first-order $FAR(1)$ with a periodic coefficient denoted $PFAR_{ap}$ if it has the following representation

$$(1 - a_t L)^d X_t = \varepsilon_t \quad (1)$$

- $\theta_t = (a_t; d) \in \mathbb{R}^2$ is an unknown vector, such that for all $t \in \mathbb{Z}$, $\exists i = \{1, \dots, p\}$ and $M \in \mathbb{Z}$ such that $t = i + pM$ and so $a_t = a_{i+pM} = a_i$, here $p \geq 2$, we also suppose that $a_t \cdot d \neq 0$.

The minimum Hellinger distance estimator

In this paper, we will define an estimator of θ and let it be denoted by $\hat{\theta}_{M,n}$. This estimator minimizes the Hellinger distance (HD^2) between the theoretical probability density of η_M and a random probability density of the estimated η_M (referred to as $\hat{\eta}_M$) denoted by $f_\eta(\cdot)$ and $f_n(\cdot)$ respectively; $\hat{\theta}_{M,n}$ is hence named the **Minimum Hellinger Distance** (MHD) estimator, and it is defined in the following form

$$\hat{\theta}_{M,n} = \arg \min_{\theta \in \Theta} HD^2(f_n; f_\eta) \quad (2)$$

where $HD^2(f_n; f_\eta)$ is the L_2 -norm of the deviation between the square-roots of f_n and f_η .

References

- [1] AN, S., Bloomfield, P., and Pantula, S.G. (2004), 'Large sample property of the likelihood estimate for fractional arima models', *Observation, Theory And Modeling Of Atmospheric Variability: Selected Papers of Nanjing Institute of Meteorology Alumni in Commemoration of Professor Jijia Zhang*, 569–581.
- [2] Baillie, R.T., and Morana, C. (2012), 'Adaptive ARFIMA models with applications to inflation', *Economic Modelling*, 29, 2451–2459.
- [3] Beran, R. (1977), 'Minimum Hellinger distance estimates for parametric models', *The annals of Statistics*, 445–463.
- [4] Corduas, M. (2000), 'Preliminary estimation of ARFIMA models', *COMPSTAT: Proceedings in Computational Statistics 14th Symposium held in Utrecht, The Netherlands, 2000*, pp. 247–252. .

3rd International Conference: Constructive Mathematical Analysis

Solvability of a System Arising in Nonlinear Ultrasonic Heating

Imen Benabbas¹, Belkacem Said-Houari²

¹ AMNEDP Laboratory, Faculty of Mathematics, University of Sciences and Technology Houari Boumediene, Algiers, Algeria

² Department of Mathematics, University of Sharjah, United Arab Emirates

imen.benabbas@usthb.edu.dz, houari@sharjah.ac.ae

key-words: JMGT equation, nonlinear acoustics, Pennes bioheat equation.

Abstract:

We analyze a mathematical model of nonlinear ultrasonic heating based on the Jordan-Moore-Gibson-Thompson equation (JMGT) with temperature-dependent medium parameters coupled to the semilinear Pennes equation for the bioheat transfer. Precisely, we consider the model

$$\begin{cases} \tau p_{ttt} + p_{tt} - c^2(\bar{\theta})\Delta p - \delta(\bar{\theta})\Delta p_t = K(\bar{\theta})(p^2)_{tt}, & \text{in } \Omega \times (0, T), \\ \rho_a C_a \bar{\theta}_t - \kappa_a \Delta \bar{\theta} + \rho_b C_b W(\bar{\theta} - \theta_a) = \mathcal{Q}(\bar{\theta}, p_t), & \text{in } \Omega \times (0, T), \\ (p, p_t, p_{tt}, \bar{\theta})|_{t=0} = (p_0, p_1, p_2, \bar{\theta}_0), & \text{in } \Omega, \\ p|_{\partial\Omega} = 0, \quad \bar{\theta}|_{\partial\Omega} = \theta_a, & \end{cases} \quad (1)$$

where Ω is a bounded domain of $\mathbb{R}^d$, $d = 1, 2, 3$. The functions p and $\bar{\theta}$ denote the acoustic pressure and the temperature fluctuations, respectively, $c(\bar{\theta})$ is the speed of sound and $\delta(\bar{\theta})$ is the sound diffusivity. The parameter $\tau > 0$ denotes a positive constant accounting for the relaxation time and the function $K(\bar{\theta})$ is also allowed to depend on $\bar{\theta}$ and is given by $K(\bar{\theta}) = \frac{\beta_{\text{acous}}}{\rho c^2(\bar{\theta})}$, where ρ is the mass density and β_{acous} is the parameter of nonlinearity. As for the parameters in the Pennes equation, ρ_b, C_b are the density and specific heat capacity of blood, and W is the tissue's volumetric perfusion. We denote by ρ_a, C_a and κ_a , respectively, the ambient density, the ambient heat capacity and the thermal conductivity of the tissue. Moreover, θ_a stands for the ambient temperature and $\mathcal{Q}$ for the acoustic energy absorbed by the tissue.

Given the quasi-linearity of the pressure equation, we use a higher-order energy method together with a fixed-point argument to prove the local well-posedness of our model. This is achieved under the assumptions that the initial data are regular, small in a lower topology and that the final time is sufficiently short. In addition, making use of the uniformity of the derived estimates with respect to the time relaxation parameter τ , we determine the convergence rate of the solution of the JMGT-Pennes model to the solution of the Westervelt-Pennes model as $\tau \rightarrow 0$. For more details, see [1].

References

- [1] Benabbas, I., Said-Houari, B. (2024). Local Well-posedness of a Coupled Jordan-Moore-Gibson-Thompson-Pennes Model of Nonlinear Ultrasonic Heating. arXiv:2401.01709.
- [2] Bongarti, M., Charoenphon, S. and Lasiecka, I. (2021). Vanishing Relaxation Time Dynamics of the Jordan-Moore-Gibson-Thompson Equation Arising in Nonlinear Acoustics. J. Evol. Equ., 21, 3553-3584.
- [3] Kaltenbacher, B., Nikolić, V. (2022). Parabolic Approximation of Quasilinear Wave Equations with Applications in Nonlinear Acoustics. SIAM J. Appl. Math., 54, 1593-1622.

3rd International Conference: Constructive Mathematical Analysis

On Modular Equation and Related Distortion Functions

Barkat Ali Bhayo

School of Energy Systems, Lappeenranta-Lahti University of Technology, Lahti, Finland

barkat.bhayo@lut.fi

key-words: Special functions, metrics, elliptic integrals.

Abstract:

In this paper, we study distortion functions derived from the solutions of modular equation arising in number theory. We establish their monotonicity and convexity properties, and derive sharp bounds. Moreover, we discuss the role of distortion functions in the theory of quasiconformal mappings.

References

- [1] Anderson, G. D., Qiu, S.-L., Vamanamurthy, M. K. and Vuorinen M. (1997). Conformal Invariants, Inequalities and Quasiconformal Maps. J. Wiley.
- [2] Bhayo, B. A., Vuorinen, M. (2012). On Generalized Trigonometric Functions with Two Parameters. J. Approx. Theory, 164, 1415-1426.
- [3] Zhao, T-H., Bhayo, B. A. and Chu, Y-M Chu. (2022). Inequalities for Generalized Grotzsch Ring Function, Comput. Methods Funct. Theory, 22, 559-574.

3rd International Conference: Constructive Mathematical Analysis

Estimation of a Finite Population Mean Under Random Non-Response Using Improved Nadaraya-Watson Kernel Weights

Nelson Kiprono Bii¹, Christopher Ouma Onyango²

¹ Institute of Mathematical Sciences, Strathmore University, Nairobi, Kenya

² Department of Mathematics and Statistics, Kenyatta University, Nairobi, Kenya

nkiprono@strathmore.edu, onyango.christopher@ku.ac.ke

key-words: Mean squared error, bias, kernel regression, two-stage cluster sampling, confidence interval lengths.

Abstract:

Non-response is a potential source of errors in sample surveys. It introduces bias and large variance in the estimation of finite population parameters. Regression models have been recognized as one of the techniques of reducing bias and variance due to random non-response using auxiliary data. In this study, it is assumed that random non-response occurs in the survey variable in the second stage of cluster sampling assuming full auxiliary information is available throughout. Auxiliary information is used at the estimation stage via a regression model to address the problem of random non-response. In particular, auxiliary information is used via an improved Nadaraya-Watson kernel regression technique to compensate for random non-response. The asymptotic bias and mean squared error of the estimator proposed are derived. Besides, a simulation study conducted indicates that the proposed estimator has smaller values of the bias and smaller mean squared error values compared to existing estimators of finite population mean. The proposed estimator is also shown to have tighter confidence interval lengths at 95% coverage rate. The results obtained in this study are useful for instance in choosing efficient estimators of finite population mean in demographic sample surveys.

References

- [1] Breidt, F. J., et. al. (2000). Local Polynomial Regression Estimators in Survey Sampling. The Annals of Statistics, 28(4), 1026-1053, Institute of Mathematical Statistics.
- [2] Malec, P., et.al. (2014). Non-parametric Kernel Density Estimation Near the Boundary. Computational Statistics & Data Analysis, Volume 72, pages 57-76, Elsevier.
- [3] Simonoff, J. S. (2012). Smoothing Methods in Statistics. Springer Science & Business Media.

3rd International Conference: Constructive Mathematical Analysis

Spectrum of the Operator $H(r, s, t, u, v)$ Defined on the Sequence Spaces ℓ_p and bv_p

Mustafa Cemil Bisgin

Department of Mathematics, Recep Tayyip Erdoğan University, Rize, Türkiye

mustafa.bisgin@erdogan.edu.tr

key-words: Spectrum of an operator, resolvent set, perturbed operator.

Abstract:

In this work, we determine the fine spectrum of quintuple band matrix operator $H(r, s, t, u, v)$ which is defined on the sequence spaces ℓ_p and bv_p , where $1 < p < \infty$. The quintuple band matrix $H(r, s, t, u, v)$ generalizes the $D(r, 0, s, 0, t)$, Δ^4 , $Q(r, s, t, u)$, Δ^3 , $D(r, 0, 0, s)$, $B(r, s, t)$, Δ^2 , $B(r, s)$, Δ , right shift and Zweier matrices.

References

- [1] Akhmedov, A., Başar, F. (2006). The Fine Spectra of the Difference Operator Δ over the Sequence Space ℓ_p ($1 \leq p < \infty$). Demonstratio Math., 39, 586-595.
- [2] Akhmedov, A., Başar, F. (2007). On the Fine Spectra of the Difference Operator Δ over the Sequence Space bv_p ($1 \leq p < \infty$). Acta. Math. Appl. Sin. Engl. Ser., 10, 1757-1768.
- [3] Bilgiç, H., Furkan, H. (2008). On the Fine Spectrum of the Generalized Difference Operator $B(r, s)$ over the Sequence Spaces ℓ_p and bv_p , ($1 < p < \infty$). Nonlinear Anal., 68, 499-506.
- [4] Bisgin, M. C. (2022). On the Fine Spectrum of the Operator $Q(r, s, t, u)$ over ℓ_p and bv_p , $1 < p < \infty$. Rocky Mountain J. Math., 52(3), 777-791.
- [5] Furkan, H., Bilgiç and H., Başar, F. (2010). On the Fine Spectrum of the Operator $B(r, s, t)$ over the Sequence Spaces ℓ_p and bv_p , ($1 < p < \infty$). Comput. Math. Appl., 60, 2141-2152.

3rd International Conference: Constructive Mathematical Analysis

Recent Results on Sampling Kantorovich Operators of the Max-Product Type in Functional Spaces

Lorenzo Boccali^{1,2}, Danilo Costarelli² and Gianluca Vinti²

¹ Department of Mathematics and Computer Science “Ulisse Dini”, University of Florence, Florence, Italy

² Department of Mathematics and Computer Science, University of Perugia, Perugia, Italy

lorenzo.boccali@unifi.it, danilo.costarelli@unipg.it, gianluca.vinti@unipg.it

key-words: Max-product sampling Kantorovich operators, Orlicz spaces.

Abstract:

In recent years, the max-product approach, originally introduced by Bede, Coroianu and Gal (for a complete overview, see the monograph [2]), has been applied to several families of linear operators. The main advantage of max-product type operators—constructed using the maximum (or supremum in the case of infinite terms) instead of the usual sum (or series for infinite terms)—is that they generally offer sharper approximations, in terms of better orders of convergence, than their linear counterparts. Motivated by these considerations, in [5], the authors introduced and studied the max-product version of the Kantorovich sampling series [1] based on generalized kernels, obtaining convergence results in the space of uniformly continuous and bounded (non-negative) functions, as well as in the L^p -spaces, for $1 \leq p < +\infty$.

In the present talk, we will show a recent study on the approximation properties of the above non-linear (sub-additive) sampling operators in the more general setting of Orlicz spaces L^φ , generated by convex φ -functions. In recently published papers [3, 4], we investigated the problem of convergence, including the rate of approximation, for sequences of max-product sampling Kantorovich operators with respect to the general notion of modular convergence, typical of such functional spaces. In addition to obtaining a convergence result via a density approach, we also established a quantitative estimate for the order of approximation in the case of functions $f : \mathbb{R} \rightarrow \mathbb{R}_0^+$, $f \in L^\varphi(\mathbb{R})$, by exploiting the Orlicz-type modulus of smoothness $\omega(f, \delta)_\varphi$, with $\delta > 0$, defined in terms of the modular functional of the corresponding space. We will show that from this result it is possible to deduce, for non-negative functions belonging to suitable Lipschitz classes in Orlicz spaces, a qualitative estimate proving that the rate of convergence is affected by the regularity of both the kernel function and the function to be approximated.

Since, as it is well-known from the literature (see, e.g., [6]), these general spaces include a wide class of functional spaces—such as Lebesgue spaces, among others—the treatment in this general context offers the advantage of formulating a unifying theory for the convergence properties of the above operators.

References

- [1] Bardaro, C., Butzer, P. L., Stens, R. L. and Vinti, G., (2007). Kantorovich-type Generalized Sampling Series in the Setting of Orlicz Spaces. *Sampling Theory in Signal and Image Processing*, 6, 29-52.
- [2] Bede, B., Coroianu, L. and Gal, S. G. (2016). *Approximation by Max-product Type Operators*. Heidelberg: Springer.
- [3] Boccali, L., Costarelli, D. and Vinti, G. (2024). Convergence Results in Orlicz Spaces for Sequences of Max-product Sampling Kantorovich Operators. *J. Comput. Appl. Math.*, 449, 115957.
- [4] Boccali, L., Costarelli, D. and Vinti, G. (2024). Max-Product Sampling Kantorovich Operators: Quantitative Estimates in Functional Spaces. *Numer. Funct. Anal. Optim.*, 45(13-14), 667-685.
- [5] Coroianu, L., Costarelli, D., Gal, S. G. and Vinti, G. (2021). Approximation by Max-product Sampling Kantorovich Operators with Generalized Kernels. *Anal. Appl.*, 19(02), 219-244.
- [6] Musielak, J. (2006). *Orlicz Spaces and Modular Spaces*, Vol. 1034, Springer.

3rd International Conference: Constructive Mathematical Analysis

Approximation Properties of a New Generalization of Brass-Stancu Operators in L^p -Spaces

Murat Bodur

Department of Engineering Basic Sciences, Konya Technical University, Konya, Türkiye

mbodur@ktun.edu.tr

key-words: Brass-Stancu-Kantorovich-type operators, L^p -convergence, averaged modulus of smoothness.

Abstract:

In this presentation, we consider the operators defined by Brass [3] and Stancu [4]. Using the Kantorovich operators in [1], we obtain another Kantorovich-type modification of these operators as in [2]. We investigate the L^p -approximation properties of these new generalized operators. Specifically, the present study focuses on L^p -convergence and provides some upper estimates for the L^p -norm of the approximation error via first-order averaged modulus of smoothness and the first-order K -functional. In addition, we provide some numerical and graphical examples to demonstrate the rate of convergence of the constructed operators.

Acknowledgements: The author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Altomare, F., Cappelletti Montano, M., Leonessa, V. and Raşa, I. (2017). A Generalization of Kantorovich Operators for Convex Compact Subsets. *Banach J. Math. Anal.*, 11(3), 591-614.
- [2] Bodur, M., Bostancı, T. and Başcanbaz-Tunca, G. (2024). On Kantorovich Variant of Brass-Stancu operators. *Demonstr. Math.*, 57(1), 1-14.
- [3] Brass, H. (1971). Eine Verallgemeinerung der Bernsteinschen Operatoren. *Abh. Math. Sem. Univ. Hamburg*, 38, 111-122.
- [4] Stancu, D. D. (1983). Approximation of Functions by Means of a New Generalized Bernstein operator. *Calcolo*, 20(2), 211-229.

3rd International Conference: Constructive Mathematical Analysis

Cross-Approximation of Functions of 2 Variables

Petr Borodin

Moscow State University, Moscow, Russia

pborodin@inbox.ru

key-words: Approximation, bivariate function, cross-function.

Abstract:

Let $f(x, y)$ be a function defined on $[0, 1]^2$. The functions $f_{ab}(x, y) = f(a, y)f(x, b)$ ($a, b \in [0, 1]$) are called crosses generated by f . S. Mazur (1936) asked whether any continuous function can be uniformly approximated by linear combinations of its crosses with any accuracy. A. Grothendieck (1955) showed that this problem is in a certain way equivalent to the famous approximation problem. Since P. Enflo (1973) solved the approximation problem negatively, the answer to Mazur's question is 'no', and A. Davie (1975) constructed the corresponding example explicitly.

We present several conditions on a continuous function f that are sufficient for linear combinations of its crosses to approximate it with any accuracy.

References

- [1] Davie, A. M. (1975). The Banach Approximation Problem. J. Approx. Th., 13, 392-394.

3rd International Conference: Constructive Mathematical Analysis

Fixed Point Results for Generalized Cyclic $\alpha - f\pi\varpi$ -Contractions

Abderrahmane Boudraa

System Dynamics and Control Laboratory, Department of Mathematics and Informatics, Oum El Bouaghi University, Algeria

abderrahmane.boudraa09@gmail.com

key-words: b -metric-like space, fixed point, cyclic contractions.

Abstract:

In this work, we introduce generalized cyclic contractions in the context of metric-like space and prove some new fixed point results concerning these contractions. We present suitable examples to make our findings worth mentioning.

References

- [1] Taieb, H., Stojan, R. (2024). Banach and Caristi Type Theorems in b -metric-like Spaces .Facta Universitatis (ni s) Ser. Math. Inform., 39(4), 713–720. <https://doi.org/10.22190/FUMI231026047H>
- [2] Parvaneh, V., Younis, M., Savic, D. and Mohammadi, B. (2022). Some New Fixed Point Results for Generalized Cyclic $\alpha - f\pi\varpi$ -Contractions in Metric-like Spaces. J. Math. Ext., 16(7), 1-26.

3rd International Conference: Constructive Mathematical Analysis

Compactness Result in the Set of Measures

Gheorghe Bucur

Institute of Mathematics of the Romanian Academy, Bucharest, Romania

gheorghe.bucur@gmail.com

key-words: σ -algebras, measure, duality, compactness, weak topology, uniform integrability.

Abstract:

We consider a σ -algebra $\mathcal{B}$ on a space X and we denote by $\underline{\mathcal{B}}$ (resp. $\mathcal{M}$) the set of all real bounded and $\mathcal{B}$ -measurable functions on X (resp. the set of all finite real measures on $\mathcal{B}$).

On the real vector space $\underline{\mathcal{B}} \times \mathcal{M}$ we consider the natural duality

$$(f, \mu) \mapsto \int f d\mu. \quad (1)$$

We give a complete characterisation of the compact subsets of $\mathcal{M}$ with respect to the weak topology on $\mathcal{M}$ associated with the above duality.

We remember that the Dunford-Pettis theorem gives a partial answer namely: if μ_0 is a positive real measure and K is a subset of $L^1(\mu_0)$, then K is relatively compact with respect to the duality $(L^1(\mu_0), L^\infty(\mu_0))$ if K is uniformly integrable with respect to μ_0 .

References

- [1] Bucur, I., Paltineanu, G. (2020). Topics in Uniform Approximation of Continuous Functions. Birkhauser.
- [2] Bourbaki, N. (1952). Integration. Paris.
- [3] Dunford, N., Schwartz, J. (1958). Linear Operators, Part I. New-York.
- [4] Meyer, P.A. (1966). Probability and Potentials. Blaisdell Publishing Company.
- [5] Nicolescu, M. (1960). Analiză Matematică III. Ed. Tehnică București.

3rd International Conference: Constructive Mathematical Analysis

A Converse of Triangle Inequality: Applications

Ileana Bucur

Department of Mathematics and Computer Science, Technical University of Civil Engineering of Bucharest, Bucharest, Romania

bucurileana@yahoo.com

key-words: Abel convergence, Banach spaces, triangle inequality, radius of convergence, power series.

Abstract:

In any metric space (M, d) , giving three points x_1, x_2, x_3 such that

$$d(x_1, x_2) < d(x_1, x_3) \quad (1)$$

we have

$$1 \leq \frac{d(x_2, x_3)}{d(x_1, x_3) - d(x_1, x_2)}. \quad (2)$$

In some cases we have

$$\sup \left\{ \frac{d(x_2, x_3)}{d(x_1, x_3) - d(x_1, x_2)} \mid x_2 \in M \right\} < \infty, \quad (3)$$

for example in Hilbert (resp. Banach) spaces we have

$$\frac{d(x, x_3)}{d(x_1, x_3) - d(x_1, x)} < \frac{d(x_2, x_3)}{d(x_1, x_3) - d(x_1, x_2)}, \forall x \in co(x_1, x_2, x_3). \quad (4)$$

$$(\text{resp. } \frac{d(x, x_3)}{d(x_1, x_3) - d(x_1, x)} < \frac{\max(d(x_2, x_3), d(x_1, x_3))}{d(x_1, x_3) - d(x_1, x_2)}). \quad (5)$$

An open question arises here.

We use this result to study the behaviour of a holomorphic function at the boundary.

References

- [1] Boboc, N. (1968). Functii Complexe. Ed. Didactic Di Pedagogic, Bucharest.
- [2] Bucur, I. (2022). Compactness Results in the Set of Differentiable Functions. Mediterr. J. Math., 19(4).
- [3] Rudin, W. (1964). Principles of Math. Analysis. Mc. Graw-Hill, New-York.
- [4] Stoilow, S. (1954). Functii Complexe. Ed. Academiei, Bucharest.

3rd International Conference: Constructive Mathematical Analysis

On Von Neumann Approximation Theorem

Gavriil Paltineanu¹, Ileana Bucur¹

¹ Department of Mathematics and Computer Science, Technical University of Civil Engineering of Bucharest, Bucharest, Romania

gavriil.paltineanu@gmail.com, bucurileana@yahoo.com

key-words: Weierstrass density theorem, Bernoulli inequality, Uryson family, Von Neumann density.

Abstract:

In one of his article on logics Von Neumann stated, without proof, that a set $\mathcal{V}$ of continuous functions on $[0, 1]^n$ with values in $[0, 1]$, uniformly approximates any $[0, 1]$ -continuous function on $[0, 1]^n$ wherever $\mathcal{V}$ is closed to the multiplication operation, contains the projection functions and

$$1 - f \in \mathcal{V} \quad \forall f \in \mathcal{V}. \quad (1)$$

We develop some technics which may be useful not only to prove the Neumann result. We improve also the methods used by Jewett in this area.

References

- [1] Bishop, E. (1981). A generalization of the Stone Weierstrass theorem. Proc. Am. Math. Soc., 81.
- [2] Bucur, I, Paltineanu, G. (2017). Some Density Theorems in the Set of Continuous Functions with Values in the Unit Interval. Mediterr. J. Math., 14:44.
- [3] Bucur, I, Paltineanu, G. (2020). Topics in Approximation of Continuous functions. Frontiers in Mathematics, Birkhauser.
- [4] Jewett, R. I. (1963). A variation of Stone-Weierstrass Theorem. Proc. of Am. Math. Soc., 14(5), 690-693.

3rd International Conference: Constructive Mathematical Analysis

Some Novel Fixed Point Results in the Sense of Interpolative Metric Space

Abdurrahman Buyukkaya¹ and Mahpeyker Ozturk²

¹ Department of Mathematics, Karadeniz Technical University, Trabzon, Türkiye

² Department of Mathematics, Sakarya University, Sakarya, Türkiye

abdurrahman.giresun@hotmail.com, mahpeykero@sakarya.edu.tr

key-words: Fixed point, interpolative metric space, simulation functions, integral equations.

Abstract:

In this study, we investigate a novel fixed point theorem within the framework of interpolative metric spaces, an enriched generalization of classical metric structures. By extending the scope of contractive-type conditions in this broader setting, we unveil new existence and uniqueness results that contribute to the ongoing development of nonlinear analysis. The proposed approach not only refines previous fixed point results but also offers greater flexibility in handling mappings that elude standard metric assumptions. Furthermore, we present a pioneering application of our main theorem to a nonlinear integral equation, demonstrating both the theoretical depth and the practical relevance of our findings. This work opens a promising avenue for future research, particularly in areas where classical fixed point results fall short due to structural constraints of the underlying space.

References

- [1] Berzig, M. (2022). First Results in Suprametric Spaces with Applications. *Mediterr. J. Math.*, 19(5), 226.
- [2] Karapınar, E. (2023). An Open Discussion: Interpolative Metric Spaces. *Adv. Theory Nonlinear Anal. Appl.*, 7, 24-27.
- [3] Karapınar, E., Agarwal, R. P. (2025). Some Fixed Point Results on Interpolative Metric Spaces. *Nonlinear Analysis: Real World Applications*, 82, 104244.
- [4] Younis, M., Ahmad, H., Ozturk, M. and Singh, D. (2025). A Novel Approach to the Convergence Analysis of Chaotic Dynamics in Fractional Order Chua's Attractor Model Employing Fixed Points, *Alex. Eng. J.*, 110, 363-375.
- [5] Cho, S. H. (2022). Fixed Point Theorems for $\mathcal{L}_\varphi$ Contractions in Branciari Distance Spaces, *Axioms*, 11, 479.
- [6] Ozturk, M., Golkarmanesh, F., Buyukkaya, A. and Parvaneh, V. (2023). Generalized Almost Simulative $\hat{Z}_{\Psi^*}^\Theta$ -contraction Mappings in Modular b -metric Spaces. *J. Math. Ext.*, 17(2), 1-37.
- [7] Girgin, E., Buyukkaya, A., Kuru, N.K., Younis, M. and Ozturk, M. (2024). Analysis of Caputo-type Nonlinear Fractional Differential Equations and Their Ulam-Hyers Stability, *Fractal Fract.*, 8(10), 558.

3rd International Conference: Constructive Mathematical Analysis

Convergence in the Variation Seminorm by Generalized Kantorovich-Type Szász-Mirakyan Operators Constructed via Appell Polynomials

Gul Canbolat¹, Harun Karsli² and Fatma Tasdelen Yesildal¹

¹ Department of Mathematics, Ankara University, Ankara, Türkiye

² Department of Mathematics, Bolu Abant İzzet Baysal University, Bolu, Türkiye

gecanbolat@ankara.edu.tr, karsli_h@ibu.edu.tr, tasdelen@science.ankara.edu.tr

key-words: Linear positive operators, generalized Szász-Mirakyan operators, variation seminorm, variation detracting property, rate of convergence, convergence in variation seminorm, TV [I]-space, Appell polynomials.

Abstract:

The aim of this study is to investigate the variation detracting property and convergence in variation of the generalized Kantorovich type Szász mirakyan operators constructed via Appell polynomials in the space of functions of bounded variation. These problems are examined with respect to the variation seminorm. Additionally, the rate of convergence is analyzed in terms of total variation.

Acknowledgements: The first author is supported by Scientific and Technological Research Council of Turkey (TUBITAK) for the graduate scholarship.

References

- [1] Altomare, F. (2019). Iterates of Markov Operators and Constructive Approximation of Semigroups. *Constr. Math. Anal.*, 2(1), 22-39.
- [2] Aral, A. (2023). Weighted approximation: Korovkin and quantitative type theorems. *Modern Math. Methods*, 1(1), 1–21.
- [3] Karsli, H. (2018). Approximation Results for Urysohn Type Two Dimensional Nonlinear Bernstein Operators. *Constr. Math. Anal.*, 1(1), 45-57.

3rd International Conference: Constructive Mathematical Analysis

A Novel Inflated Mixture Distribution with Exponential and Unit Lindley Components

Erdem Cankut¹, Kadir Karakaya¹

¹Department of Statistics, Faculty of Science, Selcuk University, Konya, Türkiye

erdem.cankut@selcuk.edu.tr, kkarakaya@selcuk.edu.tr

key-words: Inflated model, mixture distribution, bootstrap, point estimation.

Abstract:

In practical applications, there is an increasing need for more flexible distribution models to capture diverse data characteristics better. In this study, a new flexible mixture distribution is proposed. The components of this mixture are the exponential distribution and the unit-Lindley distribution [1] transformed to the (a, b) interval. This new mixture distribution differs from existing mixture distributions in the literature. Specifically, it possesses a mixture structure that inflates the (a, b) interval. Some fundamental statistical properties of the proposed distribution are examined, and graphical presentations for the probability density function and hazard rate function for various parameter settings are also provided. The maximum likelihood estimation method is used for the point estimation of the unknown parameters, and confidence intervals for the parameters are obtained based on the asymptotic properties of the maximum likelihood estimation as well as the bootstrap approach. The performances of both the point and interval estimators are evaluated through a Monte Carlo simulation study. A real data analysis is conducted to demonstrate the practical applicability of the proposed model.

References

- [1] Mazucheli, J., Menezes, A. F. B. and Chakraborty, S. (2019). On the One Parameter Unit-Lindley Distribution and Its Associated Regression Model for Proportion Data. J. Appl. Stat., 46(4), 700-714.

3rd International Conference: Constructive Mathematical Analysis

Disjoint $\mathcal{F}$ -transitive Composition Operators on Orlicz Spaces

Chung-Chuan Chen

Department of Mathematics Education, National Taichung University of Education, Taichung, Taiwan

chungchuan@mail.ntcu.edu.tw

key-words: Disjoint $\mathcal{F}$ -transitivity, composition operator, Orlicz space.

Abstract:

In this talk, we will give sufficient and necessary conditions for some families of composition operators on the Orlicz space to be disjoint $\mathcal{F}$ -transitive, where $\mathcal{F}$ is a Furstenberg family of a set. By considering some specific Furstenberg families of the set of integers, some previous results are recovered. This is a joint work [1] with Prof. Bagheri Salec, Prof. Tabatabaie and Prof. Al-Shammari.

References

- [1] Bagheri Salec, A. R., Chen, C-C., Tabatabaie, S. M. and Al-Shammari, H. H. M. (2025). Disjoint $\mathcal{F}$ -transitive Composition Operators on Orlicz Spaces, Submitted.

3rd International Conference: Constructive Mathematical Analysis

Best Proximity Points for Cyclic Operators with Applications in Image Processing

Mihaela Ancuta Chira¹, Ticala Cristina¹

¹ Department of Mathematics and Computer Science, North University Centre at Baia Mare, Technical University of Cluj-Napoca, Baia Mare, Romania

mihaela.chira@mi.utcluj.ro, cristina.ticala@mi.utcluj.ro

key-words: Best proximity point, cyclic operator, image processing.

Abstract:

Our aim is to present recent results regarding best proximity point theory with respect to cyclic operators. Inspired by the a cyclic operator and we propose a new algorithm which gives a technique to stitch a pattern.

References

- [1] Dechboon, P., Khammahawong, K. (2023). Best Proximity Point for Generalized Cyclic Enriched Contractions. *Mathematical Methods in the Applied Sciences*, 47(3).
- [2] Kirk, W. A., Srinivasan, P. S. and Veeramani, P. (2003). Fixed Points for Mappings Satisfying Cyclical Contractive Conditions, *Fixed Point Theory*, 4, 79-89.
- [3] Sadiq Basha, S., Shahzad, N. and Jeyaraj, R. (2013). Best Proximity Points: Approximation and Optimization. *Optim. Lett.*, 7, 145-155

3rd International Conference: Constructive Mathematical Analysis

Completeness Theorems on the Boundary

Alberto Cialdea

Dipartimento di Scienze di Base e Applicate, University of Basilicata, Potenza, Italy

alberto.cialdea@unibas.it

key-words: Completeness theorems, elliptic equations of higher order, elasticity system, parabolic equations.

Abstract:

Roughly speaking, there are two different kinds of completeness theorems for systems of particular solutions of partial differential equations. Results of the first kind show that in a given norm we can approximate a solution of a partial differential equation in a domain by a sequence of particular solutions of the same equation. For example, if we have a holomorphic function f of one complex variable, we can ask when f can be approximated in some norms by polynomials or by rational functions. The classical theorems of Runge and Mergelyan are the main results in this direction. These problems have been widely studied and extended to general elliptic partial differential equations.

Results of a second kind are the so-called completeness theorems in the sense of Picone. They are much more sophisticated and refer not only to a PDE, but also to a certain BVP. Given a BVP in a domain Ω , the problem is to prove that the system obtained by applying the boundary operator of the BVP to particular solutions of the PDE is complete in an appropriate space on $\partial\Omega$.

The aim of the present talk is to discuss this second topic. In particular, we will focus on the completeness on the boundary of systems obtained by means of polynomial solutions. After a review of the previous results, we present the most recent ones. They concern elasticity system [1], higher order scalar elliptic equations in two variables [2] and parabolic equations [3].

References

- [1] Cialdea, A. (2024), New completeness theorems on the boundary in elasticity. *J. Math. Anal. Appl.*, **537**(2), 128292, 21.
- [2] Cialdea, A. and Lanzara, F. (2024), Completeness theorems related to BVPs satisfying the Lopatinskii condition for higher order elliptic equations. *Constr. Math. Anal.*, **7**, 129–141.
- [3] Cialdea, A. and Mare, C.S. (2025), Completeness theorems on the boundary for a parabolic equation, to appear.

3rd International Conference: Constructive Mathematical Analysis

On Three-Layer Conjugation Problem for the Heat Conduction Equation in a Rectangle

K. Kaliyeva¹, Yue Congcong¹

¹ Al-Farabi Kazakh National University, Almaty, Kazakhstan

kaliyeva.kulyash@kazhu.kz, youshan2439545521@gmail.com

key-words: Two-layer medium, temperature field, unsteady heat conduction, method of integral transformations, conjugate conditions.

Abstract:

The analytical theory of unsteady heat conduction in a rectangular three-layer plate finds wide application in solving important technical and technological problems expressing Fourier's and Fick's laws under the corresponding dependence of the internal energy of the medium on temperature. In the article the initial boundary value problem for the heat conduction equation in a rectangular plate with a given initial temperature distribution, boundary conditions and conjugation conditions on the rupture line of the rectangular plate are considered. Thermodynamic processes are accompanied by heat transfer between elements of structures in the working space of thermal plants and the surrounding space, in which heat is a quantitative measure of energy and between bodies there is heat exchange and regularities of the process of heat transfer are considered by the theory of heat exchange and heat transfer. The analytical solution of the problem of unsteady heat conduction in a three-layer medium is constructed by the method of integral transformations: Laplace on the time variable and by the finite Fourier transform on the coordinate variables. The study of initial boundary value problems for the heat conduction equation and the creation of analytical methods for solving such problems represent an important and rapidly developing area of mathematical physics. The solution of such problems can be used to calculate unsteady temperature fields and heat fluxes in two-layer sheet products, structures and buildings, in flat samples of products and apparatuses for which thermophysical parameters functionally depend on the temperature and the interface of the three-layer medium, and in addition, can be used for the calculation of unsteady temperature fields and heat fluxes in two-layer sheet products, structures and buildings, in flat samples of products and apparatuses for which thermophysical parameters functionally depend on the temperature and the interface of the three-layer medium, and in addition, can be used for the calculation of unsteady temperature fields and heat fluxes in the three-layer medium. The boundary condition of conjugation (of the fourth kind) corresponds to the heat exchange of the body surface with the environment according to the Fourier law of heat conduction or to the heat exchange of contacting solid bodies, when the temperature of the contacting surfaces is the same (ideal, perfect thermal contact), besides, the flowing of a solid body is determined by the flow of a liquid (or gas), the heat transfer from the liquid (gas) to the body surface in the immediate vicinity of the body surface occurs according to the law of heat conduction (molecular heat transfer), which has the following properties. The problem in this formulation is considered for the first time. The practical value of the obtained results is determined by the solution of specific initial boundary value problems in a three-layer medium for the heat conduction equation.

References

- [1] Kaliyeva, K. (2014). Energy Conservation Law in the Free Atmosphere. International Journal of Engineering and Innovative Technology (IJEIT), 3(11), 50–61.
- [2] Kaliyeva, K. (2015). Existence and Uniqueness of Weak solutions for the Navier-Stokes Equations. In *Transactions on Engineering Technologies*, Chapter 4., pp. 49–65.
- [3] Kaliyeva, K. (2015). Energy Conservation Law for the Turbulent Motion in the Free Atmosphere. In Handbook of Research on Computational Simulation and Modeling in Engineering, Chapter 3, pp. 105–138. IGI Global. ISBN13: 978-1-4666-8823-0.

3rd International Conference: Constructive Mathematical Analysis

Hermite-Type Sampling Operator

Rosario Corso

Department of Mathematics and Computer Science, University of Palermo, Palermo, Italy

rosario.corso02@unipa.it

key-words: Generalized sampling operator, samplings of derivatives.

Abstract:

The generalized sampling operator is able to approximate bounded continuous functions and it is modeled on the sampling expansion for band-limited functions given by the Whittaker-Kotel'nikov-Shannon theorem. During the decades, some variations of this classical theorem have been proposed. One of them (dating back to Jagerman and Fogel and, in a more general form, to Linden and Abramson) takes into consideration also the derivative samples for the reconstruction of band-limited functions, with a consequent benefit of a larger sampling rate compared to the Whittaker-Kotel'nikov-Shannon theorem. Motivated by this new reconstruction, we modify the generalized sampling operator including the samplings of derivatives up to a generic order to approximate non necessarily band-limited functions. The definition of the new operator is

$$(G_{n,w}f)(x) = \sum_{k \in \mathbb{Z}} \left(\sum_{j=0}^n \frac{1}{j!} f^{(j)}\left(\frac{k}{w}\right) \left(x - \frac{k}{w}\right)^j \right) \chi(wx - k), \quad x \in \mathbb{R},$$

(where χ is, as usually, a kernel) and we call it an *Hermite-type sampling operator* of order $n \in \mathbb{N}^+$. One of its main features is the faster order of approximation. Besides the convergence and its rate, we discuss well-posedness, regularity, simultaneous approximation and a Voronovskaya-type formula.

This talk is based on [1].

References

- [1] Corso, R. (2025). Generalized Sampling Operators with Derivative Samples. J. Math. Anal. Appl., 547(1).

3rd International Conference: Constructive Mathematical Analysis

Nonlinear Reduced Modeling for FitzHugh-Nagumo Equation: A Comparison of Kernel Functions

Yusuf Çakır¹, Murat Uzunca¹

¹Department of Mathematics, Sinop University, Sinop, Türkiye

ycakir@sinop.edu.tr, muzunca@sinop.edu.tr

key-words: Kernel principle component analysis, pattern formation, kernel functions.

Abstract:

Linear methods remain fundamental in data analysis due to their strong theoretical grounding and broad applicability. However, many real-world systems exhibit nonlinear behaviors that linear approaches cannot fully capture. This study presents a reduced-order modeling technique using Kernel Principal Component Analysis (KPCA), a nonlinear extension of traditional PCA, applied to solution data obtained from the discrete solution vectors of the FitzHugh-Nagumo (FHN) equation. Spatial and temporal discretizations are performed using the finite difference method and the semi-implicit Euler scheme, respectively. Multidimensional scaling (MDS) is employed for non-iterative reconstruction from reduced to full space. KPCA demonstrates the ability to preserve pattern behavior of the solutions of the FHN system, while enabling efficient low-dimensional approximations. Comparative analysis of various kernel functions reveals their substantial impact on model accuracy and efficiency, underscoring the importance of tailored kernel selection for capturing underlying data structures.

References

- [1] Karasozen, B., Uzunca, M. and Kucukseyhan, T. (2015). Structure preserving integration and model order reduction of skew-gradient reaction-diffusion systems. *Annals of Operations Research*, 258, 79–106.
- [2] Çakır, Y., Uzunca, M. (2024). Kernel principal component analysis for Allen-Cahn equations. *Mathematics*, 12, 3434.
- [3] Rathi, Y., Dambreville, S. and Tannenbaum, A. (2006). Statistical shape analysis using kernel PCA. *School of Electrical and Computer Engineering Georgia Institute of Technology*, 6064, 1-8.

3rd International Conference: Constructive Mathematical Analysis

Commutator of Sharp Maximal Operator on Total Morrey Spaces

Suleyman Celik¹, Hamit C. Calisgan² and Vagif S. Guliyev^{2,3,4,5}

¹ Department of Banking Finance and Insurance, Social Sciences Vocational School, Firat University, Elazig, Türkiye

² Department of Mathematics, Kirsehir Ahi Evran University, Kirsehir, Türkiye

³ Institute of Applied Mathematics, Baku State University, Baku, Azerbaijan

⁴ Institute of Mathematics and Mechanics, Baku, Azerbaijan

⁵ Azerbaijan University of Architecture and Construction, Baku, Azerbaijan

suleyman.celik@firat.edu.tr, hcaglarcalisgan@gmail.com, vagif@guliyev.com

key-words: Sharp maximal operator, commutator, total Morrey spaces.

Abstract:

In this talk we consider the commutator of the sharp maximal operator $[b, M^\sharp]$ on the total Morrey spaces $L_{p,\lambda,\mu}(\mathbb{R}^n)$. We obtain necessary and sufficient conditions for the boundedness of the commutator of the sharp maximal operator $[b, M^\sharp]$ on the total Morrey spaces $L_{p,\lambda,\mu}(\mathbb{R}^n)$ when b belongs to the bounded space of mean oscillations $BMO(\mathbb{R}^n)$, thereby obtaining some new characterizations for some subclasses of $BMO(\mathbb{R}^n)$, see [1, 2, 3].

For $f \in L^1_{\text{loc}}(\mathbb{R}^n)$, the sharp maximal operator $M^\sharp$ and the commutator of the sharp maximal operator $[b, M^\sharp]$ are defined by

$$M^\sharp f(x) = \sup_{r>0} |B(x,r)|^{-1} \int_{B(x,r)} |f(y) - f_{B(x,r)}| dy, \quad [b, M^\sharp]f(x) = b(x)M^\sharp f(x) - M^\sharp(bf)(x),$$

where $B(x,r)$ is the ball of radius r centered at $x \in \mathbb{R}^n$ and $|B(x,r)|$ denotes the Lebesgue measure of $B(x,r)$.

Morrey spaces, introduced by C. B. Morrey 1938, play important roles in the regularity theory of PDE, including heat equations and Navier-Stokes equations. In [1] Guliyev introduce a variant of Morrey spaces called total Morrey spaces $L_{p,\lambda,\mu}(\mathbb{R}^n)$, $0 < p < \infty$, $\lambda \in \mathbb{R}$ and $\mu \in \mathbb{R}$, give basic properties of the spaces $L_{p,\lambda,\mu}(\mathbb{R}^n)$ and study some embeddings into the Morrey space $L_{p,\lambda,\mu}(\mathbb{R}^n)$, see also [2, 3]. Necessary and sufficient conditions for the boundedness of the maximal commutator operator M_b and the commutator of the maximal operator $[b, M]$ on $L_{p,\lambda,\mu}(\mathbb{R}^n)$ were also given. Some new characterizations for some subclasses of $BMO(\mathbb{R}^n)$ were obtained.

The aim of this work is to give necessary and sufficient conditions for the boundedness of the commutator of sharp maximal operator $[b, M^\sharp]$ on $L_{p,\lambda,\mu}(\mathbb{R}^n)$ when b belongs to $BMO(\mathbb{R}^n)$ spaces. New characterizations of some subclasses of $BMO(\mathbb{R}^n)$ spaces are obtained, see [2, 3].

References

- [1] Guliyev, V.S. (2022). Maximal commutator and commutator of maximal function on total Morrey spaces. J. Math. Inequal., 16(4), 1509–1524.
- [2] Guliyev, V.S. (2024). Characterizations for the fractional maximal operator and its commutators on total Morrey spaces. Positivity 28(2), Article number 51.
- [3] Guliyev, V.S. (2025). Corrigendum to : Maximal commutator and commutator of maximal function on total Morrey spaces. J. Math. Inequal., 19(1), 305–306.

3rd International Conference: Constructive Mathematical Analysis

Commutator of Anisotropic Maximal Operator on Total Anisotropic Morrey Spaces

Suleyman Celik¹, Ali Akbulut² and Mehriban N. Omarova^{3,4}

¹Department of Banking Finance and Insurance, Social Sciences Vocational School, Firat University, Elazig, Türkiye

²Department of Mathematics, Kirsehir Ahi Evran University, Kirsehir, Türkiye

³Baku State University, Baku, Azerbaijan

⁴Institute of Mathematics and Mechanics, Baku, Azerbaijan

suleyman.celik@firat.edu.tr, aakbulut@ahievran.edu.tr, mehribanomarova@yahoo.com

key-words: Total anisotropic Morrey spaces, anisotropic maximal function, anisotropic fractional maximal function, commutator, anisotropic Lipschitz spaces.

Abstract:

In this talk we consider the commutators of the anisotropic maximal operator $[b, M^d]$ in total anisotropic Morrey spaces $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$ when b belongs to anisotropic Lipschitz spaces $\dot{\Lambda}_{\beta,d}(\mathbb{R}^n)$.

Morrey spaces, introduced by C. B. Morrey 1938, play important roles in the regularity theory of PDE, including heat equations and Navier-Stokes equations. In [3] Guliyev introduce a variant of Morrey spaces called total Morrey spaces $L_{p,\lambda,\mu}(\mathbb{R}^n)$, $0 < p < \infty$, $\lambda \in \mathbb{R}$ and $\mu \in \mathbb{R}$, see also [4]. In [1] the authors was consider the total anisotropic Morrey spaces $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$, give basic properties of the spaces $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$ and study some embeddings into the Morrey space $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$. Necessary and sufficient conditions for the boundedness of the anisotropic maximal commutator operator M_b^d and the commutator of the anisotropic maximal operator $[b, M^d]$ on $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$ were also given. Some new characterizations for some subclasses of $BMO(\mathbb{R}^n)$ were obtained.

The aim of this work is to give necessary and sufficient conditions for the boundedness of the commutator of anisotropic maximal operator $[b, M^d]$ on $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$ when b belongs to anisotropic Lipschitz spaces $\dot{\Lambda}_{\beta,d}(\mathbb{R}^n)$. New characterizations of some subclasses of anisotropic Lipschitz spaces $\dot{\Lambda}_{\beta,d}(\mathbb{R}^n)$ are obtained, see [2].

References

- [1] Abasova, G. A., Omarova, M. N. (2022). Commutator of Anisotropic Maximal Function with BMO Functions on Total Anisotropic Morrey Spaces. Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Mathematics, 43(1), 3-15.
- [2] Celik, S., Akbulut, A. and Omarova, M. N. (2025). Characterizations of Anisotropic Lipschitz Functions via the Commutators of Anisotropic Maximal Function in Total Anisotropic Morrey Spaces. Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Mathematics, 45(1), 1-13.
- [3] Guliyev, V. S. (2022). Maximal Commutator and Commutator of Maximal Function on Total Morrey Spaces. J. Math. Inequal., 16(4), 1509-1524.
- [4] Guliyev, V. S. (2025). Corrigendum to : Maximal Commutator and Commutator of Maximal Function on Total Morrey Spaces. J. Math. Inequal., 19(1), 305-306.

3rd International Conference: Constructive Mathematical Analysis

Approximation Theorem via Statistical ψ -convergence on $H_\omega(K)$

Selin Cinar

Department of Mathematics, Sinop University, Sinop, Türkiye

scinar@sinop.edu.tr

key-words: Positive linear operators, statistical convergence, the Bleimann, Butzer and Hahn operators.

Abstract:

In this study, a general Korovkin-type theorem is presented for sequences of positive linear operators from $H_\omega(K)$ to $C_B(K)$ where $K = [0, \infty)$. This theorem is proved using the concept of statistical ψ -convergence, which is a generalization of the concept of statistical convergence stated independently by Fast and Steinhauss ([2], [2]). Then, an example that satisfies the conditions of our main theorem will be presented. Finally, we calculate the corresponding rate of the statistical ψ -convergence of positive linear operators through the continuity modulus.

References

- [1] Fast, H. (1951). Sur la Convergence Statistique. In Colloquium mathematicae, 2(3-4), 241-244.
- [2] Steinhaus, H. (1951). Sur la Convergence Ordinaire et la Convergence Asymptotique. In Colloquium mathematicae, 2(1), 73-74.

3rd International Conference: Constructive Mathematical Analysis

Generalized k -Lauricella Functions

Yahya Çin¹, Nejla Özmen¹

¹Department of Mathematics, Duzce University, Duzce, Türkiye

yahyacin2525@gmail.com, nejlaozmen06@gmail.com

key-words: Gamma and Beta k -function, Pochhammer k -symbol, k -Appell functions, k -Lauricella functions, generalized k -Lauricella functions.

Abstract:

The main purpose of this study is to introduce k -analogue of the Lauricella functions and the generalized Lauricella functions. Firstly, definition of k -Lauricella functions are defined with the help Pochhammer k -symbol. Later Generalized k -Lauricella functions are explained. Then theorems and implementations are given.

References

- [1] Srivastava, H. M., Daoust, M. C. (1969). Certain Generalized Neumann Expansions Associated with the Kampé de Fériet Function. *Nederl. akad. Wetensch. Indag. Math.*, 31, 449-457.
- [2] Srivastava, H. M., Manocha, H. L. (1984). *Treatise on Generating Functions*. John Wiley & Sons, Inc.: New York, USA.
- [3] Diaz, R., Pariguan E. (2007). On Hypergeometric Functions and Pochhammer k -symbol. *Divulgaciones Matemáticas.*, 15, 179-192.
- [4] Mubeen, S. (2012). k -Analogue of Kummer's First Formula. *Journal of Inequalities and Special Functions*, 3(3), 41-44.
- [5] Özmen, N., Erkus-Duman, E. (2013). Some Results for a Family of Multivariable Polynomials. *AIP Conf. Proc.*, 1558, 1124-1127.
- [6] Mubben, S., Rehman, A. (2014). (n, k) -factorials. *Journal of Inequalities and Special Functions*, 5(3), 14-20.

3rd International Conference: Constructive Mathematical Analysis

A New Formula of Delta Laplace Transform for the Anti-causal System Defined on Isolated Time Scales

Mufit San¹, Huriye Dag¹

¹ Department of Mathematics, Cankiri Karatekin University, Cankiri, Türkiye

mufitsan@karatekin.edu.tr, huriyedg348@gmail.com

key-words: Anti-causal system, Laplace transform, time scales.

Abstract:

This talk aims to present recent findings related to the unilateral and bilateral Laplace transforms for anti-causal systems defined on isolated time scales. The foundational work by Ortigueira et al. [4] introduced the Laplace transform for anti-causal systems, which was later generalized for arbitrary time scales by Ortigueira et al. [5] and San and Ortigueira [6]. We will compare our results with previously published findings by the authors listed in the bibliography, emphasizing the relevance and implications of our research in the context of existing literature.

References

- [1] Bohner, M., Peterson, A. (2001). Dynamic Equations on Time Scales: An Introduction with Applications. Springer Science & Business Media.
- [2] Bohner, M., Guseinov, G. S. (2010). The Laplace Transform on Isolated Time Scales. Comput. Math. Appl., 60(6), 1536-1547.
- [3] Davis, J. M., Gravagne, I. A. and Marks, R. J. (2010). Bilateral Laplace Transforms on Time Scales: Convergence, Convolution, and the Characterization of Stationary Stochastic Time Series. Circuits Syst. Signal Process., 29(6), 1141-1165.
- [4] Ortigueira, M. D., Coito, F. J. and Trujillo, J. J. (2015). Discrete-time Differential Systems. Signal Processing, 107, 198-217.
- [5] Ortigueira, M. D., Torres, D. F. and Trujillo, J. J. (2016). Exponentials and Laplace Transforms on Nonuniform Time Scales. Commun. Nonlinear Sci. Numer. Simul., 39, 252-270.
- [6] San, M., Ortigueira, M. D. (2022). Unilateral Laplace Transforms on Time Scales. Mathematics, 10(23), 4552.

3rd International Conference: Constructive Mathematical Analysis

Construction of New Curves on n^{th} Degree Polynomial-Type Curves Using the Flc Frame

Süleyman Senyurt¹, Esra Damar²

¹ Department of Mathematics, Ordu University, Ordu, Türkiye

² Department of Motor Vehicles and Transportation Technologies, Vocational School of Technical Sciences, Hitit University, Çorum, Türkiye

ssenyurt@odu.edu.tr, esradamar@hitit.edu.tr

key-words: Pedal curves, Frenet like frame, polynomial curves.

Abstract:

In this study, first, the pedal curves were defined as the geometric locus of perpendicular projections onto the Frenet-like frame (Flc) vectors T , D_2 and D_1 of a given polynomial curve. Second, the Frenet elements of the new pedal curves were expressed in terms of Flc frame vectors T , D_2 and D_1 , such that the interrelations among these frames could be established. Next the curvature and torsion functions of each pedal curve were examined, separately. Then, by considering the Frenet elements of these pedal curves as position vectors new Smarandache curves were defined. Furthermore, for each Smarandache curve, the expressions of curvature and torsion functions based on Frenet-like frame were obtained. Finally, new type of curves were introduced and contributed to the curve family, and most importantly an applet with animation was created and shared on the freeware Geogebra program for anyone to define new pedal curves according to Flc frame, simultaneously.

References

- [1] Ali, A. T. (2010). Special Smarandache Curves in the Euclidean Space. International Journal of Mathematical Combinatorics, 2, 30-36.
- [2] Dede, M. (2022). Why Flc-Frame is Better than Frenet Frame on Polynomial Space Curves?. Mathematical Sciences and Applications E-Notes., 10, 190-198.
- [3] Canli, D. Senyurt, S. Kaya, F. E. and Grilli, L. (2024). The Pedal Curves Generated by Alternative Frame Vectors and Their Smarandache Curves. Symmetry, 16(8).
- [4] Li, Y., Pei, D. (2018). Pedal Curves of Frontals in the Euclidean Plane. Math Meth Appl Sci., 41(5), 1988-1997.

3rd International Conference: Constructive Mathematical Analysis

Local Laplacian Equienergetic Graphs

Cahit Dede¹, Ayse Dilek Maden¹

¹Department of Mathematics, Selçuk University, Konya, Türkiye

cahitdede@gmail.com, aysedilekmeden@selcuk.edu.tr

key-words: Graph energy, local Laplacian energy, local Laplacian equienergetic graphs.

Abstract:

Let G be a graph with vertex set $\{v_1, v_2, \dots, v_n\}$ and L be the *Laplacian matrix* of G . The eigenvalues of L are called the *Laplacian eigenvalues* of G . Then, *Laplacian energy* of G is defined as

$$LE(G) = \sum_{i=1}^n |\lambda_i(L) - \bar{d}|,$$

where $\bar{d} = \frac{\sum_{i=1}^n d_i}{n}$ see [3, 4]. It is our main concern in this paper to study the problem of the variation of the energy when a vertex is deleted, which was first studied in Espinal and Rada [2], and more recently in [1, 5]. Let $G^{(v)}$ be the graph obtained from G by deleting vertex v . The quantity

$$\mathcal{LE}_G(v) = LE(G) - LE(G^{(v)})$$

is the variation of the Laplacian energy in G when the vertex v is deleted, and called as the *local Laplacian energy of G at vertex v* . Then, the *local Laplacian energy of G* is defined as:

$$e_L(G) = \sum_{v \in V(G)} \mathcal{LE}_G(v).$$

In this study, we define and characterize local-equienergetic graphs whose local Laplacian energies remain invariant under specific structural perturbations. In particular, we show that the deletion of an edge can preserve local Laplacian energy under certain conditions. Our results contribute to the broader understanding of energy-based invariants in spectral graph theory and offer new insights into the resilience of graph structures under local modifications.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Dede, C., Popat, K. M. (2025). Locally Equienergetic Graphs. MATCH Commun. Math. Comput. Chem, 93, 759–766.
- [2] Espinal, C., Rada, J. (2024). Graph Energy Change due to Vertex Deletion. MATCH Commun. Math. Comput. Chem, 92, 89–103.
- [3] Gutman, I. (2001). The Energy of a Graph: Old and New results. In Anton Betten, Axel Kohnert, Reinhard Laue, and Alfred Wassermann, editors, Algebraic Combinatorics and Applications, pages 196–211, Springer Berlin Heidelberg.
- [4] Nikiforov, V. (2007). The Energy of Graphs and Matrices. J. Math. Anal., 326(2), 1472–1475.
- [5] Rakshith, B. R., Das, K. C. (2024). A Note on (local) Energy of a Graph. Comput. Appl. Math., 43(7), 399.

3rd International Conference: Constructive Mathematical Analysis

Approximation by Bivariate Modified Generalized Sampling Operators

Samet Degle¹, Metin Turgay¹, Tuncer Acar¹

¹Department of Mathematics, Selcuk University, Konya, Türkiye

smt.degle@gmail.com, metinturgay@yahoo.com, tunceracar@ymail.com

key-words: Generalized sampling series, rate of convergence, bivariate modulus of continuity.

Abstract:

In this study, we introduce a new family of bivariate generalized sampling operators by modifying the classical operators through a function ρ that satisfies the suitable conditions. We investigate the pointwise and uniform convergence of our newly defined operator. We give the rate of convergence of the family of operators via the classical bivariate modulus on continuity. We also obtain an asymptotic formula in the sense of voronovskaja.

Acknowledgements: The author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports. Authors has been supported by Scientific and Technological Research Council of Turkey (TUBITAK) under the Grant Number 1001-Project 123F123. The authors thank to TUBITAK for their supports.

References

- [1] Turgay, M., Acar, T., (2024). Approximation by Modified Generalized Sampling Series. *Mediterr. J. Math.*, 21:107.
- [2] Butzer, P. L. (2001). The Sampling Theorem of Signal Processing. In *Trends in Industrial and Applied Mathematics: Proceedings of the 1st International Conference on Industrial and Applied Mathematics of the Indian Subcontinent* (p. 23). Kluwer Academic Publishers.
- [3] Acar, T., Turgay, M. (2023). Approximation by Bivariate Generalized Sampling Series in Weighted Spaces of Functions. *Dolomites Res. Notes Approx.*, 16, 11-22.
- [4] Acar, T., Alagoz, O., Aral, A., Costarelli, D., Turgay, M. and Vinti, G. (2022). Convergence of Generalized Sampling Series in Weighted Spaces. *Demonstr. Math.*, 55(1), 153-162.

3rd International Conference: Constructive Mathematical Analysis

Result for Multivalued Contractions on Ultrametric Spaces

Ozlem Acar¹, Tugce Delen¹

¹Department of Mathematics, Faculty of Science, Selcuk University, Konya, Türkiye

acarozlem@gmail.com, tugcedelen@yahoo.com

key-words: Fixed point, F-contraction, ultrametric space.

Abstract:

Recently, fixed point theory is studied on ultrametric space. The present talk deals with new fixed point theorems by F-contraction especially rational type and integral type mappings. The special cases of the results which reduce the existence are presented and some examples are given illustrate our results.

References

- [1] Acar, Ö. (2022). Some Fixed Point Results on Ultrametric space. *Topol. Algebra Appl.*, 10(1), 227-232.
- [2] Acar, Ö., Delen, T. (2025). A Result for F-contraction on Ultrametric Spaces. Submitted.
- [3] Acar, Ö., Delen, T. and Özkapu, A. S. (2025). The New Fixed Point Theorem on Ultrametric Spaces. Accepted.
- [4] Banach, S. (1922). Sur les operations dans les ensembles abstraits et leur application aux equations integrales. *Fund. Math.*, 3, 133-181.
- [5] Brianciari, A. (2001). A Fixed Point Theorem for Mappings Satisfying a General Contractive Condition of Integral Type, *IJMMS*, 9(29), 531-536.
- [6] Gajić, L. (2001). On Ultrametric Space, *Novi Sad J. Math.*, 31(2), 69-71.
- [7] Gajić, L. (2002). A Multivalued Fixed Point Theorem in Ultrametric Spaces. *Mat. Vesnik*, 54(3-4), 89-91.
- [8] Ginişwamy, J., Maheshwari, P.G. (2017). Fixed Point Theorems Under F-Contraction In Ultrametric Space. *Adx. Fixed point Theory*, 7(1), 144-154.
- [9] Kirk, W. A., Shahzad, N. (2012). Some Fixed Point Results in Ultrametric Spaces. *Topology Appl.*, 159(15), 3327-3334.
- [10] Liu, Z., Li, X., Kang, S.M. and et al. (2011). Fixed Point Theorems for Mappings Satisfying Contractive Conditions of Integral Type and Applications, *Fixed Point Theory Appl.*, 2011, 64.
- [11] Liu, Z., Li, J. and Kang, S.M. (2013). Fixed Point Theorems of Contractive Mappings of Integral Type. *Fixed Point Theory Appl.*, 2013, 300.
- [12] Mamghaderi, H., Masiha, H. P. (2016). Some Fixed Point Theorems for G-nonexpansive Mappings on Ultrametric Spaces and Non-Archimedean Normed Spaces with a Graph. *J. Math. Appl.*, 39, 81-90.
- [13] Mamghaderi, H., Masiha, H. P. (2018). Fixed Point Theorem in Ultrametric Space, *Appl. Anal. Discrete Math.* 12(2), 336-346.
- [14] Pitchaimani, M., Kumar, D.R. (2017). On Nadler Type Results in Ultrametric Spaces with Application to Well-posedness, *Asian-Eur. J. Math.*, 10(4), 1750073, 15 pp.
- [15] Popa, V., (2008). A General Fixed Point Theorem for Mappings in Ultrametric Spaces. *Stud. Cercet. Ştiinţ. Ser. Mat. Univ. Bacău*, 18, 249-254.
- [16] Robert, A. M. (2000). A Course in p-adic Analysis, vol. 198 of Graduate Texts in Mathematics, Springer-Verlag, New York.
- [17] Van Roovij, A. C. M. (1978). Non Archimedean Functional Analysis Marcel Dekker, New York.
- [18] Wardowski, D. (2012). Fixed Points of a New Type of Contractive Mappings in Complete Metric Spaces, *Fixed Point Theory Appl.*, 94, 6 pp.

3rd International Conference: Constructive Mathematical Analysis

Stability to Perturbations of S -fractions with Applications

Marta Dmytryshyn¹, Volodymyr Hladun² and Roman Rusyn³

¹ West Ukrainian National University, Ternopil, Ukraine

² Lviv Polytechnic National University, Lviv, Ukraine

³ Vasyl Stefanyk Precarpathian National University, Ivano-Frankivsk, Ukraine

m.dmytryshyn@wunu.edu.ua, volodymyr.r.hladun@lpnu.ua, roman.rusyn@pnu.edu.ua

key-words: Stability to perturbations, continued fraction, convergence, relative errors.

Abstract:

This presentation is devoted to studying stability under perturbations for a class of functional continued fractions known as S -fractions (Stieltjes continued fractions) [4]. While these mathematical constructs are powerful tools in applied problems, their practical application requires a thorough understanding of their stability under perturbations. We establish formulas quantifying the relative errors of the approximants that result directly from perturbations applied to the elements of the continued fraction itself. Utilizing these formulas and leveraging techniques from the analytical theory of continued fractions, we derive sufficient conditions for the stability of these continued fractions under perturbations. A significant outcome of this research is the derivation of explicit bounds for the relative errors of the approximants, demonstrating how the perturbation error of a continued fraction's approximant depends on the magnitude of the perturbations and the values of the fraction's coefficients.

References

- [1] Jones, W. B., Thron, W. J. (1980). Continued Fractions: Analytic Theory and Applications, Addison-Wesley Pub. Co., Reading.

3rd International Conference: Constructive Mathematical Analysis

On the Approximation of the Lauricella-Saran's Hypergeometric Functions F_K and Their Ratios by Branched Continued Fractions

Roman Dmytryshyn¹, Vitaliy Goran¹

¹Vasyl Stefanyk Precarpathian National University, Ivano-Frankivsk, Ukraine

dmytryshynr@hotmail.com, vitaliy.goran.23@pnu.edu.ua

key-words: Hypergeometric function, branched continued fraction, approximation by rational functions, convergence, analytic continuation.

Abstract:

We consider the problem of approximating Lauricella-Saran's hypergeometric functions by a special family of functions – branched continued fractions. We establish domains of the analytical continuation of Lauricella-Saran's hypergeometric function F_K with real and complex parameters, using their branched continued fraction expansions whose elements are polynomials of three complex variables [1, 2]. To do this, we used the so-called extension theorem, which allows us to extend the domain of convergence of the branched continued fraction, which is already known for a small domain, to a larger domain, as well as the PC method (so called correspondence property) to prove that it is also the domain of analytical continuation. At the end, several numerical experiments are presented that illustrate the efficient approximation of special functions by branched continued fractions.

References

- [1] Antonova, T., Dmytryshyn, R., Goran, V. (2023). On the Analytic Continuation of Lauricella-Saran Hypergeometric Function $F_K(a_1, a_2, b_1, b_2; a_1, b_2, c_3; \mathbf{z})$. *Mathematics*, 11(21), 4487.
- [2] Dmytryshyn, R., Goran, V. (2024). On the Analytic Extension of Lauricella-Saran's Hypergeometric Function F_K to Symmetric Domains. *Symmetry*, 16(2), 220.

3rd International Conference: Constructive Mathematical Analysis

Approximation by Kantorovich Sampling Operators in Weighted Variable Exponent Lebesgue Spaces

Borislav R. Draganov^{1,2}

¹ Department of Mathematics and Informatics, Sofia University “St. Kliment Ohridski”, Sofia, Bulgaria

² Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, Sofia, Bulgaria

bdraganov@fmi.uni-sofia.bg

key-words: Kantorovich sampling operator, direct estimate, Voronovskaya-type estimate, strong converse inequality, saturation, modulus of smoothness, weighted variable exponent Lebesgue space.

Abstract:

Bardaro, Butzer, Stens and Vinti [1] introduced the Kantorovich-type sampling operators

$$(S_w^X f)(x) := \sum_{k \in \mathbb{Z}} \frac{w}{\theta_k} \int_{t_k/w}^{t_{k+1}/w} f(u) du \chi(wx - t_k), \quad x \in \mathbb{R}, \quad w > 0.$$

Here f is a locally Lebesgue integrable function, defined on $\mathbb{R}$, and $\chi : \mathbb{R} \rightarrow \mathbb{R}$ is the kernel of the sampling operator. The nodes $\{t_k\}_{k \in \mathbb{Z}}$ satisfy the assumptions $t_k < t_{k+1}$ and $\lim_{k \rightarrow \pm\infty} t_k = \pm\infty$, as, in addition, $\theta \leq \theta_k := t_{k+1} - t_k \leq \Theta$ for all $k \in \mathbb{Z}$ with some positive constants θ, Θ .

We will present a direct estimate of the norm of the error of S_w^X by a modulus of smoothness in certain variable exponent Lebesgue spaces with the weight

$$\rho_{\alpha, \beta}(x) := \begin{cases} |x|^{-\alpha}, & x < -1, \\ 1, & -1 \leq x \leq 1, \\ x^{-\beta}, & x > 1, \end{cases}$$

where $\alpha, \beta \geq 0$. The variable exponent is such that the Hardy-Littlewood maximal operator is bounded.

Further, we will complement the direct error estimate with a converse one, which enables us to establish the saturation property of this approximation process and describe its saturation class and its trivial class.

The results to be presented have recently appeared in [2].

Acknowledgements: This study is financed by the European Union-NextGenerationEU, through the National Recovery and Resilience Plan of the Republic of Bulgaria, project No BG-RRP-2.004-0008.

References

- [1] Bardaro, C., Butzer, P. L., Stens, R. L. and Vinti, G. (2007). Kantorovich-type Generalized Sampling Series in the Setting of Orlicz Spaces. *Sampl. Theory Signal Image Process.*, 6(1), 29-52.
- [2] Draganov, B. R. (2025). A Characterisation of the Rate of Approximation of Kantorovich Sampling Operators in Weighted Variable Exponent Lebesgue Spaces. *Dolomites Res. Notes Approx.*, 18(1), 72-90.

3rd International Conference: Constructive Mathematical Analysis

On λ_f –Statistical Convergence of Sequences of Sets

Basıl Elsibli¹, Mikail Et¹

¹Department of Mathematics, Fırat University , Elazığ, Türkiye

221121102@firat.edu.tr, mikail.et68@gmail.com

key-words: Modulus function, Wijsman statistical convergence, statistical convergence.

Abstract:

In this work, using an unbounded modulus function f we introduce and examine the concepts of Wijsman λ_f –statistical convergence and Wijsman strongly λ_f –convergence, and give some relations between Wijsman λ_f –statistical convergence and Wijsman strongly λ_f –convergence.

References

- [1] Aizpuru, A., Listà-García, M. C. and Rambla-Barreno, F. (2014). Density by Moduli and Statistical Convergence. Quaest. Math., 37, 525-530.
- [2] Bhardwaj, V. K., Dhawan, S. (2015). f -Statistical Convergence of Order α and Strong Cesàro Summability of Order α with respect to a Modulus. J. Inequal. Appl., 2015, 14 pp.
- [3] Bhardwaj, V. K., Dhawan, S. (2017). Density by Moduli and Wijsman Lacunary Statistical Convergence of Sequences of Sets. J. Inequal. Appl., 2017:25.
- [4] Fast, H. (1951). Sur la convergence statistique, Colloq. Math., 2, 241-244.
- [5] Fridy, J. A. (1985). On Statistical Convergence. Analysis, 5(4), 301-313.
- [6] Nakano, H. (1953). Concave Modulars. J. Math. Soc. Japan 5, 29-49.
- [7] Nuray F., Rhoades, B. (2012). Statistical Convergence of Sequences of Sets. Fasc. Math., 49, 87-99.
- [8] Šalát, T. (1980). On Statistically Convergent Sequences of Real Numbers. Math. Slovaca, 30(2), 139-150.

3rd International Conference: Constructive Mathematical Analysis

On the Least Squares and Weighted Least Squares Estimation Based on Ranked Set Sampling

Tenzile Erbayram¹, Yunus Akdoğan¹

¹Department of Statistics, Faculty of Science, Selcuk University, Konya, Türkiye

tenzile.erbayram@selcuk.edu.tr, yakdogan@selcuk.edu.tr

key-words: Rank set sampling, estimation, perfect ranking, Monte Carlo simulation order statistics.

Abstract:

Rank set sampling stands out as an effective and practical method for parameter estimation, especially in cases where complete and accurate measurement of observations is difficult or costly. In this study, the least squares and weighted least squares methods is conserved based on order statistics and used in estimating the parameters of the Poisson new XLindley distribution. An extensive Monte Carlo simulation study is performed to compare the proposed least squares and weighted least squares methods with classical estimation methods. The results of the simulation studies show that the proposed modification has lower mean squared errors compared to the estimation methods used in classical rank set sampling, especially in the case of small samples. Applications carried out on real data sets also support the theoretical findings. As a result, estimators based on ranked set sampling offer a strong alternative in terms of obtaining more robust and reliable results in statistical modeling and inference studies.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/ 2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Augusto Taconeli, C., Rodrigues de Lara, I. A. (2022). Discrete Weibull distribution: Different estimation methods under ranked set sampling and simple random sampling. *Journal of Statistical Computation and Simulation*, 92(8), 1740-1762.
- [2] Erbayram, T., Akdoğan, Y. and Chesneau, C. (2025, March 11). Stress-strength Reliability for the Discrete Poisson-Lindley Distribution based on the Rank Set Sample Method. *Bulletin of the Iranian Mathematical Society*. [Accepted for publication]

3rd International Conference: Constructive Mathematical Analysis

Trees Families with Algebraic Connectivity Greather than or Equal to $2(1 - \cos(\pi/9))$

Emrehan Erdem¹, Ayse Dilek Maden¹ and Masood Ur Rehman²

¹Department of Mathematics, Selçuk University, Konya, Türkiye

²School of Mathematics and Physics, Anqing Normal Universit, Anqing, China

ecanerdem@selcuk.edu.tr, aysedilekmeden@selcuk.edu.tr, masood@aqnu.edu.cn

key-words: Trees, Laplacian matrix, algebraic connectivity.

Abstract:

For a simple and undirected graph G with n vertices and adjacency matrix A . The Laplacian spectrum of G is set of eigenvalues $\lambda_n(G) \geq \lambda_{n-1}(G) \geq \dots \geq \lambda_2(G) \geq \lambda_1(G) = 0$ of the Laplacian matrix L of G , given by $L = D - A$, where D is the diagonal matrix of vertex degrees. The second smallest eigenvalue of the Laplacian matrix is called algebraic connectivity and is often denoted by $\alpha(G)$ in the literature.

Let T be a tree (a connected acyclic graph) on n vertices and $\alpha(T)$ be algebraic connectivity of T . Classifying trees according to their algebraic connectivity values has been an interesting area of study. Yuan et al. classified trees such that $\alpha(T) \geq 2 - \sqrt{3}$ [2], Wang and Tan classified trees such that $\alpha(T) \geq \frac{5-\sqrt{21}}{2}$ [3]. Later, Belay et al. showed tree classes such that $\alpha(T) \geq 2(1 - \cos(\pi/7))$ [1]. The lower bounds $2 - \sqrt{3}$, $\frac{5-\sqrt{21}}{2}$ and $2(1 - \cos(\pi/7))$ given for algebraic connectivity in [1, 2, 3] were characterized actually by Kirkland in [4]. In this study, we give new limit points in a similar way and find new tree classes with algebraic connectivity greater than or equal to $2(1 - \cos(\pi/9))$.

References

- [1] Belay, D. D. M., de Freitas, M. A. A. and da Silva, C. M. (2021). On Trees with Algebraic Connectivity Greater than or Equal to $2(1 - \cos(\frac{\pi}{7}))$. Computational and Applied Mathematics, 40, 215.
- [2] Yuan, X. Y., Shao, J. Y. and Zhang, L. (2008). The Six Classes of Trees with the Largest Algebraic Connectivity. Discrete Applied Mathematics, 156, 757-769.
- [3] Wang, X. K. and Tan, S. W. (2012). Ordering Trees by Algebraic Connectivity. Linear Algebra Applications, 436, 3684-3691.
- [4] Kirkland, S. (2003). A Note on Limit Points for Algebraic Connectivity. Linear Algebra and its Applications, 373, 5-11.
- [5] Kirkland, S., Neumann, M. and Shader, B. L. (1996). Characteristic Vertices of Weighted Trees via Perron Values. Linear and Multilinear Algebra, 40, 311-325.

3rd International Conference: Constructive Mathematical Analysis

Fuzzy Soft Element and Its Application to Decision-making

Buse Erdiñç¹, Kemal Taşköprü¹

¹ Department of Mathematics, Faculty of Science, Bilecik Şeyh Edebali University, Bilecik, Türkiye

5497789@ogrenci.bilecik.edu.tr, kemal.taskopru@bilecik.edu.tr

key-words: Fuzzy soft set, fuzzy soft element, decision-making.

Abstract:

This paper proposes the novel concept of fuzzy soft elements, thereby extending the traditional notion of fuzzy soft points through the more flexible assignment of alternatives to attributes. Consequently, this concept offers a new vantage point for understanding fuzzy soft set operations. Through this, after some of the properties of fuzzy soft set operations are given, a decision-making example is presented that demonstrates the practical application.

References

- [1] Alcantud, J. C. R. (2022). The Relationship Between Fuzzy Soft and Soft Topologies. *Int. J. Fuzzy Syst.*, 24, 1653-1668.
- [2] Das, S., Samanta, S. K. (2012). Soft Real Sets, Soft Real Numbers and Their Properties. *J. Fuzzy Math.*, 20(3), 551-576.
- [3] Maji, P. K., Biswas, R. and Roy, A. R. (2001). Fuzzy Soft Sets. *J. Fuzzy Math.*, 9(3), 589-602.
- [4] Majumdar, P., Samanta, S. (2010). Generalised Fuzzy Soft Sets. *Comput. Math. Appl.*, 59(4), 1425-1432.
- [5] MokoY, J. (2020). Powerset Theory of Fuzzy Soft Sets. *Int. J. Fuzzy Log. Intell. Syst.*, 20(4), 298-315.
- [6] Molodtsov, D. (1999). Soft Set Theory—First results. *Comput. Math. Appl.*, 37(4), 19-31.
- [7] Taşköprü, K., Altıntaş, İ. (2021). A New Approach for Soft Topology and Soft Function via Soft Element. *Math. Meth. Appl. Sci.*, 44(9), 7556-7570.
- [8] Taşköprü, K., Karaköse, E. (2023). A Soft Set Approach to Relations and Its Application to Decision Making. *Math. Sci. Appl. E-Notes*, 11(1), 1-13.
- [9] Taşköprü, K. (2023). Soft Order Topology and Graph Comparison based on Soft Order. *AIMS Math.*, 8(4), 9761-9781.
- [10] Varol, B. P., Aygünoglu, A. and Aygün, H. (2014). Neighborhood Structures of Fuzzy Soft Topological Spaces. *J. Intell. Fuzzy Syst.*, 27(4), 2127-2135.
- [11] Zadeh, L. A. (1965). Fuzzy Sets. *Inf. Control*, 8(3), 338-353.
- [12] Zimmermann, H.-J. (2011). *Fuzzy Set Theory and Its Applications*. Springer, New York.

3rd International Conference: Constructive Mathematical Analysis

New Fixed Point Theorem for Rational F-Contractions

Esref Erdogan¹, Ozlem Acar¹

¹Department of Mathematics, Faculty of Science, Selcuk University, Konya, Türkiye

esreferdogan@yahoo.com, acarozlem@ymail.com

key-words: Fixed point, rational type, metric space.

Abstract:

In [1], Wardowski introduced the concept of F -contraction, proving that every F -contraction mapping on a complete metric space has a unique fixed point, with consecutive approximations converging to this fixed point. For the function $F = \ln(x)$ on the interval $x \in (0, 1)$, a Banach contraction is an example of an F -contraction, which extends the main theorem of [1] to the Banach Fixed Point Theorem. There are many papers have been studied on this subject set [1, 2]. The most evident results are concentrated on applications of F -contraction to differential, integral equation, homotopy,... In this talk, we extend Wardowski's technique to present fixed point results for rational type F -contraction mappings in spaces equipped with two metrics.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Wardowski, D. (2012). Fixed Points of A New Type of Contractive Mappings in Complete Metric Spaces, Fixed Point Theory Appl. 2012, 1-6.
- [2] Acar, Ö., Altun, I. (2017). A Fixed Point Theorem for φF -Geraghty Contraction on Metric-like Spaces, Fasciculi Mathematici, 2017, 5-12.
- [3] Acar, Ö. (2023). Some Fixed Point Results with Integral Type $(H, \phi)F$ -contraction, Annals of the University of Craiova-Mathematics and Computer Science Series, 50(1), 53-59.

3rd International Conference: Constructive Mathematical Analysis

A Theoretical Approach to Superfluous Parameters in Constrained Multivariate Linear Models

Melek Eriş Büyükkaya¹, Nesrin Güler²

¹Department of Computer Sciences, Karadeniz Technical University, Trabzon, Türkiye

²Department of Econometrics, Sakarya University, Sakarya, Türkiye

melekeris@ktu.edu.tr, nesring@sakarya.edu.tr

key-words: BLUP, multivariate linear models, superfluous parameters.

Abstract:

In this study, we develop a comprehensive methodology for analyzing constrained multivariate linear models (CMLMs) with superfluous parameters. The proposed model is constructed by augmenting new regressors to a base constrained multivariate linear model. Our analysis considers both the original constrained model and its extended version with superfluous parameters. With these models, the constraint added to the models is written in a common matrix form in terms of unknown parameters and converted to their unconstrained equivalents, thus providing an easier to process analysis. We derive explicit analytical formulas for computing the best linear unbiased predictors (BLUPs) and best linear unbiased estimators (BLUEs) of all unknown parameter matrices by solving systems of linear equations. The properties and analytical expressions of these predictors and estimators are examined within the framework of both models. In particular, several novel and valuable properties of BLUPs are established. Additionally, we provide analytical solutions for special cases involving the general vector of all unknown parameter vectors. For methodologies and related models, see [1] and [2, 3], respectively.

References

- [1] Güler, N., Eriş Büyükkaya, M. (2023). Statistical Inference of a Stochastically Restricted Linear Mixed Model. AIMS Mathematics, 8(10), 24401-24417.
- [2] Güler, N., Eriş Büyükkaya, M. (2024). Further Remarks on Constrained Over-parameterized Linear Models. Stat. Paper, 65(2), 975-988.
- [3] Jiang, B., Tian, Y. (2024). Equivalent Analysis of Different Estimations Under A Multivariate General Linear Model. AIMS Mathematics, 9(9), 23544-23563.

3rd International Conference: Constructive Mathematical Analysis

$d(d^{-1})$ -Continuous Multi-valued Mappings and Some Fixed Point Results

Ismat Beg¹, Irem Eroglu²

¹Department of Mathematics and Statistical Sciences, Lahore School of Economics, Lahore, Pakistan

² Department of Mathematics, Ordu University, Ordu, Türkiye

iremeroglu@odu.edu.tr

key-words: Quasi-metric, fixed point, d -continuity, d^{-1} -continuity, multi-valued map.

Abstract:

Banach contraction theorem [1] has been examined and improved by many researcher due to its applicability to many fields. In [4], Nieto et al. obtained a new extension of Banach contraction mapping theorem to partially ordered sets. The concept of quasi-metric space was introduced by Wilson [6]. In [5], Schellekens showed that quasi-metric spaces are very suitable tools for analyzing the complexity analysis of algorithms. Recently, in [2] new fixed point results in partially ordered bicomplete quasi-metric spaces have been obtained. In this work, we introduce the concept of d -continuity and d^{-1} -continuity for multivalued mappings defined on a partially ordered quasi-metric spaces and we prove some fixed point theorems for multivalued maps. Moreover, we give an application of our results to homotopy theory.

References

- [1] Banach, S. (1992). Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. Fund. Math., 3, 133-181.
- [2] Beg, I., Eroglu, I. and Valero, O. (2024). Fixed Points on Partially Ordered Quasi-metric Spaces. Fixed Point Theory, 25(2), 473-494.
- [3] Beg, I., Eroglu, I. (2024). Fixed Point of Multivalued Maps on Bicomplete Partially Ordered Quasi-metric Spaces. Discussiones Mathematicae Differential Inclusions, Control and Optimization, 44, 167-184.
- [4] Nieto, J. J., Rodríguez-López, R. (2005). Contractive Mapping Theorems in Partially Ordered Sets and Applications to Ordinary Differential Equations. Order, 22, 223-239.
- [5] Schellekens, M. (1995). The Smyth Completion: A Common Foundation for Denotational Semantics and Complexity Analysis. Proc.MFPS11, Electronic Notes in Theoretical Computer Science 1, 211-232.
- [6] Wilson, W. A. (1931). On Quasi-Metric Spaces. American Journal of Mathematics, 53(3), 675-684.

3rd International Conference: Constructive Mathematical Analysis

Non-parametric Estimation for Doubly Geometric Process

Gultac Eroglu Inan

Department of Statistics, Ankara University, Ankara, Türkiye

geroglu@ankara.edu.tr

key-words: Geometric process, doubly geometric process, non-parametric estimation, least square method, inverse Gaussian distribution.

Abstract:

The geometric process (GP) is a widely used stochastic model in the reliability theory. Lam [1, 2] first introduced the model as a generalization of the renewal process (RP) and used the process as a model in the reliability context. Lam [5] introduced the GP and its applications. Although the GP is an important and widely used model in many areas, it has some limitations. Wu [4] proposed the doubly geometric process (DGP) as an alternative model that can overcome the limitations of the GP. The parameter estimation problem is very important for both GP and DGP. Lam [3] introduces a least square (LSE) method for nonparametric (NP) estimation problem in GP. In this study two NP methods proposed by Jasim and Al-Qazaz [1] for DGP which are (LSE) and log-LSE are presented. Additionally, modified moment (MM) estimators are obtained. The efficiencies of the maximum likelihood (ML) estimators are compared with (MM) estimators by an extensive simulation study.

References

- [1] Jasim, O. R., Qazaz, Q. N. N. A. (2022). Nonparametric Estimation in Doubly Geometric Stochastic Process. Int. J. Agricult .Stat. Sci., 17.
- [2] Lam, Y. (1988). Geometric Processes and Replacement Problem. Acta Math Appl Sin., 4 366-377.
- [3] Lam, Y. (1988). A Note on the Optimal Replacement Problem. Adv. Appl. Probab., 20 ,479-482.
- [4] Lam, Y. (1992). Nonparametric Inference for Geometric Process. Commun. Stat.Theory Methods, 21, 2083-2105.
- [5] Lam, Y. (2007). The Geometric Processes and Its Applications. World Scientific, Singapore.
- [6] Wu, S. (2018). Doubly Geometric Process and Applications. Journal of the Operational Research Society, 69(1), 66-67.

3rd International Conference: Constructive Mathematical Analysis

Parameter Estimation for Doubly Geometric Process with the Inverse Gaussian Distribution

Gultac Eroglu Inan

Department of Statistics, Ankara University, Ankara, Türkiye

geroglu@ankara.edu.tr

key-words: Geometric process, doubly geometric process, parametric estimation, maximum likelihood method, inverse Gaussian distribution.

Abstract:

The geometric process (GP) is a widely used stochastic model in the reliability theory. Lam [1, 2] first introduced the model as a generalization of the renewal process (RP) and used the process as a model in the reliability context. Lam [3] introduced the GP and its applications. Although the GP is an important and widely used model in many areas, it has some limitations. The two limitations can be given as follows: The GP cannot be used for non-monotone interarrival times whose distributions have varying shape parameters. The second limitation is that only logarithmic growth or explosive growth is allowed. Wu [4] proposed the doubly geometric process (DGP) as an alternative model that can overcome the limitations of the GP. The parameter estimation problem is very important for both GP and DGP. In this study, the parameter estimation problem for a DGP is considered under the assumption that the first inter-arrival time has inverse Gaussian (IG) distribution. The estimators of the parameters are obtained by maximum likelihood method. Additionally, the asymptotic unbiasedness and consistency of the estimators are investigated. The asymptotic distributions of the maximum likelihood (ML) estimators are obtained and some statistical properties such as unbiasedness and consistency of the estimators are investigated. It is shown in the Monte Carlo simulation study that the ML estimators are highly efficient estimators.

References

- [1] Lam, Y. (1988). Geometric Processes and Replacement Problem. Acta Math Appl Sin., 4, 366-377.
- [2] Lam, Y. (1988). A Note on the Optimal Replacement Problem. Adv. Appl. Probab., 20, 479-482.
- [3] Lam, Y. (2007). The Geometric Processes and its Applications. World Scientific, Singapore
- [4] Wu, S. (2018). Doubly Geometric Process and Applications. Journal of the Operational Research Society, 69 (1), 66-67.

3rd International Conference: Constructive Mathematical Analysis

Quasi-metric Spaces and Related Fixed Point Results

Andreea Fulga

Department of Mathematics and Computer Science, Transilvania University of Brasov, Brasov, Romania

afulga@unitbv.ro

key-words: Quasi-metric space, fixed point.

Abstract:

We present a series of further results, in different settings, examining the fixed point results for such spaces.

References

- [1] Fulga, A., Karapinar, E. and Petrusel, G. (2020). On Hybrid Contractions in the Context of Quasi-Metric Spaces. Mathematics, 8(5). <https://doi.org/10.3390/math8050675>
- [2] Zhou, M., Liu, X., Saleem, N., Fulga, A. and Ozgur, N. (2022). A New Study on the Fixed Point Sets of Proinov-Type Contractions via Rational Forms. Symmetry, 14(1). <https://doi.org/10.3390/sym14010093>.
- [3] Karapinar E., Fulga A., Rashid M. and Shahid L., Aydi H. (2019). Large Contractions on Quasi-Metric Spaces with an Application to Nonlinear Fractional Differential Equations. Mathematics, MDPI, 7(5), 1-11.

3rd International Conference: Constructive Mathematical Analysis

On the Two-layer Adjoint Problem for the Heat Conduction Equation in a Rectangle

K.Kaliyeva¹, Ajiaheer Gemingnuer¹

¹ Al-Farabi Kazakh National University, Almaty, Kazakhstan

Kaliyeva.kulyash@kazhu.kz, giminnur9@gmail.com

key-words: Two-layer medium, temperature field, unsteady heat conduction, method of integral transformations, conjugation conditions.

Abstract:

The analytical theory of unsteady heat conduction in a rectangular plate finds wide application in solving important technical and technological problems expressing Fourier's and Fick's laws at the corresponding dependence of the internal energy of the medium on temperature. In the article the initial boundary value problem for the equation of heat conduction in a rectangular plate with a given initial temperature distribution, boundary conditions and conjugation conditions on the break line of the rectangular plate are considered. Thermodynamic processes are accompanied by heat transfer between elements of structures in the working space of thermal plants and the surrounding space, in which heat is a quantitative measure of energy and between bodies there is heat exchange and regularities of the process of heat transfer are considered by the theory of heat exchange and heat transfer. The analytical solution of the problem of unsteady heat conduction in a two-layer medium is constructed by the method of integral transformations: Laplace on the time variable and finite Fourier transform on the coordinate variables. The study of initial boundary value problems for the heat conduction equation and the creation of analytical methods for solving such problems represent an important and rapidly developing area of mathematical physics. The solution of such problems can be used to calculate unsteady temperature fields and heat fluxes in two-layer sheet products, structures and buildings, in flat samples of products and apparatuses for which thermophysical parameters functionally depend on the temperature and the interface of the two-layer medium, and in addition can be used to calculate unsteady temperature fields and heat fluxes in two-layer sheet products, structures and buildings. The boundary condition of conjugation (of the fourth kind) corresponds to the heat exchange of the body surface with the surrounding medium according to the Fourier law of heat conduction or to the heat exchange of adjoining solid bodies, when the temperature of the adjoining surfaces is the same (ideal, perfect thermal contact), besides, the flowing of a solid body is determined by the flow of liquid (or gas), the heat transfer from liquid (gas) to the body surface in close proximity to the body surface occurs according to the law of heat conduction (molecular heat transfer), which has the following properties

References

- [1] Kaliyeva, K. (2014). Energy Conservation Law in the Free Stmosphere. Int. J. Eng. Innov. Technol. (IJEIT), 3(11), 50–61. Florida, USA. ISSN: 2277-3754.
- [2] Kaliyeva, K. (2014). Existence and Uniqueness of the Navier-Stokes Problem in Infinite Space. IAENG J. Eng. Comput. Sci., Proc. of Int. Conf. on Eng. and Comput. Sci., July 2–4, London, U.K. ISBN: 978-988-1952-37.
- [3] Kaliyeva, K. (2015). Chapter 3. Energy Conservation Law for the Turbulent Motion in the Free Atmosphere. In: Handbook of Research on Computational Simulation and Modeling in Engineering, IGI Global, pp. 105–138. ISBN: 978-1-4666-8823-0.

3rd International Conference: Constructive Mathematical Analysis

On Controlled Interpolative Metric Spaces

Ekber Girgin

Department of Engineering Fundamental Sciences, Sakarya University of Applied Sciences, Sakarya, Türkiye

ekbergirgin@subu.edu.tr

key-words: Graphical interpolative metric space, controlled metric space, Kannan contraction.

Abstract:

Recently, Karapinar [1] introduced a new generalized metric space (so called interpolative metric space). After that, Panda [2] generalized this concept which is called extended interpolative metric spaces. Finally, Girgin et.al. [3] defined the controlled interpolative metric spaces. This study introduces new type fixed point theorems using Kannan contraction in the novel framework of controlled interpolative metric spaces.

References

- [1] Karapinar, E. (2023). An Open Discussion: Interpolative Metric Spaces. Adv. Theory Nonlinear Anal. Appl., 7, 24-27.
- [2] Panda, S. K. (2025). Rossler attractor-based numerical solution of the fractal-fractional operator: A fixed point approach. Letters in Nonlinear Analysis and its Applications.
- [3] Girgin, E., Büyükkaya A. and Öztürk, M. (2025). New Fixed Point Approach in Controlled Interpolative Metric Spaces. Submitted.

3rd International Conference: Constructive Mathematical Analysis

Commutator of Sharp Maximal Operator on Lorentz Spaces

Suleyman Celik², Sezgin Girginol¹ and Ali Akbulut¹

¹Department of Mathematics, Kirsehir Ahi Evran University, Kirsehir, Türkiye

²Department of Banking Finance and Insurance, Social Sciences Vocational School, Firat University, Elazig, Türkiye

suleyman.celik@firat.edu.tr, sezgingirginol@hotmail.com, aakbulut@ahievran.edu.tr

key-words: Maximal operator, commutator, Lorentz space.

Abstract:

In this talk we consider the commutator of the sharp maximal operator $[b, M^\sharp]$ on the Lorentz spaces $L^{p,q}(\mathbb{R}^n)$. We obtain necessary and sufficient conditions for the boundedness of the commutator of the sharp maximal operator $[b, M^\sharp]$ on the Lorentz spaces $L^{p,q}(\mathbb{R}^n)$ when b belongs to the bounded space of mean oscillations $BMO(\mathbb{R}^n)$, thereby obtaining some new characterizations for some subclasses of $BMO(\mathbb{R}^n)$, see [1, 2].

For $f \in L^1_{loc}(\mathbb{R}^n)$, the maximal operator M and the commutator of the anisotropic sharp maximal operator $[b, M^\sharp]$ are defined by

$$M^\sharp f(x) = \sup_{r>0} |B(x,r)|^{-1} \int_{B(x,r)} |f(y) - f_{B(x,r)}| dy, \quad [b, M^\sharp]f(x) = b(x)M^\sharp f(x) - M^\sharp(bf)(x),$$

where $B(x,r)$ is the ball of radius r centered at $x \in \mathbb{R}^n$ and $|B(x,r)|$ denotes the Lebesgue measure of $B(x,r)$. Suppose that f is a measurable function on $\mathbb{R}^n$, then we define

$$f^*(t) = \inf\{s > 0 : d_f(s) \leq t\}, \quad d_f(s) := |\{x \in \mathbb{R}^n : |f(x)| > s\}|, \quad \forall s > 0.$$

The Lorentz space $L^{p,q} \equiv L^{p,q}(\mathbb{R}^n)$, $0 < p, q \leq \infty$ is the collection of all measurable functions f on $\mathbb{R}^n$ such the quantity

$$\|f\|_{L^{p,q}} := \|t^{\frac{1}{p} - \frac{1}{q}} f^*(t)\|_{L^q(0,\infty)}$$

is finite. Clearly $L^{p,p} \equiv L^p$ and $L^{p,\infty} \equiv WL^p$. The functional $\|\cdot\|_{L^{p,q}}$ is a norm if and only if either $1 \leq q \leq p$ or $p = q = \infty$.

Theorem 1 Let $p, q \in (1, \infty)$. The following assertions are equivalent:

(i) $b \in BMO(\mathbb{R}^n)$ and $b^- \in L^\infty(\mathbb{R}^n)$.

(ii) The operator $[b, M^\sharp]$ is bounded on $L^{p,q}(\mathbb{R}^n)$.

(iii) There exist a constant $C > 0$ such that $\sup_B \frac{\|(b(\cdot) - 2M^\sharp(b)(\cdot))\chi_B\|_{L^{p,q}(\mathbb{R}^n)}}{\|\chi_B\|_{L^{p,q}(\mathbb{R}^n)}} \leq C$.

(iv) There exist a constant $C > 0$ such that $\sup_B |B|^{-1} \|(b(\cdot) - 2M^\sharp(b)(\cdot))\chi_B\|_{L^1(\mathbb{R}^n)} \leq C$.

References

- [1] Akbulut, A., Isayev, F.A. and Serbetci, A. (2024). Anisotropic Maximal Commutator and Commutator of Anisotropic Maximal Operator on Lorentz Spaces. Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., Mathematics, 44(4), 5-12.
- [2] Guliyev, V. S. (2024). Maximal Commutator and Commutator of Maximal Operator on Lorentz Spaces. Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., Mathematics, 44(4), 43-49.

3rd International Conference: Constructive Mathematical Analysis

Comparison of Various Types of Inverse Soft Covering Upper Approximations

Zehra Güzel Ergül

Department of Mathematics, Kırşehir Ahi Evran University, Kırşehir, Türkiye

zguzel@ahievran.edu.tr

key-words: Inverse soft set, rough set, inverse soft rough set, decision making.

Abstract:

This study presents a comparison of various inverse soft covering upper approximations, focusing specifically on the differences between them. A key finding of our work is the introduction of a new type of inverse soft covering upper approximation which is a new approach in developing inverse soft covering based rough sets. We demonstrate that this new approximation is smaller than the three existing approximations described in the literature. Based on this result, we aim to manage the decision-making processes, especially for the uncertainty problems, in a more ideal way.

References

- [1] Çetkin, V., Aygünöğlu, A. and Aygün H. (2016). A New Approach in Handling Soft Decision Making Problems. J. Nonlinear Sci. Appl., 9, 231-239.
- [2] Demirtaş N., Hussain, S. and Dalkilic, O., (2020), New Approaches of Inverse Soft Rough Sets and Their Applications in a Decision Making Problem. J. Appl. Math. Informatics, 38(3-4), 335-349.
- [3] Feng, F., Li C., Davvaz, B. and Ali, I.M., (2010). Soft Sets Combined with Fuzzy Sets and Rough Sets: A Tentative Approach. Soft Computing, 14(9), 899-911.
- [4] Molodtsov, D., (1999). Soft Set Theory First Results. Comput. Math. Appl., 37, 19-31.
- [5] Pawlak, Z., Rough, Sets, (1982). International Journal of Computer and Information Sciences, 11(5), 341-356.
- [6] Zhu, W., Wang F., (2007), On Three Types of Covering-Based Rough Sets. IEEE Transactions on Knowledge and Data Engineering, 19(8), 1131-1143.

3rd International Conference: Constructive Mathematical Analysis

Generalizations of Certain Classes of Single-Valued Mappings to Multivalued Cases

Emirhan Hacıoglu

Department of Mathematics, Trakya University, Edirne, Türkiye

emirhanhacioglu@trakya.edu.tr

key-words: Fixed point, multivalued mappings, existence.

Abstract:

In this study, multivalued generalizations of certain classes of single-valued transformations defined on metric spaces are obtained. Building upon recently introduced concepts such as mappings contracting perimeters of triangles, new structural properties and fixed-point results for the multivalued counterparts are explored. Further examples and applications are also provided to illustrate the theoretical framework.

References

- [1] Popescu, O., Popescu, C. M. (2024). Periodic Points of Mappings Contracting Total Pairwise Distances. arXiv preprint, arXiv:2403.19488.
- [2] Popescu, O., Popescu, C. M. (2024). Fixed Point Theorem for Generalized Chatterjea Type Mappings. Acta Math. Hungar. 173, 500–509.
- [3] Petrov, E. (2023). Fixed Point Theorem for Mappings Contracting Perimeters of Triangles. arXiv preprint, arXiv:2308.01003.
- [4] Popescu, O., Popescu, C. M. (2024). Mappings Contracting triangles. arXiv:2403.19488v1.

3rd International Conference: Constructive Mathematical Analysis

On Groups and Vector Spaces with Easy Direct Limits

Emília Halušková¹, Małgorzata Jastrzebska²

¹Mathematical Institute, Slovak Academy of Sciences, Košice, Slovakia

²Siedlce University, Konarskiego 2, Siedlce, Poland

ehaluska@saske.sk, malgorzata.jastrzebska@uws.edu.pl

key-words: Algebra, direct limit, retract, group, vector space.

Abstract:

The direct limit construction belongs to basic tools used in Universal algebra, see e.g. [2]. Despite its frequent application, a systematic account of properties of this construction is rare in the literature, for a recent one see [1].

We will focus on direct limit families in which only one algebra occurs. Groups and vector spaces will be considered as algebras.

Let $\mathcal{A}$ be an algebra. We denote by $\underline{\mathbf{L}}\mathcal{A}$ the class of all isomorphic copies of direct limits which can be obtained from $\mathcal{A}$ and we denote by $\mathbf{R}\mathcal{A}$ the set of all retracts of $\mathcal{A}$. Then $\mathbf{R}\mathcal{A} \subseteq \underline{\mathbf{L}}\mathcal{A}$. We will say that $\mathcal{A}$ has *easy direct limits* if every algebra from $\underline{\mathbf{L}}\mathcal{A}$ is isomorphic to a retract of $\mathcal{A}$.

If $\mathcal{A}$ is finite, then $\mathcal{A}$ is an algebra with easy direct limits, cf.[3]. Infinite mono-unary algebras with easy direct limits are studied in [4].

We will present a mono-unary approach which helps to recognize algebras with easy direct limits. It works for an algebra whose some term operation is also an endomorphism. E.g., for multiplicative group of non-zero rational numbers we can use the term operation x^2 to see that this group is not with easy direct limits. Further, we show that

- every finite dimensional vector space has easy direct limits.
- the additive group of rational numbers has easy direct limit and additive groups of integers and real numbers do not.

Acknowledgements: The research is supported with the Grant VEGA 2/0104/24.

References

- [1] Gan, A., Muzammal, A. and Yang, Y. (2021). Limits of Quantum B-Algebras. *Mathematics*, 9(24), 12p.
- [2] Grätzer, G.(2008). *Universal Algebra*. 2nd printing of the 2nd ed. 1979. New York, Springer.
- [3] Halušková, E., Ploščica, M.(1999). On Direct Limits of Finite Algebras. *Contributions to General Algebra* 11, Heyn, Klagenfurt, 101-104.
- [4] Halušková, E.(2018). Monounary Algebras with Easy Direct Limits. *Miskolc Math. Notes*, 19(1), 291-302.

3rd International Conference: Constructive Mathematical Analysis

Solution of Split Equality Fixed Point Problem for Finite Family of Demigeneralized Mappings

Murtala Haruna Harbau¹, Sahura Badamasi Muhammad^{2,3}

¹Department of Science and Technology Education, Bayero University, Kano, Nigeria

²Department of Mathematical Sciences, Bayero University, Kano, Nigeria

³Department of Mathematics, Aliko Dangote University of Science and Technology, Wudil, Kano, Nigeria

murtalaharbau@yahoo.com, sahurabadamasi77@gmail.com

key-words: Split equality fixed point problem, demigeneralized mappings, weak convergence.

Abstract:

In this paper, we constructed and study an iterative algorithm for approximating solutions of split equality fixed point problem (SEFPP) for finite family of demigeneralized mappings in uniformly convex and uniformly smooth real Banach spaces. Weak convergence of the sequence generated by the algorithm was established. Strong convergence theorem was also proved under the assumption that the operators are demi-compact. Moreover, we give application of our result to split equality problem (SEP)

References

- [1] Chidume, C. E., Adamu, A. (2021). Solving Split Equality Fixed Point Problem for Quasi- ϕ -nonexpansive Mappings. Thai Journal of Mathematics, 19(4), 1699-1717.
- [2] Giang, D. M., Strodiot, J. J. and Nguyen, V. H. (2017). Strong Convergence of an Iterative Method for Solving the Multiple-set split Equality Fixed Point problem in a Real Hilbert Space. Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat., 111(4), 983-998.
- [3] Ugwunnadi, G. C., Jolaoso, L. O. and Okeke, C. C. (2022). A generalized Explicit Iterative Method for Solving Generalized Split Feasibility Problem and Fixed Point Problem in Real Banach Spaces. Symmetry 14:335.

3rd International Conference: Constructive Mathematical Analysis

Preserving Non-Negativity in Rational Approximation of Fractional Diffusion Problems

Stanislav Harizanov¹, Svetozar Margenov¹

¹Institute of Information and Communication Technologies, Bulgarian Academy of Sciences, Sofia, Bulgaria

sharizanov@parallel.bas.bg, margenov@parallel.bas.bg

key-words: Fractional diffusion, rational approximation, non-negativity preserving.

Abstract:

Maximum principles are among the basic qualitative properties of the elliptic or parabolic operators. In numerically solving PDEs, it is a natural requirement that these properties be preserved at the discrete level.

The presented analysis is dedicated to problems in a multidimensional domain Ω with general geometry. Let $\mathcal{A}$ be a self-adjoint elliptic operator of second order. Our study is focussed on the case of spectral fractional diffusion where $\mathcal{A}^\alpha = \mathcal{W}^T \mathcal{D}^\alpha \mathcal{W}$. Here $\mathcal{W}$ is the row-matrix of eigenfunctions and $\mathcal{D}$ is the diagonal matrix of eigenvalues of $\mathcal{A}$. Let the symmetric and positive definite matrix A be obtained after a finite difference or finite element discretization of $\mathcal{A}$.

We first discuss the BURA (*Best Uniform Rational Approximation*) method for $\alpha \in (0, 1)$ (sub-diffusion), which is introduced in [2]. At the discrete level, the method is defined in the form $A^{-\alpha} \approx \mu_1^{-\alpha} r_{\alpha,k}(\mu_1 A^{-1})$. Here $r_{\alpha,k}(z)$ is the best uniform rational approximation of degree (k, k) to z^α , $z \in [0, 1]$ and $\mu_1 > 0$ is the minimal eigenvalue of A . Following [1] we further develop sufficient conditions for positive approximation of the operator $A^{-\alpha}$. Next, we investigate the case of $\alpha > 1$ (super diffusion). Here, the conditions for non-negativity preservation require a more careful analysis on both BURA and BURA-based methods (see, e.g. [3]).

The last part of the talk is devoted to the analysis of parabolic problems. In time stepping, θ -methods are utilised with $\theta \in \{0, 0.5, 1\}$. For implicit schemes, we need BURA or a BURA-based approximation of A^α . It may be surprising to some readers that numerically multiplying a vector by A^α is a more complicated task than solving the corresponding fractional diffusion linear system. A significant part of the presented results concerns the dependence of the non-negativity conditions on the fractional power of the diffusion operator.

References

- [1] Harizanov, S., Margenov, S. (2018). Positive Approximations of the Inverse of Fractional Powers of SPD M-matrices. *Control Systems and Mathematical Methods in Economics*, 687, 147-163.
- [2] Harizanov, S., Lazarov, R., Margenov, S., Marinov, P. and Pasciak, J. (2020). Analysis of Numerical Methods for Spectral Fractional Elliptic Equations based on the Best Uniform Rational Approximation. *Journal of Computational Physics*, 408, 109285.
- [3] Slavchev, D., Harizanov, S. and Kosturski, N., (2024). Analysis on Computational Issues when Approximating Fractional Powers of Sparse SPD Matrices. *International Journal of Applied Mathematics*, 37(6), 699-711.

3rd International Conference: Constructive Mathematical Analysis

Anti-Periodic Boundary Value Problems for Singular Fractional p -Laplacian Equations

Mahir Hasansoy

Department of Mathematics, Istanbul Beykent University, Istanbul, Türkiye

mahirhasansoy@beykent.edu.tr

key-words: Anti-periodic boundary value problems, Caputo fractional differential equations, fractional p -Laplacian, fixed point theorem, global existence.

Abstract:

We study the following anti-periodic boundary value problems for Caputo fractional differential equations with a singular nonlinearity involving p -Laplacian operator ([1], [2], [3], [4], [5], [6] and [7]).

$$\begin{cases} D_*^\beta \phi_p(D_*^\alpha x(t)) = t^{-\gamma} f(t, x(t)), \\ x(0) = -x(1), \quad D_*^\alpha x(0) = -D_*^\alpha x(1), \\ x \in \text{Dom}(D_*^\alpha), \quad \phi_p(D_*^\alpha x) \in \text{Dom}(D_*^\beta), \quad D_*^\alpha x \in C[0, 1], \end{cases} \quad (1)$$

where $f \in C([0, 1] \times \mathbb{R})$, $0 \leq \gamma < \alpha$, $\beta \leq 1$, $\phi_p(t) = |t|^{p-2}t$, $p > 1$. Our main goal is to prove the existence of a solution to the problem (0.1). We first, show that this problem is equivalent to a class of Volterra integral equations.

Theorem 1 Let $f \in C([0, 1] \times \mathbb{R})$ and $0 \leq \gamma < \alpha$, $\beta \leq 1$. A function $x \in C[0, 1]$ is a solution of the anti-periodic boundary value problem (0.1) if and only if $x \in C[0, 1]$ satisfies the Volterra integral equation

$$x(t) = I^\alpha \phi_q [I^\beta (t^{-\gamma} f(t, x(t))) + L_f x(t)] + M_f x(t). \quad (2)$$

where $L_f x(t) = -\frac{1}{2} I^\beta (t^{-\gamma} f(t, x(t)))|_{t=1}$ and $M_f x(t) = -\frac{1}{2} I^\alpha \phi_q [I^\beta (t^{-\gamma} f(t, x(t))) + L_f x(t)]|_{t=1}$.

The main result regarding the existence of the solution is given in the following theorem.

Theorem 2 Let $f \in C([0, 1] \times \mathbb{R})$ and there exist nonnegative functions $a, b \in C[0, 1]$ such that $|f(t, u)| \leq a(t) + b(t)|u|^{p-1}$, for all $t \in [0, 1]$ and $u \in \mathbb{R}$. If $\frac{3^p}{2^p \Gamma(\alpha+1)^{p-1}} \frac{\Gamma(1-\gamma)}{\Gamma(1-\gamma+\beta)} \|b\| < 1$ then the Volterra integral equation (0.2) has a solution in $C[0, 1]$.

Finally, we give an example as an application of Theorem 0.2.

Example 0.1 If $0 < b < \frac{1}{14}$ then the following problem has a solution in $C[0, 1]$.

$$\begin{aligned} D_*^{\frac{3}{4}} \phi_3(D_*^{\frac{1}{2}} x(t)) &= t^{-1/3} (1 + bx^2(t)), \\ x(0) &= -x(1), \quad D_*^\alpha x(0) = -D_*^\alpha x(1). \end{aligned}$$

References

- [1] Chen, T., Liu, W. (2012). An Anti-periodic Boundary Value Problem for the Fractional Differential Equation with a p -Laplacian Operator. Applied Mathematics Letters, 25, 1671-1675.
- [2] Chen, T., Liu, W. and Yang, C. (2010). Antiperiodic Solutions for Lienard-type Differential Equation with p -Laplacian Operator, Bound. Value Probl., 2010, 1-12.
- [3] Chen, Y., Nieto, J. J. and O'Regan, D. (2007). Anti-periodic Solutions for Fully Nonlinear First-order Differential Equations. Math. Comput. Model., 46, 1183-1190.
- [4] Fazli, H., Nieto, J. J. (2018). Fractional Langevin Equation with Anti-periodic Boundary Conditions. Chaos, Solitons and Fractals, 114, 332-337.
- [5] Faranco, D., Nieto, J. J. and O'Regan, D. (2003). Anti-periodic Boundary Value Problem for Nonlinear First Order Ordinary Differential Equations. J. Math. Inequal. Appl., 6, 477-485.
- [6] Hasanov, M. (2024). Initial Value Problems for Fractional p -Laplacian Equations with Singularity. AIMS Mathematics, 9, 12800-12813.
- [7] Webb, J. R. L. (2019). Initial Value Problems for Caputo Fractional Equations with Singular Nonlinearities. Electronic Journal of Differential Equations. 2019(117), pp. 1-32.

3rd International Conference: Constructive Mathematical Analysis

Squared Basis Operators Related to Modified Bessel Functions

Monika Herzog

Department of Applied Mathematics, Cracow University of Technology, Cracow, Poland

mherzog@pk.edu.pl

key-words: Squared basis, Bessel function, Voronovskaya-type results.

Abstract:

In our talk, we propose considering approximation properties of positive linear operators related to the modified Bessel function of the first kind. We focus on these operators because the operators related to the modified Bessel function of order 0 come in the King-type operators associated with the squared Szász-Mirakyan basis.

Similar to, for example, the case of Bernstein operators, it can be proved that the error of approximation by our operators is smaller than the classical Szász-Mirakyan ones.

References

- [1] Acu, A.M., Heilmann, M., Raşa, I. and Seserman, A. (2023). Poisson Approximation to the Binomial Distribution: Extensions to the Convergence of Positive Operators. *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.*, 117, 1-13.
- [2] Gupta, V., Acu, A. M. (2019). On Difference of Operators with Different Basis Functions. *Filomat*, 33, 3023-3034.
- [3] Herzog, M. (2003). Approximation of Functions from Exponential Weight Spaces by Operators of Szász-Mirakyan Type. *Commentat. Math.*, 43(1), 77-94.
- [4] Holhoş, A. (2020). King-type Operators related to Squared Szász-Mirakyan basis. *Studia Univ. "Babeş-Bolyai" Math.*, 65(2), 279-290.
- [5] King, J.P. (2003). Positive Linear Operators which Preserve x^2 . *Acta Math. Hung.*, 99, 203-208.

3rd International Conference: Constructive Mathematical Analysis

Non-Instantaneous Impulsive Differential Equations with a Caputo Fractional Derivative with respect to Other Functions and Strict Stability

Snezhana Hristova

Faculty of Mathematics and Informatics, Plovdiv University, Plovdiv, Bulgaria

snehri@gmail.com

key-words: Strict stability, noninstantaneous impulses, Caputo derivatives with respect to another function, fractional differential equations, Lyapunov functions.

Abstract:

A nonlinear system of fractional differential equations with noninstantaneous impulses is considered. Caputo fractional derivatives with respect to others function (CFDF) are applied (see,[1]). The main characteristics of these fractional derivatives are two: first, the lower limit of CFDF changes at any starting time of action of the noninstantaneous impulses, the applied function in CFDF is changeable at each interval without impulses. The strict stability for the studied system is defined (see, for example, [2, 3, 4]) As an auxiliary system we consider a system of two linear scalar noninstantaneous impulsive fractional equations with CFDF and define a strict stability in couple. The introduced definitions are illustrated on several examples. We use appropriate Lyapunov functions to obtain sufficient conditions for strict stability of the studied system. These sufficient conditions depends significantly on the type of the impulsive functions.

Acknowledgments: This research was partially funded by Bulgarian National Science Fund under Project KP-06-N62/1.

References

- [1] Almeida, R. (2017). A Caputo Fractional Derivative of a Function with respect to Another Function. Commun. Nonl. Sci. Numer. Simul. 44, 460-481.
- [2] Lakshmikantham, V. Mohapatra, R. N. (2001). Strict Stability of Differential equations. Nonlinear Anal., 46(7), 915-921.
- [3] Agarwal, R. P., Hristova, S. and O'Regan, D. (2015). Lyapunov Functions and Strict Stability of Caputo Fractional Differential Equations. Adv. Differ. Equ., 2015, 346. <https://doi.org/10.1186/s13662-015-0674-5>
- [4] Liu, K., Yang G. (2007). Strict Stability Criteria for Impulsive Functional Differential Systems. J. Inequal. Appl., 2008, 243863. <https://doi.org/10.1155/2008/243863>

3rd International Conference: Constructive Mathematical Analysis

Analysis of the mKdV Equation in Magnetohydrodynamics for Exploring Solitons and Nonlinear Dynamics

Muhammad Abubakar Isah¹, Asif Yokus²

¹Department of Mathematics, Istanbul Ticaret University, Istanbul, Türkiye

²Department of Mathematics, Firat University, Elazig, Türkiye

maisah@ticaret.edu.tr, asfyokus@firat.edu.com.tr

key-words: The mKdV equation, the φ^6 -model expansion technique, soliton solution.

Abstract:

In this study, we construct the traveling wave solutions for the nonlinear mKdV equation using the φ^6 -model expansion approach. The KdV equation is applied in shallow water and stratified flows, classical and magnetohydrodynamics, plasma physics and other scientific fields. The explicit retrieval of a wide range of solution types, including bright, singular, dark, and singular soliton solutions, is made possible by the φ^6 -model expansion technique. Another kind of solution that are explicitly retrieved are kink-type solitons, which are referred to as topological solitons in the context of water waves. The nonlinear dynamical features of the equation may be improved by the outcomes of this research. The method proposes a useful and effective approach for solving a large class of non-linear partial differential equations. Interesting graphs are used to illustrate and highlight the dynamical characteristics of the solutions obtained.

References

- [1] Zayed, E. M. E., Alurfi, K. A. E. (2014). The $(G'/G, 1/G)$ -Expansion Method and Its Applications for Solving Two Higher Order Nonlinear Evolution Equations. Math. Probl. Eng., 2014(1), 746538.
- [2] Yokus, A., Isah, M. A. (2025). Unveiling Novel Insights: Nonlinear Dispersion Dynamics of Optical Solitons Through New φ^6 -model Expansion in the Klein–Gordon equation. Chin. J. Phys., 95, 476–492.
- [3] Duran, S., Yokus, A. and Kaya, D. (2025). Evolution Dynamics of Fundamental Mechanisms of Wave Propagation in Gas Bubble-liquid Interactions and Soliton Solutions. Phys. Scr., 100(2), p.025229.

3rd International Conference: Constructive Mathematical Analysis

Fixed Points in Δ -symmetric Quasi Modular Metric Spaces of Non-Archimedean Type

Dilek Karabacak Kesik¹, Mahpeyker Öztürk¹

¹ Department of Mathematics, Sakarya University, Sakarya, Türkiye

dilek.kesik2@ogr.sakarya.edu.tr, mahpeykero@sakarya.edu.tr

key-words: Non-Archimedean quasi modular metric spaces, Δ -symmetry, fixed point theorems.

Abstract:

This study aims to present a novel and transformative concept of a non-Archimedean Δ -symmetric quasi modular metric space, which represents a significant generalization that effectively integrates the characteristics of quasi metrics, modular metrics, and Δ -symmetric frameworks. By employing this innovative structure, we derive Matkowski-type fixed point theorems under relaxed contractive conditions, thereby extending classical findings into a more intricate and asymmetric realm. Furthermore, we undertake a comprehensive analysis of the topological and convergence properties elicited by quasi modular metrics, revealing their complex interrelationships with traditional metric topologies. Our approach not only pushes the boundaries of existing research in fixed point theory but also significantly broadens its scope by removing continuity assumptions and incorporating variable iterations within the transformative context of the non-Archimedean and Δ -symmetric paradigm.

References

- [1] Banach, S. (1922). Sur les operations dans les ensembles abstraits et leurs applications aux equations integrales, Fund. Math., 3(1), 133-181.
- [2] Karapınar, E., Roldan Lopez de Hierro, A. F., Samet, B. (2016). Matkowski theorems in the context of quasi metric spaces and consequences on G-metric spaces, Anal. Stiint. Ale Univ., Ovidius Constanța-Ser. Math., 24, 309-333.
- [3] Chistyakov, V. V. (2006). Metric modulars and their application, Dokl. Math., 73, 32-35.
- [4] Paknazar, M., De la Sen, M. (2017). Best proximity point results in non-Archimedean modular metric spaces, Mathematics, 5(2), 23.
- [5] Matkowski, J. (1977). Fixed point theorems for mappings with a contractive iterate at a point, Proc. Amer. Math. Soc., 62(2), 344-348.

3rd International Conference: Constructive Mathematical Analysis

A Comparative Stability Analysis of Fixed Point Iterations

Osman Alagoz¹, Rumeysa Karaboga²

¹ Department of Mathematics, Bilecik Seyh Edebali University, Bilecik, Türkiye

² Department of Mathematics, Bilecik Seyh Edebali University, Bilecik, Türkiye

osman.alagoz@bilecik.edu.tr, rumeysakaraboga2025@gmail.com

key-words: Stability, fixed point, functional analysis.

Abstract:

In this work, we conduct a comparative stability analysis of several well-known fast iterative processes for contractive mappings in Banach spaces. Specifically, we examine the M-iteration and Kadioğlu and Yildirim iteration processes providing a theoretical assessment of their stability conditions and convergence rates. The theoretical findings are supported by numerical experiments to illustrate their practical performance. Our results highlight how these iterative methods respond to variations in contraction parameters and initial conditions, offering valuable insights for selecting appropriate approaches in applied mathematics and computational sciences.

References

- [1] Alagoz, O. (2018). Numerical Reckoning Fixed Points for Berinde Mappings via a Faster iteration Process. Ser. Math. Inform., 33(2), 295-305.
- [2] Sahu, O., and Banerjee, A. (2024). Convergence, stability, and data dependence results for a new iteration method in Banach space. Electronic Journal of Mathematical Analysis and Applications, 12(1), Article No: 12
- [3] Hyers, D. H. (1941). On the stability of the linear functional equation. *Proceedings of the National Academy of Sciences of the United States of America*, 27(4), 222–224. <https://doi.org/10.1073/pnas.27.4.222>

3rd International Conference: Constructive Mathematical Analysis

Benchmarking Iterative Processes on Contractive Mappings with Numerical Applications

Osman Alagoz¹, Halil Ibrahim Karaboga¹

¹ Department of Mathematics, Bilecik Seyh Edebali University, Bilecik, Türkiye

osman.alagoz@bilecik.edu.tr, halil39572@gmail.com

key-words: Rate of convergence, fixed point, M-iteration.

Abstract:

In this study, we compare the convergence speed of several iterative methods for contractive mappings, focusing on the M-iteration and Kadioğlu–Yıldırım iteration. Using Berinde’s definition, we theoretically and numerically show that these methods converge faster than classical ones such as Picard, Mann, and Ishikawa. The findings help identify more efficient schemes for solving fixed point problems.

References

- [1] Alagoz, O., Gunduz, B., and Akbulut, S. (2018). Numerical reckoning fixed points for Berinde mappings via a faster iteration process. **Facta Universitatis, Series: Mathematics and Informatics**, 33(2), 295–305. <https://doi.org/10.22190/FUMI1802295A>
- [2] Alagoz, O., Gunduz, B., and Akbulut, S. (2020). Convergence theorems with a faster iteration process for Suzuki’s generalized non-expansive mapping with numerical examples. **Erzincan University Journal of Science and Technology**, 13(SPECIAL ISSUE-I), 162–174. <https://doi.org/10.18185/erzifbed.639027>
- [3] Thakur, B. S., and Others. (2016). A new iteration scheme for approximating fixed points of nonexpansive mappings. **Filomat**, 30(10), 2711–2720.

3rd International Conference: Constructive Mathematical Analysis

Remark on Neumann Problem in the Ring Domain

Bahriye Karaca

Fundamental Sciences, Izmir Bakircay University, Izmir, Türkiye

bahriye.karaca@bakircay.edu.tr, bahriyekaraca@gmail.com

key-words: Neumann problem, ring domain, polyanalytic functions.

Abstract:

Recently, several complex model partial differential equations, which hold significant importance in various applied science fields, have been thoroughly examined, particularly in relation to the Neumann problem (see [1]-[3]). Additionally, numerous mathematicians have explored the solvability and solutions of complex partial differential equations under various boundary conditions (see [3]). This paper addresses the Neumann boundary value problem for harmonic functions in a ring-shaped domain within the complex plane. We investigate the existence and uniqueness of solutions to this problem by employing methods from complex analysis and potential theory. The problem is formulated with prescribed normal derivatives on the inner and outer boundaries of the annular domain. Using techniques such as Cauchy Pompeiu Representation Formula and Green's function representation, we derive explicit solution formulas and establish the necessary conditions for the solvability of the problem.

Acknowledgements: The author has been supported by the Scientific and Technological Research Council of Turkey (TUBITAK) under the 2224-B Grant Program for Participation in Scientific Meetings within the Country. The author thanks TUBITAK for their support.

References

- [1] Begehr, H. (2024). Polyanalytic Neumann-n Problem in the Unit Disk. Complex Variables and Elliptic Equations, 1-9.
- [2] Akel, M, Begehr, H. and Mohammed, A. (2022). Integral Representations in the Complex Plane and Iterated Boundary Value Problems. Rocky Mt J Math., 52, 381-413.
- [3] Karaca, B. (2018). Dirichlet Problem for Higher-Order Equations in a Ring Domain. Complex Variables and Elliptic Equations, 68 (12), 2134-2143.

3rd International Conference: Constructive Mathematical Analysis

Continued Fractions in Solving Congruences and Diophantine Equations

Gulsum Ulusoy Ada¹, Busra Karaca¹

¹Institute of Graduate Studies, Cankiri Karatekin University, Cankiri, Türkiye

gulsumulusoy@karatekin.edu.tr, busrkarac@gmail.com

key-words: Fractional differential equations, Riemann-Liouville operators, α -trigonometric functions, mittag-leffler functions, oscillatory solutions, memory effects.

Abstract:

Continued fractions provide an effective framework for representing real numbers and functions. Historically, algorithms such as Euclid's algorithm were employed to address number-theoretic problems, including the solution of Diophantine equations and congruences. This paper explores the historical development and mathematical foundations of continued fractions, emphasizing their role in solving linear Diophantine equations and congruences. We demonstrate how simple continued fractions can be utilized to find rational approximations and solutions to such equations. In particular, we show that a number aa has a multiplicative inverse modulo mm if and only if specific conditions related to continued fractions are satisfied.

References

- [1] Hayek, N., Trujillo, J. J., Rivero, M., Bonilla, B. and Moreno, J. C. (1999). An Extension of Picard-Lindelöf Theorem to Fractional Differential Equations. *Applied Analysis*, 70(3-4), 347-361.
- [2] Kilbas, A. A., Bonilla, B. and Trujillo, J. J. (2000). Fractional Integrals and Derivatives, and Differential Equations of Fractional Order in Weighted Spaces of Continuous Functions. *Doklady Mathematics*, 62(2), 222-226.
- [3] Metzler, R. Klafter, J. (2000). The Random Walk's Guide to Anomalous Diffusion: A Fractional Dynamic Approach. *Physics Reports*, 339(1), 1-77.
- [4] Miller, K. S. Ross, B. (1993). *An Introduction to the Fractional Calculus and Fractional Differential Equations*. Wiley and Sons, New York.
- [5] Mittag-Leffler, M. G. (1903). Sur la nouvelle fonction. *Comptes Rendus de l'Académie des Sciences Paris*, 137, 554-558.
- [6] Podlubny, I. (1999). *Fractional Differential Equations (Mathematics in Science and Engineering 198)*. Academic Press, London.
- [7] Samko, S. G., Kilbas, A. A. and Marichev, O. I. (1993). *Fractional Integrals and Derivatives. Theory and Applications*. Gordon and Breach Science Publ., Switzerland.
- [8] Sokolov, I. M., Klafter, J. and Blumen, A. (2002). Fractional Kinetics. *Physics Today*, 55(11), 48-54.
- [9] Baleanu, D., Mohammadi, H. and Rezapour, S. (2016). On the Existence of Solutions for a Class of Nonlinear Fractional Differential Equations. *Journal of Mathematical Analysis and Applications*, 433(1), 593-602.
- [10] Abbas, S., Benchohra, M. and Agarwal, R. P. (2018). *Topics in Fractional Differential Equations*. Cambridge University Press.

3rd International Conference: Constructive Mathematical Analysis

Approximation by a Generalization of the Szász-Baskakov Operators

Kadir Kanat¹, Melek Sofyalioglu Aksoy¹ and Verda Karadas¹

¹ Mathematics, Ankara Hacı Bayram Veli University, Polatli Faculty of Science and Letters, Ankara, Türkiye

verda.karadas@hbv.edu.tr

key-words: Szász-Baskakov operators, modulus of continuity, Voronovskaya-type theorem.

Abstract:

In this work we obtain a generalization of Szász-Baskakov operators with the help of special polynomials. The moments of the new operator are calculated. For our new operator, an approximation ratio is found by using the first and second modulus of continuity. Also, an asymptotic approximation is obtained by utilizing Voronovskaya-type theorem. Finally, error tables are given with the help of the determined functions.

References

- [1] Altomare, F., Campiti, M. (1994). Korovkin-type Approximation Theory and Its Applications. vol 17. Walter de Gruyter., Berlin.
- [2] Szász, O. (1950). Generalization of S. Bernstein's Polynomials to the Infinite Interval. J. Res. Nat. Bur. Standards., 45(3),239-245.
- [3] Khan, S., Raza, N. (2013). General-Appell Polynomials within the Context of Monomiality Principle. Int. J. Anal., 328032:11.

3rd International Conference: Constructive Mathematical Analysis

Surface Pencil along a Common Spatial Geodesic According to Positional Adapted Frame in Euclidean 3-space

Bilge Karahan¹, Muhammed T. Sariaydin¹

¹ Selcuk University, Department of Mathematics, 42130, Konya, Türkiye.

bilgeugurlu12@gmail.com, talatsariaydin@gmail.com

key-words: Surface pencil, positional adapted frame, Euclidean space

Abstract:

In this study, surface families that pass through a given space curve and consider this curve as a geodesic investigated. Using the positional adapted frame vectors of the curve, the surface family will be constructed as a linear combination of the components of the local coordinate frame, and the necessary and sufficient conditions examined. The surface families obtained theoretically will be supported with examples and visualized using drawing software. Finally, the geometric properties of the resulting new equations analyzed.

References

- [1] Atalay, G. S., Bayram, E., & Kasap, E. (2018). Surface family with a common asymptotic curve in Minkowski 3-space. *Journal of Science and Arts*, 18(2), 357-368.
- [2] Atalay, G. Ş., & Kasap, E. (2015). Surfaces family with common null asymptotic. *Applied Mathematics and Computation*, 260, 135-139.
- [3] Bayram, E., Güler, F., & Kasap, E. (2012). Parametric representation of a surface pencil with a common asymptotic curve. *Computer-Aided Design*, 44(7), 637-643.
- [4] Bilici, M., & Çalışkan, M. (2011). Some New Notes on the involutes of the timelike curves in Minkowski 3-space. *Int. J. Math. Sciences*, 6(41), 2019-2030.
- [5] Camci, C., Ilarslan, K., & Sucurovic, E. (2003). On pseudohyperbolical curves in Minkowski space-time. *Turkish Journal of Mathematics*, 27(2), 315-328.
- [6] Chen, B. Y. (2003). When does the position vector of a space curve always lie in its rectifying plane?. *The American mathematical monthly*, 110(2), 147-152.
- [7] Do Carmo, M. P. (2016). *Differential geometry of curves and surfaces: revised and updated second edition*. Courier Dover Publications.
- [8] Ergün, E., Bayram, E., & Kasap, E. (2014). Surface pencil with a common line of curvature in Minkowski 3-space. *Acta Mathematica Sinica, English Series*, 30, 2103-2118.

3rd International Conference: Constructive Mathematical Analysis

Convergence of Steklov Type Neural Network Operators

Serife Nur Karaman¹, Tuncer Acar¹

¹Department of Mathematics, Selcuk University, Konya, Türkiye

serifenurkaraman@yahoo.com, tunceracar@ymail.com

key-words: Neural networks, Steklov integrals, uniform convergence.

Abstract:

In this talk, we deal with construction of newly family of Neural Network operators, that is, Steklov Neural Network operators. By using Steklov type integral, we introduce a new version of Neural Network operators. We obtain some convergence theorems for the family, such as, pointwise and uniform convergence, rate of convergence via modulus of continuity.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Acar, T., Alagoz, O., Aral, A., Costarelli, D., Turgay, M. and Vinti, G. (2022). Convergence of Generalized Sampling Series in Weighted Spaces. *Demonstr. Math.*, 55, 153-162.
- [2] Costarelli, D. (2024). Convergence and High Order of Approximation by Steklov Sampling Operators. *Banach Journal of Mathematical Analysis*, 18(4), 70.
- [3] Acar, T., Kursun, S. (2023). Pointwise Convergence of Generalized Kantorovich Exponential Sampling Series. *Dolomites Res. Notes Approx.*, 16, 1-10.

3rd International Conference: Constructive Mathematical Analysis

Quantitative Convergence and Multivariate Extensions of Activated Convolution Operators via Positive Linear Methods

Seda Karateke

Department of Software Engineering, Istanbul Atlas University, Istanbul, Türkiye

seda.karateke@atlas.edu.tr

key-words: Trainable half-hyperbolic tangent function, positive linear operator, multidimensional convolution operator, simultaneous approximation.

Abstract:

In this work, we study trainable half-hyperbolic tangent function and convolution-based operators enhanced with activation mechanisms. The work highlights the importance of the "positive linear operators method" in understanding operator behavior and learning dynamics. Using the modulus of continuity, we derive quantitative convergence results towards the identity operator. We also extend the framework to multivariate settings by analyzing both simultaneous and iterative generalizations.

References

- [1] Anastassiou, G. A. (2023). Parametrized, Deformed and General Neural Networks; Springer: Heidelberg/Berlin, Germany; New York, NY, USA, 2023.
- [2] Anastassiou, G. A. (2024). Multivariate Approximation by Symmetrized and Perturbed Hyperbolic Tangent Activated Multidimensional Convolution Type Operators, *Axioms*, 13:779, <https://doi.org/10.3390/axioms13110779>.
- [3] Moldovan, G. (1974). Discrete Convolutions in Connection with Functions of Several Variables and Positive Linear Operators. (Romanian) *Studia Univ. Babes-Bolyai Ser. Math.-Mech.*, 19(1), 51-57.
- [4] Karateke, S. (2025). Some Mathematical Properties of Flexible Hyperbolic Tangent Activation Function with Application to Deep Neural Networks. *Revista de la Real Academia de Ciencias Exactas, Fisicas y Naturales. Serie A. Matematicas*, 119(2), 1-21.
- [5] Bojanic, R., Shisha, O. (1973). On the Precision of Uniform Approximation of Continuous Functions by Certain Linear Operators of Convolution Type. *J. Approximation Theory*, 8, 101-113.

3rd International Conference: Constructive Mathematical Analysis

Stability and Solution of Triple i - f -Hom-Der on Triple Banach Algebras

Vahid Keshavarz¹ and Thabet Abdeljawad²

¹ Department of Mathematics, Shiraz University of Technology, Shiraz, Iran

² Department of Mathematics and Science, Prince Sultan University, Riyadh, Saudi Arabia

v.keshavarz68@yahoo.com, tabdeljawad@psu.edu.sa

key-words: System of additive-quadratic functional equations, triple i - f -hom-der, Hyers-Ulam stability.

Abstract:

In this work, we introduce a new concept of additive and quadratic equations, a type of system of additive-quadratic functional equations on triple Banach algebras. In the following, we define triple i - f -homomorphism-derivation (briefly, triple i - f -hom-der), where $i \in \{1, 2\}$ associated with the additive-quadratic equation where f is a triple linear homomorphism for $i = 1$ and a triple quadratic homomorphism for $i = 2$ on triple Banach algebras. Finally, we employ a fixed point approach to explore the stabilities of the Hyers-Ulam property for system of additive-quadratic functional equations and triple i - f -hom-der, utilizing two control functions from Găvruta and Rassias on triple Banach algebras.

References

- [1] Hyers D. H. (1941). On the stability of the linear functional equation, Proc. Natl. Acad. Sci. U.S.A., 27, 222–224.
- [2] Jahedi S. and Keshavarz V. (2022). Approximate generalized additive-quadratic functional equations on ternary Banach algebras, Journal of Mathematical Extension, 16(1), 1-11.
- [3] Keshavarz V. and Heydari M. T. (2024). Hyers–Ulam stability of solution for generalized Lie bracket of derivations, Journal of Mathematics, 2024(1), 1015443.
- [4] Paokanta S., Dehghanian M., Park C. and Sayyari Y.(2023). A system of additive functional equations in complex Banach algebras, Demonstratio Mathematica, 56, 1-10, 20220165.
- [5] Ulam S. M., (1960). Problems in Modern Mathematics, Chapter IV, Science edn. Wiley, New York.

3rd International Conference: Constructive Mathematical Analysis

An Inverse Source Problem for Time-fractional Pseudoparabolic Equations with Memory Term

Moldir Shazyndayeva^{1,2}, Khomysh Khonatbek^{1,2}

¹Al-Farabi Kazakh National University, Almaty, Kazakhstan

²Institute of mathematics and mathematical modeling, Almaty, Kazakhstan

konat_k@mail.ru, moldir.trz@gmail.com

key-words: Inverse problem, Caputo-type fractional derivative, pseudoparabolic equation.

Abstract:

Let Ω be a bounded domain in R^d , $d \geq 1$, with a smooth boundary and $Q_T := \Omega \times [0, T]$ is a bounded cylinder $0 < T < \infty$. In this work, we consider the following inverse problems for the pseudoparabolic equation with time-fractional Caputo derivative

$$\mathcal{D}_t^\alpha (u - \kappa \Delta u) - \beta \Delta u - \gamma \int_0^t K(t - \tau) \Delta u(x, \tau) d\tau = f(x)g(x, t) + h(x, t), (x, t) \in Q_T, \quad (1)$$

which is supplemented with the initial condition

$$u(x, 0) = u_0(x), \quad x \in \Omega \quad (2)$$

and the boundary condition

$$u(x, t) = 0, \quad t \in [0, T], \quad x \in \partial\Omega, \quad (3)$$

and one of the following overdetermination condition

$$u(x, T) = a(x), \quad x \in \Omega \quad (4)$$

or

$$\int_0^T u(x, t)w(t)dt = a(x), \quad x \in \Omega. \quad (5)$$

Inverse problem I: Finding a pair of functions $\{u(x, t), f(x)\}$ that satisfy (1)-(3) and the final overdetermination condition (4).

Inverse problem II: Finding a pair of functions $\{u(x, t), f(x)\}$ that satisfy (1)-(3) and the integral overdetermination condition (5).

Here, the functions $K(t)$, $g(x, t)$, $u_0(x)$, $a(x)$ and $h(x, t)$ and the constants $\kappa > 0$, $\beta > 0$ and $|\gamma| < \infty$ are given. In this talk, we study the questions of existence and uniqueness of a strong generalized solution of (1)-(4) or (1)-(3),(5).

Funding: This research work has been funded by Grant numbers AP19676624 and AP23486218 the Ministry of Education and Science of the Republic of Kazakhstan (MES RK), Kazakhstan.

3rd International Conference: Constructive Mathematical Analysis

Analysis of Approximation Processes for Fuzzy Numbers via Truncated Max-Product Operators

Sevilay Kirci Serenbay^{1,2}, Rabia Aktas Karaman²

¹Department of Mathematics, Harran University, Şanlıurfa, Türkiye

²Department of Mathematics, Ankara University, Ankara, Türkiye

skserenbay@harran.edu.tr, skserenbay@ankara.edu.tr, raktas@science.ankara.edu.tr

key-words: Max-product operators, shape characteristics, fuzzy numbers.

Abstract:

In this paper, we introduce a class of truncated max-product type operators defined over arbitrary compact intervals and investigate their approximation properties. It is demonstrated that these operators maintain the same order of uniform approximation, along with preserving monotonicity and shape characteristics, similarly to the classical constructions on unbounded domains. Additionally, comparisons and graphical illustrations are presented to emphasize the effectiveness of the proposed operators in the approximation of fuzzy functions.

References

- [1] Majeed, S. Y., Serenbay Kirci, S. (2023). Analyzing Convergence of Fuzzy Numbers Using Generalized Max-Product Bernstein-Chlodowsky Operators on Compact Intervals. *Rare Metal Materials and Engineering*, 52(9), 38-52.
- [2] Majeed, S. Y., Serenbay Kirci, S. (2024). On Approximation by Truncated Max-product Baskakov Operators of Fuzzy Numbers. *Filomat*, 38(9), 3179-3191.
- [3] Majeed, S. Y. (2024). Kesilmiş ve kesilmemiş Maksimum-çarpım Tipi Operatörlerle Bulanık Sayıların Yaklaşımı. Doktora Tezi, Fen Bilimleri Enstitüsü, Harran Üniversitesi, Şanlıurfa.

3rd International Conference: Constructive Mathematical Analysis

Some Basic Results of Eccentricity Sombor Index

Havva Kirgiz¹, Ayse Dilek Maden²

¹ Office of Big Data, Necmettin Erbakan University, Konya, Türkiye

² Department of Mathematics, Selcuk University, Konya, Türkiye

havva.kirgiz@erbakan.edu.tr, aysedilekmeden@selcuk.edu.tr

key-words: Eccentricity, lower bounds, upper bounds.

Abstract:

This paper is related to investigating the upper and lower bounds of eccentricity Sombor index, stated as $SO_{ec}(\mathcal{G})$. In mathematical terms, this index is defined as $SO_{ec}(\mathcal{G}) = \sum_{uv \in E(\mathcal{G})} \sqrt{ec^2(u) + ec^2(v)}$ where $E(\mathcal{G})$ is the set of edges in the graph $\mathcal{G}$, and $ec(.)$

is the eccentricity of a vertex of the edge in $E(\mathcal{G})$. The eccentricity of a vertex is defined as the maximum distance between the vertex and any other vertex in the graph. By analyzing graph structural aspects, the work provides new perspectives on the features and extremal behaviors of this graph invariant. The calculated constraints contribute to our knowledge of the eccentricity Sombor index and its possible applications in graph theory and other domains.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Das, K. C., Çevik, A. S., Cangul, I. N. and Shang, Y. (2021). On Sombor Index. *Symmetry*, 13(1), 140.
- [2] Gutman, I. (2021). Geometric Approach to Degree-based Topological Indices: Sombor Indices. *MATCH Commun. Math. Comput. Chem.*, 86, 11-16.
- [3] Kaya Gök, G., Celik, K. (2024). New Bounds for Sombor Index. *Bull. Int. Math. Virtual Inst.*, 14(2), 267-274.
- [4] Nacaroglu, Y. (2019). Some Bounds for Eccentric Version of Harmonic Index of Graphs. *Ikonion Journal of Mathematics*, 1(1), 11-17.
- [5] Wilson, R. J. (1979). *Introduction to Graph Theory*. Pearson Education India.

3rd International Conference: Constructive Mathematical Analysis

On Optimal BV-control Problem in Coefficients for Elliptic Equation with Variable Order of Nonlinearity

Ismail Aydin¹, Peter Kogut²

¹Department of Mathematics, Sinop University, Sinop, Türkiye

²Mathematical Analysis and Optimization, Oles Honchar Dnipro National University, Dnipro, Ukraine

iaaydinmath@gmail.com, p.kogut@i.ua

key-words: Dirichlet problem, $p(x)$ -Laplace operator, optimal control, control in coefficients.

Abstract:

We deal with the following optimal control problem (OCP) in coefficients for nonlinear elliptic equation with nonstandard growth conditions (see [1] for more details)

$$\begin{aligned} &\text{Minimize } I(u, y) = \int_{\Omega} |y - z_d|^2 dx + \int_{\Omega} |Du| \\ &\text{subject to the constraints} \\ &-\operatorname{div} \left(u |\nabla y|^{p(x)-2} \nabla y \right) = -\operatorname{div} f \quad \text{in } \Omega, \\ &y = 0 \text{ on } \partial\Omega, \\ &u \in \mathfrak{A}_{ad} = \left\{ u \in BV(\Omega) \cap L^\infty(\Omega) \mid \xi_1(x) \leq u(x) \leq \xi_2(x) \text{ a.e. in } \Omega \right\}, \\ &2 \leq \alpha = \operatorname{ess\,inf}_{x \in \Omega} p(x) \leq p(x) \leq \operatorname{ess\,sup}_{x \in \Omega} p(x) = \beta < \infty, \quad \text{a.e. in } \Omega, \end{aligned}$$

where Ω is a bounded open connected subset of $\mathbb{R}^N$ with a Lipschitz boundary $\partial\Omega$, the exponent $p : \Omega \rightarrow [2, \infty)$ is a Lebesgue-measurable function, the function $u \in BV(\Omega) \cap L^\infty(\Omega)$ is taken as a control, and $z_d \in L^2(\Omega)$ and $f \in L^\infty(\Omega; \mathbb{R}^N)$ are given distributions.

The characteristic feature of the considered OCP is the fact that the exponent $p(x)$ is assumed to be Lebesgue-measurable, and we do not impose any additional assumptions on $p(x)$ like to be a Lipschitz function or satisfy the so-called log-Hölder continuity condition.

In order to handle the inherent degeneracy of the $p(x)$ -Laplacian, we use a special two-parametric regularization scheme. We derive existence and uniqueness of variational V -solutions to the underlying boundary value problem and the corresponding optimal control problem. We also discuss the asymptotic behaviour of the solutions to regularized problems on each (ε, k) -level as the parameters tend to zero and infinity, respectively.

References

- [1] Aydin, I, Kogut, P. (2024). Approximation of an Optimal BV -Control Problem in the Coefficient for the $p(x)$ -Laplace Equation. *J. of Optimiz., Diff. Eq. and Their Appl.*, 32(2), 175-204.

3rd International Conference: Constructive Mathematical Analysis

Korovkin Type Approximation Theorem via Statistical Relative Quasi Uniform Convergence

Burçak Kolay

Department of Mathematics, Sinop University, Sinop, Türkiye

burcakyilmaz@sinop.edu.tr

key-words: Relative quasi uniform convergence, positive linear operator, Korovkin-type approximation theorem.

Abstract:

In this paper we present the notion of statistical relative quasi uniform convergence and using this type of convergence, we give a Korovkin-type approximation theorem. Finally, we give an example showing that our results are proper extensions of the corresponding classical ones.

References

- [1] Bukovská, Z., Bukovski, L., & Ewert, J. (1992). Quasi-uniform convergence and L-spaces, 321-329.
- [2] Caserta, A., Di Maio, G., & Kočinac, L. D. (2011). Statistical convergence in function spaces. In Abstract and Applied Analysis (Vol. 2011, No. 1, p. 420419). Hindawi Publishing Corporation.
- [3] Chittenden, E. W. (1914). Relatively uniform convergence of sequences of functions. Transactions of the American Mathematical Society, 15(2), 197-201.
- [4] Hobson, E. W. (1904). On modes of convergence of an infinite series of functions of a real variable. Proceedings of the London Mathematical Society, 2(1), 373-387.
- [5] Gadjiev, A. D., & Orhan, C. (2002). Some approximation theorems via statistical convergence. The Rocky Mountain Journal of Mathematics, 129-138.
- [6] Steinhaus, H. (1951). Sur la convergence ordinaire et la convergence asymptotique. In Colloq. math (Vol. 2, No. 1, pp. 73-74).
- [7] Fast, H. (1951). Sur la convergence statistique. In Colloquium mathematicae (Vol. 2, No. 3-4, pp. 241-244).
- [8] Demirci, K., Boccuto, A., Yıldız, S., & Dirik, F. (2020). Relative uniform convergence of a sequence of functions at a point and Korovkin-type approximation theorems. Positivity, 24, 1-11.
- [9] Gordon, R. A. (1998). When is a limit function continuous?. Mathematics magazine, 71(4), 306-308.

3rd International Conference: Constructive Mathematical Analysis

Convergence of Remote Projections onto Convex Sets

Eva Kopecká

Department of Mathematics, Universität Innsbruck, Innsbruck, Austria

eva.kopecka@uibk.ac.at

key-words: Hilbert space, nearest point projection, convex set, convergence, symmetry.

Abstract:

Let $\{C_\alpha\}_{\alpha \in \Omega}$ be a family of closed and convex sets in a Hilbert space H , having a nonempty intersection C . We consider a sequence $\{x_n\}$ of remote projections onto them. This means, $x_0 \in H$, and x_{n+1} is the projection of x_n onto such a set $C_{\alpha(n)}$ that the ratio of the distances from x_n to this set and to any other set from the family is at least $t_n \in [0, 1]$. We study properties of the weakness parameters t_n and of the sets C_α which ensure the norm or weak convergence of the sequence $\{x_n\}$ to a point in C . We show that the condition

$$\forall \{a_\nu\} \in \ell_2 \text{ with } a_\nu \geq 0 : \liminf_{m \rightarrow \infty} \frac{a_m}{t_m} \sum_{\nu=1}^m a_\nu = 0 \quad (1)$$

is necessary and sufficient for the norm convergence of x_n to a point in C for any starting element and any family of closed, convex, and symmetric sets C_α . This generalizes a result of Temlyakov who introduced (1) in the context of greedy approximation theory. We give examples explaining to what extent the symmetry condition on the sets C_α can be dropped. Condition (1) is stronger than $\sum t_n^2 = \infty$ and weaker than $\sum t_n/n = \infty$. The condition $\sum t_n^2 = \infty$ turns out to be necessary and sufficient for the sequence $\{x_n\}$ to have a partial weak limit in C for any family of closed convex sets C_α and any starting element.

References

- [1] Borodin, P. A., Kopecká, E. (2023). Convergence of Remote Projections onto Convex Sets. Pure and Appl. Funct. Anal., 8(6), 1603-1620.

3rd International Conference: Constructive Mathematical Analysis

Modified Exponential Sampling Series in Logarithmic Weighted Spaces of Functions

Sadettin Kursun¹, Tuncer Acar²

¹ Department of Basic Sciences, National Defence University, Ankara, Türkiye

² Department of Mathematics, Selcuk University, Konya, Türkiye

sadettinkursun@yahoo.com, tunceracar@gmail.com

key-words: Exponential sampling series, logarithmic weighted spaces of functions, modulus of continuity.

Abstract:

In this talk, we consider a modified form of the exponential sampling series constructed by Bardaro et al. [4] in 2017. In constructing the modified exponential sampling series, we consider the remaining part of the Mellin-Taylor formula based on the Mellin derivatives given in [5]. Then, we determine the approximation properties of the newly constructed series in logarithmic weighted spaces of functions. One can refer to [3], [1] and [2] as the first papers where the approximation properties of sampling series of exponential type in logarithmic weighted spaces of functions have been investigated.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports. Authors have been supported by Scientific and Technological Research Council of Turkey (TUBITAK) under the Grant Number 1001-Project 123F123. The authors thank to TUBITAK for their supports.

References

- [1] Acar, T., Kursun, S. (2023). Pointwise Convergence of Generalized Kantorovich Exponential Sampling series. Dolomites Research Notes on Approximation, 16, 1-10.
- [2] Acar, T., Kursun, S. and Acar, O. (2024). Approximation Properties of Exponential Sampling series in Logarithmic Weighted Spaces. Bulletin of the Iranian Mathematical Society, 50:36
- [3] Aral, A., Acar, T. and Kursun, S. (2022). Generalized Kantorovich Forms of Exponential Sampling Series. Analysis and Mathematical Physics, 12(2).
- [4] Bardaro, C. Faina, L. and Mantellini, I. (2017). A Generalization of the Exponential Sampling Series and Its Approximation Properties. Mathematica Slovaca, 67(6), 1481-1496.
- [5] Butzer, P. L., Jansche, S. (1997). A Direct Approach to the Mellin Transform. Journal of Fourier Analysis and Applications, 3, 325-375.

3rd International Conference: Constructive Mathematical Analysis

On the Generic Configurations of Two Dimensional Strongly Competing Systems

Flavia Lanzara

Mathematics Department, Sapienza University, Rome, Italy

flavia.lanzara@uniroma1.it

key-words: Segregation of species, competition-diffusion system, Pattern formation, free boundary problem.

Abstract:

We study some qualitative properties of a well-known free boundary problem modelling strong segregation of N competing populations in planar domains. The main goal is to show that the domain generically divides into subdomains with only triple points, i.e. at most 3 populations meet at any point of the free boundary. To achieve this, we relate the solutions of the problem to a particular class of harmonic maps in singular spaces, which can be seen as the real part of certain holomorphic functions.

This is a joint work with E. Montefusco, V. Nesi and E. Spadaro (Mathematics Department, Sapienza University, Rome, Italy).

References

- [1] Lanzara, F., Montefusco, E. (2021). Some Remarks on Segregation of k Species in Strongly Competing Systems. *Interfaces Free Bound.*, 23, 403-419.
- [2] Lanzara, F., Montefusco, E., Nesi, V. and Spadaro, E. (2025). Generic Configurations in 2D Strongly Competing Systems. Submitted.

3rd International Conference: Constructive Mathematical Analysis

Resolvent Approaches to Elliptic Regularity in Stationary Fokker-Planck Equations

Haesung Lee

Department of Mathematics and Big Data Science, Kumoh National Institute of Technology, Gumi, Republic of Korea

fthslt@kumoh.ac.kr, fthslt14@gmail.com

key-words: Stationary Fokker-Planck equations, Dirichlet forms, resolvents.

Abstract:

In this talk, we present the regularity results established in [1] for solutions to the stationary Fokker-Planck equation with general coefficients in dimensions $d \geq 3$. Specifically, we show that when the solution is interpreted as a measure with an L^2 -density, this density not only belongs to the Sobolev space $W^{1,2}$ but also enjoys Hölder continuity. Our approach relies on constructing a reference measure $\mu = \rho dx$ through elliptic regularity and existence results, ensuring that the associated divergence-type operator fits within the framework of sectorial Dirichlet forms. By combining elliptic regularity results for both divergence-type and non-divergence-type equations, we establish $W^{2,2}$ -regularity for the image of the resolvent operator. Furthermore, we show that the solution density arises as the weak limit of $W^{1,2}$ -functions constructed via the resolvent, revealing the essential role of Dirichlet form techniques and resolvent approximations in understanding the regularity of stationary Fokker-Planck equations. We also anticipate that with refined estimates, our approach can extend to the two-dimensional case.

References

- [1] Lee, H. (2025). Local Elliptic Regularity for Solutions to Stationary Fokker-Planck Equations via Dirichlet Forms and Resolvents. *Boundary Value Problems*, 2025, Article number: 68. <https://doi.org/10.1186/s13661-025-02056-0>

3rd International Conference: Constructive Mathematical Analysis

Boundary Integral Equations of the First Kind: Solvability and Applications

Alberto Cialdea¹, Vita Leonessa² and Angelica Malaspina¹

¹Department of Basic and Applied Sciences, University of Basilicata, Potenza, Italy

²Department of Engineering, University of Basilicata, Potenza, Italy

alberto.cialdea@unibas.it, vita.leonessa@unibas.it, angelica.malaspina@unibas.it

key-words: Boundary integral equations, potential theory, reducible operator, differential form.

Abstract:

In this talk, we focus on a method for solving boundary integral equations of the first kind, which naturally arise when seeking solutions to certain boundary value problems for elliptic differential equations in terms of single or double layer potentials. The resulting integral representations differ from those obtained through direct methods. One potential advantage of this alternative formulation is that the corresponding boundary integral equation may be more manageable, either numerically or analytically, depending on the specific context (see, e.g., [5]).

Coming back to the method used, it was introduced by Cialdea in [1] and it is based on the theory of reducible operators and the theory of differential forms.

Such a method has been extended to different boundary value problems associated with partial differential equations and systems, both in simply and multiple connected domains (see, e.g. [2,3,4] and the recent paper [6]).

References

- [1] Cialdea, A. (1988). On the Oblique Derivative Problem for the Laplace Equation, and Related Topics. *Rendiconti della Accademia Nazionale delle Scienze detta dei XL*, 12(1), 181-200.
- [2] Cialdea, A., Leonessa, V. and Malaspina, A. (2012). On the Dirichlet and the Neumann Problems for Laplace Equation in Multiply Connected Domains. *Complex Variables and Elliptic Equations*, 57(10), 1035-1054.
- [3] Cialdea, A., Leonessa, V. and Malaspina, A. (2015). The Dirichlet Problem for Second-order Divergence form Elliptic Operators with Variable Coefficients: the Simple Layer Potential Ansatz. *Abstract and Applied Analysis* 2015, Art. ID 276810, 11 pp.
- [4] Cialdea, A., Leonessa, V. and Malaspina, A. (2023). On the Traction Problem for Steady Elastic Oscillations Equations: the Double Layer Potential Ansatz. *Rendiconti del Circolo Matematico di Palermo, II. Ser.*, 72, 1947-1960.
- [5] Hsiao, G. (1989). On Boundary Integral Equations of the First Kind. *Journal of Computational Mathematics*, 7(2), 121-31.
- [6] Leonessa, V. Malaspina, A. (2025). On the Dirichlet Problem for the System of Thermoelasticity with Microtemperatures. *Mediterr. J. Math.*, 22, article number 95.

3rd International Conference: Constructive Mathematical Analysis

Existence and Monotonicity of Principal Eigenvalues for a Class of Fractional Boundary Value Problems

Yulong Li

Department of Mathematics, University of Dayton, Dayton, USA

liyulong0807101@gmail.com

key-words: Fractional boundary value problem, principal eigenvalue, fixed point theory.

Abstract:

In this talk, we investigate the spectral problem

$$\begin{cases} (-\mathcal{D}_{a+}^{\alpha}u)(x) = \lambda u, & x \in (a, b), \\ u(a) = u(b) = 0, & 1 < \alpha < 2, \end{cases}$$

where $\mathcal{D}_{a+}^{\alpha}$ represents the Riemann-Liouville fractional derivative. Using fixed point theory and positive operator theory, we establish the existence of principal eigenvalues for this problem. Interestingly, while the principal eigenvalue $\lambda(\alpha)$ is generally non-monotonic as a function of the order α , we show that the normalized quantity

$$\frac{\Gamma(\alpha - 1)}{(b - a)^{\alpha - 1}} \frac{1}{\lambda(\alpha)}$$

is strictly decreasing with respect to α .

If time permits, we will also discuss extensions to nonlinear cases of $f(u, x)$ and to a broader range of the fractional order parameter $1 < \alpha < \infty$.

3rd International Conference: Constructive Mathematical Analysis

Design Construction and Model Selection for Small Mixture-Process Variable Experiments with High-dimensional Model Terms

Kashinath Chatterjee¹, Chang-Yun Lin²

¹ Department of Population, Augusta University, Augusta, Georgia, USA

² Department of Applied Mathematics and Institute of Statistics, National Chung Hsing University, Taichung, Taiwan

kashinathchatterjee077@gmail.com, chlin6@nchu.edu.tw

key-words: Bayesian, D-optimality criterion, generalized Bayesian-D criterion, split-plot design.

Abstract:

This paper considers the design construction and model selection for mixture-process variable experiments where the number of variables is large. For such experiments the generalized least squares estimates cannot be obtained and hence it will be difficult to identify the important model terms. To overcome these problems, here we employ the generalized Bayesian-D criterion to choose the optimal design and apply the Bayesian analysis method to select the best model. Two algorithms are developed to implement the proposed methods. A fish-patty experiment demonstrates how the Bayesian approach can be applied to a real experiment. Simulation studies show that the proposed method has a high power to identify important terms and well controls the type I error.

3rd International Conference: Constructive Mathematical Analysis

Sensitivity Approach for the Evaluation of Statistical Test Batteries

Elena Almaraz Luengo

Department of Statistic and Operational Research, Complutense University of Madrid (UCM), Spain, Interdisciplinary Mathematics Institute (IMI), Group of Analysis, Security and Systems (GASS, UCM)

ealmaraz@ucm.es

key-words: FIPS 140-2, pseudo-random numbers, sensitivity, tests battery, tests suite.

Abstract:

For the verification of statistical properties such as randomness or uniformity, different hypothesis tests, generally non-parametric, are used to validate or not the generators that produced such sequences. In professional practice, in areas such as simulation, Monte Carlo techniques, applications in security and encrypted communication systems, etc., such verifications are performed by applying sets of tests called suites or batteries [1]. These batteries must be robust and contain uncorrelated tests to avoid possible interpretation biases and high computational times.

A current line of research is focused precisely on the analysis of the structure of batteries in order to make them as complete, robust and efficient as possible, see for example [2, 3, 4] or [6] among others.

In this paper we will discuss the main techniques used for battery analysis and also focus on the sensitivity of the FIPS 140-2 [5] tests battery.

References

- [1] Almaraz Luengo, E., García Villalba, L. J. (2021). Recommendations on Statistical Randomness Test Batteries for Cryptographic Purposes. *ACM Computing Surveys*, 54(4).
- [2] Almaraz Luengo, E., Leiva, M., García Villalba, L. J., Hernández-Castro, J. and Hurley-Smith, D. (2022). Critical Analysis of Hypothesis Tests in Federal Information Processing Standard (140-2). *Entropy*, 24(5), 1-16.
- [3] Almaraz Luengo, E., Alaña, B., García Villalba, L. J. and Hernández-Castro, J. D. (2022). Weaknesses in ENT Battery Design. *Applied Sciences*, 12(9), 1-7.
- [4] Burciu, P., Simion, E. (2019). A Systematic Approach of NIST Statistical Tests Dependencies. *Journal of Electrical Engineering, Electronics, Control and Computer Science*, 5(1), 1-6.
- [5] Information Technology Laboratory. National Institute of Standards and Technology (2001). FIPS PUB 140-239. url: <https://csrc.nist.gov/publications/detail/fips/140/2/final>
- [6] Haramoto, H., Matsumoto, M. (2019). Checking the Quality of Approximation of p -values in Statistical Tests for Random Number Generators by Using a Three-level Test. *Math. Comput. Simul.*, 161, 66-75.

3rd International Conference: Constructive Mathematical Analysis

Fake Nodes Approach and Some Applications

Stefano De Marchi

Department of Mathematics “Tullio Levi-Civita”, University of Padova, Padova, Italy

stefano.demarchi@unipd.it

key-words: Mapped basis, fake nodes approach.

Abstract:

The concept of mapped bases has been widely studied, but all the proposed methods show convergence provided that the function is resampled at the mapped nodes. In applications, this is often physically unfeasible. We discuss the extension of the so-called *mapped bases without resampling* interpolation, also known as *Fake Nodes Approach* (FNA) [1, 2], to any basis and dimension. The univariate case has been discussed in [1] and some of its applications have been collected in the recent paper [2]. In the talk we show some relevant applications of this “fancy” approach.

References

- [1] De Marchi, S., Marchetti, F., Perracchione, E., and Poggiali, D. (2020). Polynomial Interpolation via Mapped Bases without Resampling. *J. Comput. Appl. Math.*, 364, p.112347.
- [2] De Marchi, S., Marchetti, F., Elefante, G. and Francomano, E. (2022). Polynomial Mapped Bases: Theory and Applications. *SIMAI Comm. Appl. Ind. Math.*, 22, 1-9.
- [3] De Marchi, S., Marchetti, F., Perracchione, E., and Poggiali, D. (2021). Multivariate Approximation at Fake Nodes. *Appl. Math. Comput.*, 391, 125628.
- [4] Poggiali, D., Cecchin, D. and De Marchi, S. (2022). Reducing the Gibbs Effect in Multimodal Medical Imaging by the Fake Nodes Approach. *J. Comput. Math. and Data Science*, 4, 100040.

3rd International Conference: Constructive Mathematical Analysis

Rational Approximations of Fractional Elliptic Operators

Stanislav Harizanov¹, Svetozar Margenov¹

¹Institute of Information and Communication Technologies, Bulgarian Academy of Sciences, Sofia, Bulgaria

sharizanov@parallel.bas.bg, margenov@parallel.bas.bg

key-words: Fractional elliptic operator, rational approximation, exponential convergence.

Abstract:

Let $\mathcal{A}$ be a second-order elliptic differential operator in a bounded multidimensional domain Ω with general geometry. We consider the spectral definition of $\mathcal{A}^\alpha$. Over the past decade, several approaches have been proposed to approximate $\mathcal{A}^{-\alpha}$, $\alpha \in (0, 1)$. Despite their different origin, they can all be written as a rational approximation.

We first discuss the BURA method introduced in [2] (see also the survey [1]). It is defined in the form $\mathcal{A}^{-\alpha} \approx \lambda_1^{-\alpha} r_{\alpha,k}(\lambda_1 \mathcal{A}^{-1})$. Here $r_{\alpha,k}(z)$ is the best uniform rational approximation of degree k to z_1^α , $z \in [0, 1]$ and $\lambda_1 > 0$ is the minimal eigenvalue of $\mathcal{A}$. We demonstrate how the additive representation of the BURA approximation is a useful tool for qualitative analysis of $\mathcal{A}^\alpha$. We also show that the error decreases exponentially as k increases.

Let now the symmetric and positive definite matrix A be a mesh approximation of $\mathcal{A}$. A^α is dense, which is consistent with the nonlocality of $\mathcal{A}^\alpha$. The computational efficiency of the BURA method is due to the representation $A^{-\alpha} \approx \tilde{c}_0(I + \sum_{i=1}^k \tilde{c}_i(A - \tilde{d}_i I)^{-1})$, where $\tilde{c}_i \geq 0$ and $\tilde{d}_i < 0$. Near-optimal estimates of computational complexity are obtained that do not depend on the condition number of A .

The last part of the talk is devoted to generalizations regarding the power α . The presentation follows and further develops the results of [3]. First, the advantages and disadvantages of the BURA and BURA-based methods for $\alpha > 0$ are discussed. Finally, the case $\alpha < 0$ is considered. It turns out that it is the most complex. Thus, approximating the fractional elliptic operator is more difficult than approximating its inverse. Accordingly, in the discrete case, developing a numerical method for multiplying a vector by A^α is a more difficult problem than solving a linear system with the same matrix.

References

- [1] Harizanov, S., Lazarov, R. and Margenov, S. (2020). A Survey on Numerical Methods for Spectral Space-fractional Diffusion Problems. *Fractional Calculus and Applied Analysis*, 23(6), 1605–1646.
- [2] Harizanov, S., Lazarov, R., Margenov, S., Marinov, P. and Pasciak, J. (2020). Analysis of Numerical Methods for Spectral Fractional Elliptic Equations based on the Best Uniform Rational Approximation. *Journal of Computational Physics*, 408, 109285.
- [3] Kosturski, N., Margenovi, S. (2024). Analysis of BURA and BURA-based Approximations of Fractional Powers of Sparse SPD Matrices. *Fractional Calculus and Applied Analysis*, 27, 706–724.

3rd International Conference: Constructive Mathematical Analysis

On the Complete Controllability of Fractional-Order Integrodifferential System Involving Generalized Conformable Derivatives

Maher Jneid¹, Rola Masri¹

¹Department of Mathematics and Computer Science, Beirut Arab University, Tripoli, Lebanon

m.jneid@bau.edu.lb, rmm462@student.bau.edu.lb.

key-words: Integrodifferential system, generalized conformable derivatives, semigroup theory, contraction mapping principle, complete controllability.

Abstract:

Controllability concepts have been essential across various disciplines, including engineering, robotics, optimal control, and applied mathematics. Following Kalman's definition, controllability is characterized by the ability to steer the solution of a dynamical system from any initial state to a desired target at a given finite time. In this paper, the complete controllability of fractional integrodifferential systems involving generalized conformable derivatives is investigated. A set of sufficient conditions for the complete controllability of Fractional-Order Integrodifferential semilinear systems is established, assuming that the associated linear part is completely controllable. The obtained results based on employing advanced mathematical tools, including the semigroup theory, fractional calculus, and the contraction mapping principle. A concrete example is provided at the end to illustrate the practical effectiveness of the theoretical findings.

Throughout this paper, the following fractional nonlinear system will be considered:

$$\begin{cases} {}^{GC}D^p z(t) = Az(t) + Bv(t) + h\left(t, z(t), \int_0^t g(t, r, z(r)) dr\right) & t \in]0, a] \\ z(0) = z_0 \in Z, \end{cases} \quad (1)$$

Z is a Hilbert space, ${}^{GC}D^p$ is the generalized conformable derivative of order $0 < p < 1$, $A : Z \rightarrow Z$ is the infinitesimal generator of a strongly continuous semi-group $\{S(t), t > 0\}$, $B : V \rightarrow Z$ is a bounded linear operator, $h : [0, a] \times Z \times Z \rightarrow Z$, $g : [0, a] \times [0, a] \times Z \rightarrow Z$ are non-linear continuous functions with appropriate conditions.

Here, some selected references:

References

- [1] Zhao, D., Luo, M. (2017). General Conformable Fractional Derivative and Its Physical Interpretation. *Calcolo*, 54, 903-917.
- [2] Sarıkaya, M. Z., Budak, H. and Usta, H. (2019). On Generalized the Conformable Fractional Calculus. *TWMS Journal of Applied and Engineering Mathematics*, 9(4), 792-799.
- [3] Azouz, F., Boucenna, D., Ben Makhlouf, A., Mchiri, L. and Benchaabane, A. (2021). Controllability of Differential Systems with the General Conformable Derivative. *Complexity*, 2021, (1), 2817092.
- [4] Atraoui, M., Bouaoud, M. (2021). On the Existence of Mild Solutions for Nonlocal Differential Equations of the Second Order with Conformable Fractional Derivative. *Advances in Difference Equations*, 2021, 1-11.

3rd International Conference: Constructive Mathematical Analysis

On Some Results Related to Positive Definite Matrices by Means of Cauchy-Schwarz Inequality

Mahmut Furkan Meniz¹, Ramazan Turkmen¹

¹Department of Mathematics, Selcuk University, Konya, Türkiye

menizfurkan1@gmail.com, rturkmen@selcuk.edu.tr

key-words: Cauchy-Schwarz inequality, singular values, majorization, unitarily invariant norms.

Abstract:

In this study, the relationship between the Cauchy-Schwarz inequality and matrix norms and singular values is discussed. First, the classical form of the Cauchy-Schwarz inequality is recalled, and its extended versions in matrix space are examined. Then, the connections between singular values and unitarily invariant norms are addressed, and various inequalities are derived in this context. In particular, the majorization properties of singular values for Hermitian and positive semi-definite matrices are investigated. In addition, using some special matrix functions like Lieb functions and matrix convex functions, we get some inequalities for singular values and norms of positive semidefinite matrices.

References

- [1] Kittaneh, F. (1988). Notes on Some Inequalities for Hilbert Space Operators. Publ. Res. Inst. Math. Sci., 24(2), 283–293.
- [2] Kato, T. (1952). Notes on Some Inequalities for Linear Operators. Math. Ann., 125, 208-212.
- [3] Papat, R. B. (1987). Majorization and Singular Values. Linear and Multilinear Algebra, 21, 211-214.

3rd International Conference: Constructive Mathematical Analysis

A New Generalization of the Frank Matrix

Efruz Ozlem Mersin

Department of Mathematics, Aksaray University, Aksaray, Türkiye

efruzmersin@aksaray.edu.tr

key-words: Binomial matrices, max matrix, Frank matrix.

Abstract:

Matrices play a fundamental role in various areas of mathematics, including combinatorics, numerical analysis, and linear algebra. The literature includes significant papers on matrices whose entries are binomial coefficients, such as Pascal matrices. Max matrices, whose elements are defined as the maximum of their respective row and column indices, are other special matrices in matrix theory. Among them, the Frank matrix has recently attracted significant attention from researchers. Several studies have examined various generalizations of the Frank matrix and analysed their basic properties. This paper introduces a new generalization of the Frank matrix, called the binomial Frank matrix, and investigates some of its properties, including the determinant, inverse, adjoint matrix, LU decomposition, and certain norms. This study contributes to the literature by offering a novel approach to extending Frank matrices and providing insights into their algebraic and analytical properties.

References

- [1] Frank, W. L. (1958). Computing Eigenvalues of Complex Matrices by Determinant Evaluation and by Methods of Danilewski and Wielandt. *Journal of the Society for Industrial and Applied Mathematics*, 6(4), 378-392.
- [2] Hake, J.-F. (1985). A Remark on Frank Matrices. *Computing*, 35(3-4), 375-379.
- [3] Higham, N. J. (2024). The Power of Bidiagonal Matrices. *Electronic Journal of Linear Algebra*, 40, 453-474.
- [4] Shi, B., Kizilates, C. (2024). A New Generalization of the Frank Matrix and Its Some Properties. *Computational and Applied Mathematics*, 43(1), 19.
- [5] Strum, J. E. (1977). Binomial Matrices. *The Two-Year College Mathematics Journal*, 8(5), 260-266.
- [6] Mersin, E. O., Bahsi, M. (2021). Sturm Theorem for the Generalized Frank matrix. *Hacettepe Journal of Mathematics and Statistics*, 50(4), 1002-1011.
- [7] Mersin, E. O, Bahsi, M. (2024). Bounds for the Maximum Eigenvalues of the Fibonacci-Frank and Lucas-Frank Matrices. *Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics*, 73(2), 420-436.
- [8] Mersin, E. O, Bahsi, M. and Maden, A. D. (2020). Some Properties of Generalized Frank Matrices. *Mathematical Sciences and Applications E-Notes*, 8(2), 170-177.
- [9] Varah, J. M. (1986). A Generalization of the Frank Matrix. *SIAM Journal on Scientific and Statistical Computing*, 7(3), 835-839.

3rd International Conference: Constructive Mathematical Analysis

Highly Efficient Computational Technique for Equal Width Equation Modelling Dispersive Waves

Emre Kirli¹, Mehmet Ali Mersin²

¹Mathematics Department, Aksaray University, Aksaray, Türkiye

²Computer Programming, Aksaray University, Aksaray, Türkiye

emre.26@windowslive.com, mehmetalimersin@gmail.com

key-words: Collocation method, trigonometric B-splines, equal width equation, solitary waves.

Abstract:

The Equal Width (EW) equation, introduced by Morrison et al. [1], is given by

$$w_t + ww_x - \mu w_{xxt} = 0 \quad (1)$$

as a model for nonlinear dispersive waves. Recently, numerical solutions of the EW equation have gained popularity due to their simplicity compared to analytical solutions, which are limited to specific initial and boundary conditions. In this study, we aim to obtain an improved numerical solution by employing a fourth-order two-step scheme for temporal discretization and quartic trigonometric B-splines for spatial discretization. To validate the approach, the propagation of a single solitary wave is simulated. The accuracy of the proposed method is assessed by computing the L_∞ error norm and comparing it with results from the literature. Moreover, the temporal order of convergence and three conserved quantities are numerically evaluated and found to be consistent with their theoretical values. The results indicate that our method offers an efficient alternative compared to recent approaches.

References

- [1] Morrison, P. J., Meiss, J. D. and Carey J. R. (1984). Scattering of RLW solitary waves. *Physica D*, 11, 324-336.
- [2] Roshan, T. (2011). A Petrov-Galerkin Method for Equal Width Equation. *Applied Mathematics and Computation*, 218(6), 2730-2739.
- [3] Irk, D. (2012). B-Spline Galerkin Solutions for the Equal Width Equation. *Physics of Wave Phenomena*, 20(2), 122-130.
- [4] Oruc Esen, A., Bulut, F. (2021). Highly Accurate Numerical Scheme based on Polynomial Scaling Functions for Equal Width Equation, *Wave Motion*, 105, 102760.

3rd International Conference: Constructive Mathematical Analysis

On Close to Convexity of a Class of Bounded Harmonic Mappings

Asena Cetinkaya¹, S. Melike Aydogan², Oya Mert Coskun³ and F. Muge Sakar⁴

¹ Department of Mathematics and Computer Sciences, Istanbul Kultur University, Istanbul, Türkiye

² Department of Mathematics, Istanbul Technical University, Istanbul, Türkiye

³ Department of Mathematics, Tekirdag Namik Kemal University, Tekirdag, Türkiye

⁴ Department of Management, Dicle University, Diyarbakir, Türkiye

asnfigen@hotmail.com, aydogansm@itu.edu.tr, oyamert@nku.edu.tr, mugesakar@hotmail.com

key-words: Convex, close-to-convex, harmonic.

Abstract:

The present study introduces a new subclass of bounded harmonic mappings. This subclass will represent existing concepts in the field and aims to provide new insights and potential applications to study harmonic functions. The purpose of defining these mappings is twofold: firstly, to deepen the understanding of their properties, and secondly, to broaden the scope of their usage in various mathematical contexts. This analysis investigates the conditions necessary for certain mappings to exhibit properties that make them approach close to be convex. Furthermore, sharp coefficient estimates are determined.

Acknowledgments: The authors are thankful for the support by the Scientific and Technological Research Council of Turkey (TUBITAK 1002-Short Term R&D Funding Program), Project Number: 124F430

References

- [1] Clunie, J., Sheil-Small, T. (1984). Harmonic Univalent Functions. *Annales Academiae Scientiarum Fennicae. Series A. I. Mathematica*, 9, 3-25.
- [2] Chuaqui, M., Duren, P. and Osgood, B. (2004). Curvature Properties of Planar Harmonic Mappings. *Computational Methods and Function Theory*, 4(1), 127-142.
- [3] Dorff, M., Rolf, J. (2012). Anamorphosis, Mapping Problems, and Harmonic Univalent Functions. In *Explorations in Complex Analysis* (pp. 197-269), Mathematical Association of America.
- [4] Duren, P. (2004). *Harmonic Mappings in the Plane*. Cambridge University Press.
- [5] Nagpal, S., Ravichandran, V. (2014). A Subclass of Close-to-convex Harmonic Mappings. *Complex Variables and Elliptic Equations*, 59(2), 204-216.

3rd International Conference: Constructive Mathematical Analysis

Solving Sequential Linear Fractional Differential Equations Using α -Trigonometric Functions

Gülsüm Ulusoy Ada¹, Mussa Yousef Mohamed¹

¹Institute of Graduate Studies, Çankiri Karatekin University, Çankiri, Türkiye

gulsumulusoy@karatekin.edu.tr, mussablhaj1980@gmail.com

key-words: Fractional differential equations, Riemann–Liouville operators, α -trigonometric functions, mittag-leffler functions, oscillatory solutions, memory effects.

Abstract:

This paper introduces a novel analytical framework for solving sequential linear fractional differential equations (SLFDEs) with constant coefficients using α -trigonometric functions. These functions extend classical trigonometric forms via generalized Mittag-Leffler series to incorporate power-law memory and oscillatory dynamics. The approach enables the explicit construction of both homogeneous and non-homogeneous solutions, preserving non-local memory effects intrinsic to fractional-order systems. A new concept the α -Wronskian is defined to assess the linear independence of solutions, paralleling the classical Wronskian in ordinary differential equations. Applications to fractional harmonic oscillators illustrate the framework's capability in capturing damped and amplified responses driven by memory kernels. The proposed method significantly enhances the modeling of complex physical and engineering systems governed by fractional dynamics.

References

- [1] Carpinteri, A., Mainardi, F. (Eds.). (1997). *Fractals and Fractional Calculus in Continuum Mechanics* (CIAM Courses and Lectures 378). Springer-Verlag, New York.
- [2] Diethelm, K. (2010). *The Analysis of Fractional Differential Equations*. Springer, Berlin.
- [3] Hayek, N., Trujillo, J. J., Rivero, M., Bonilla, B. and Moreno, J. C. (1999). An Extension of Picard-Lindelöf Theorem to Fractional Differential Equations. *Applied Analysis*, 70(3–4), 347–361.
- [4] Kilbas, A. A., Bonilla, B., Trujillo, J. J. (2000). Fractional Integrals and Derivatives, and Differential Equations of Fractional Order in Weighted Spaces of Continuous Functions. *Doklady Mathematics*, 62(2), 222–226.
- [5] Kilbas, A. A., Bonilla, B. and Trujillo, J. J. (2000). Existence and Uniqueness Theorems for Nonlinear Fractional Differential Equations. *Demonstratio Mathematica*, 33(3), 583–602.
- [6] Kilbas, A. A., Srivastava, H. M. and Trujillo, J. J. (2003). Fractional Differential Equations: An Emergent Field in Applied and Mathematical Sciences. In S. Samko, A. Lebre, and A. F. dos Santos (Eds.), *Factorization, Singular Operators and Related Problems* (pp. 151–174). Kluwer Academic Publishers, London.
- [7] Metzler, R. and Klafter, J. (2000). The Random Walk's Guide to Anomalous Diffusion: A Fractional Dynamic Approach. *Physics Reports*, 339(1), 1–77.
- [8] Miller, K. S. Ross, B. (1993). *An Introduction to the Fractional Calculus and Fractional Differential Equations*. Wiley and Sons, New York.
- [9] Mittag-Leffler, M. G. (1903). Sur la nouvelle fonction. *Comptes Rendus de l'Académie des Sciences Paris*, 137, 554–558.
- [10] Podlubny, I. (1999). *Fractional Differential Equations* (Mathematics in Science and Engineering 198). Academic Press, London.
- [11] Samko, S. G., Kilbas, A. A. and Marichev, O. I. (1993). *Fractional Integrals and Derivatives. Theory and Applications*. Gordon and Breach Science Publ., Switzerland.
- [12] Sokolov, I. M., Klafter, J. and Blumen, A. (2002). Fractional Kinetics. *Physics Today*, 55(11), 48–54.
- [13] Baleanu, D., Mohammadi, H. and Rezapour, S. (2016). On the Existence of Solutions for a Class of Nonlinear Fractional Differential Equations. *Journal of Mathematical Analysis and Applications*, 433(1), 593–602.
- [14] Abbas, S., Benchohra, M. and Agarwal, R. P. (2018). *Topics in Fractional Differential Equations*. Cambridge University Press.
- [15] Atangana, A. (2020). *Fractional Calculus with Applications in Engineering and Physics*. Academic Press.
- [16] Jarad, F., Abdelwohab, A. A. and Baleanu, D. (2021). New Fractional Derivatives with Non-singular Kernels. *Advances in the Theory of Nonlinear Analysis and its Applications*, 5(1), 1–14.

3rd International Conference: Constructive Mathematical Analysis

Generalized Gamma-Type Operators

Mirella Cappelletti Montano¹, Vita Leonessa² and Arianna Travaglini²

¹Department of Mathematics, University of Bari Aldo Moro, Bari, Italy

²Department of Engineering, University of Basilicata, Potenza, Italy

mirella.cappellettimontano@uniba.it, vita.leonessa@unibas.it, arianna.travaglini@unibas.it

key-words: Generalized Gamma distribution, approximation by positive linear operators, asymptotic formula, rate of convergence.

Abstract:

In this talk we present a class of integral operators of probabilistic type, recently introduced in [1], which are constructed by means of the generalized Gamma distribution. In particular, we investigate their approximation properties in weighted continuous functions spaces and L^p -spaces, providing some estimates of the rate of convergence by means of different moduli of smoothness (such as, e.g., the ones introduced in [2]). An asymptotic formula is also discussed.

References

- [1] Cappelletti Montano, M., Leonessa V. and Travaglini A. (2025). On the Approximation Properties of Generalized Gamma-type Operators in Several Function Spaces, *Annals of Functional Analysis*, accepted.
- [2] Ditzian, Z., Totik, V. (1987). *Moduli of Smoothness*. Springer Verlag, New York.

3rd International Conference: Constructive Mathematical Analysis

The Probabilistic Characteristics of Ion-Acoustic Waves in a Randomized Plasma Model with Non-extensive Electrons

A. Hussein¹, T. H. Khalil² and M. M. Selim²

¹Applied College, Umm Al-Qura University, Makkah, Saudi Arabia

²Department of Physics, Faculty of Science, Damietta University, New Damietta P. O. 34517, Egypt

ahmostafa@uqu.edu.sa, thkhalil90@gmail.com, selim@du.edu.eg

key-words: Plasmas with uncertain properties, random variable transformation, probability density function, mean, variance, confidence interval.

Abstract:

In this study, a randomized plasma model consisting of positive ions and non-extensive electrons is presented. The uncertain properties of the plasma system are treated as continuous random variables. A probabilistic characterization of ion-acoustic wave propagation is then analyzed by calculating the first probability density functions, along with the corresponding means, variances, and confidence intervals. To achieve this, the random variable transformation (RVT) method [1, 2, 3] is applied as a probabilistic approach. The theoretical findings are validated through numerical simulations, which show strong agreement with physical expectations.

References

- [1] A. Hussein, M. M. Selim, "A general probabilistic solution of randomized radioactive decay chain (RDC) model using RVT technique," *Eur. Phys. J. Plus*, vol. 135, 2020, p. 418. <https://doi.org/10.1140/epjp/s13360-020-00389-6>
- [2] H. Slama, N. A. El-Bedwhey, A. El-Depsy, M. M. Selim, "Solution of the finite Milne problem in stochastic media with RVT Technique," *Eur. Phys. J. Plus*, vol. 132, 2017, p. 505.
- [3] A. Elshekhipy et al., "A numerical computing for the CDF of injective RVT with optimization analysis using Weibull distribution: Applications to stochastic heat transfer," *Alexandria Engineering Journal*, vol. 122, 2025, pp. 665–680. <https://doi.org/10.1016/j.aej.2025.03.028>

3rd International Conference: Constructive Mathematical Analysis

Propagation of Solitons and Nonlinear Behavior in Nonlinear Power Law Fibers

Muhammad Abubakar Isah¹, Ahmad Muhammad²

¹Department of Mathematics, Istanbul Ticaret University, Istanbul, Türkiye

²Department of Physics and Materials Sciences, College of Arts and Sciences, Qatar University, Doha, Qatar.

maisah@ticaret.edu.tr, a.muhammad@qu.edu.qa

key-words: The complex Ginzburg–Landau equation, soliton solutions, the φ^6 -model expansion technique.

Abstract:

In this work, we study the soliton propagation in the context of a detuning factor in nonlinear optics, represented by the complex Ginzburg–Landau equation (CGLE). The φ^6 -model expansion technique is used to explicitly obtain various solutions, including trigonometric, hyperbolic, and rational function solutions. Additionally, singular soliton solutions are seen as positive. Using nonlinear power law fibers, the model is examined. The findings of this work are expected to contribute to the enhancement of the nonlinear dynamical properties of the CGLE. The approach suggests a useful and efficient technique for getting reliable solutions to a large number of nonlinear partial differential equations. Some of the results that were discovered are shown in contour, two-dimensional, and three-dimensional plots.

References

- [1] Isah, M. A., Yokus, A., (2023). A Novel Technique to Construct Exact Solutions for the Complex Ginzburg-Landau Equation Using Quadratic-cubic Nonlinearity Law. *Mathematics in Engineering, Science & Aerospace (MESA)*, 14(1).
- [2] Yokus, A., Isah, M. A., (2025). Unveiling Novel Insights: Nonlinear Dispersion Dynamics of Optical Solitons through New φ^6 -model Expansion in the Klein–Gordon Equation. *Chin. J. Phys.*, 95, 476–492.
- [3] Isah, M.A., Yokus, A. and Kaya, D., (2024). Exploring the Influence of Layer and Neuron Configurations on Boussinesq Equation Solutions via a Bilinear Neural Network Framework. *Nonlinear Dyn*, 112(15), 13361-13377.
- [4] Duran, S., Yokus, A. and Kaya, D., (2025). Evolution Dynamics of Fundamental Mechanisms of Wave Propagation in Gas Bubble-liquid Interactions and Soliton Solutions. *Phys. Scr.*, 100(2), 025229.

3rd International Conference: Constructive Mathematical Analysis

A Constructive Definition of the Riemann Integral on a Separable Banach Space

Timothy Myers

Mathematics Department, Howard University, Washington DC, United States

tmyers@howard.edu

key-words: Riemann integral, Lebesgue integral, separable Banach space.

Abstract:

The goal of this talk is to construct a Riemann integral on a separable Banach space which possesses all of the fundamental properties of the Riemann integral on $\mathbf{R}^n$. Let $\mathcal{B}$ represent a separable Banach space. The paper [1] presents a proof that $\mathcal{B}$ has an isomorphic, isometric embedding in $\mathbf{R}^\infty$. In this work we will use this embedding to define a Riemann integral on special subsets of $\mathcal{B}$, which makes the derivations of most of its properties virtually identical to those of its finite-dimensional analogue. Similar to the Lebesgue integral on $\mathcal{B}$, this Riemann integral has the advantage of equaling a limit of Riemann integrals on $\mathbf{R}^n$ as $n \rightarrow \infty$.

We will use this convergence to study some probability density functions on $\mathcal{B}$.

References

- [1] Gill, T.L., Myers, T. (2019). Constructive Analysis on Banach Spaces. Real Analysis Exchange, 44, 1-36.

3rd International Conference: Constructive Mathematical Analysis

Geometry of Curves as a way Establish Connections between Integrable Nonlinear Equations

Gulgassyl Nugmanova¹, Akbota Myrzakul² and Aidana Azhikhan²

¹ Eurasian National University, Astana, Kazakhstan

² Astana IT University, Astana, Kazakhstan

nugmanovagn@gmail.com, akbotamyrzakul23@gmail.com, aidana.azhikhan@gmail.com

key-words: Spin systems, integrable nonlinear equations, geometry of curves.

Abstract:

We have investigated the spin equation with self-consistent potentials of the form [1]:

$$S_t + S_{xxt} + (vSS_x)_x + \frac{1}{4}(\{S_x, S_t\}S)_x - (wS)_x = 0, \quad (1)$$

$$w_x + \frac{1}{4}(tr(S_x^2))_t = 0, \quad (2)$$

$$v_x + \frac{1}{4}tr(S[S_x, S_t]) = 0, \quad (3)$$

here

$$S = \begin{pmatrix} S_3 & S^- \\ S^+ & -S_3 \end{pmatrix}, \quad S^\pm = S_1 \pm iS_2, \quad S^2 = \Sigma, \quad \Sigma = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

It has been established that equation (1) and equation [1]:

$$q_t + q_{xxt} - vq - (wq)_x = 0, \quad (4)$$

$$r_t + r_{xxt} + vr - (wr)_x = 0, \quad (5)$$

$$v_x + 2(r_{xt}q - r q_{xt}) = 0, \quad (6)$$

$$w_x - 2(rq)_t = 0, \quad (7)$$

where $q(x, t)$ and $r(x, t)$ are complex functions.

are equivalent to each other by means of the geometry of curves and surfaces. Exact soliton solutions of equation (1)-(1) have been found, and the same solutions of equation (4)-(7) have been constructed through that geometric equivalence [2]-[3].

The approach proposed in this study makes it possible to clarify the nature of the close connection between integrable nonlinear partial differential equations through the differential geometry of curves and surfaces.

References

- [1] Sagidullayeva, Z., Nugmanova, G., Myrzakulov, R. and Serikbayev, N. Integrable Kuralay Equations: Geometry, Solutions and Generalizations. <http://arxiv.org/abs/2206.05348>.
- [2] Myrzakul, A., Myrzakulov, R. (2017). Integrable Geometric Flows of Interacting Curves/Surfaces, Multilayer Spin Systems and the Vector Nonlinear Schrödinger Equation. Int. J. Geom. Methods Mod. Phys, 14.
- [3] Myrzakul, A., Nugmanova, G., Serikbayev, N. and Myrzakulov, R. (2021). Surfaces and Curves Induced by Nonlinear Schrödinger-Type Equations ant Their Spin Systems. Symmetry, 13.

3rd International Conference: Constructive Mathematical Analysis

Zero Divisors of a Pythagorean Ring

Raymond Calvin Ochieng¹, Maurice Owino Oduor² and Vitalis Onyang-Otieno¹

¹Strathmore Institute of Mathematical Sciences, Strathmore University, Nairobi, Kenya

²Department of Mathematics, Actuarial and Physical Sciences, University of Kabianga, Kericho, Kenya

rochieng@strathmore.edu, moduor@kabianga.ac.ke, vonyango@strathmore.edu

key-words: Pythagorean ring, zero divisors, quotient ring.

Abstract:

Let P denote the ring of the Pythagorean triples and $Z(P)$ is its subset of zero divisors. It is well known that P is isomorphic to Z^2 . Even though it is straight forward to identify the zero divisors in Z^2 , the identification of zero divisors in P is quite laborious. Since $Z(Z^2)$ is not an ideal of Z^2 , the isomorphism between the two rings implies that $Z(P)$ is not an ideal of P . However, some subsets of $Z(P)$ form ideals of P . In this paper, we have answered the following question: Given that $\varphi : P \rightarrow Z^2$ is an isomorphism, what is $\varphi^{-1}(Z(Z^2))$? The zero divisors in P have been identified and partitioned into ideal forming subsets and consequently constructing, new (quotient) rings P/I and I being ideal forming subset of $Z(P)$. Further, some interesting results on the zero divisor graph of P are obtained.

References

- [1] Anderson, D. F., Livingston, P. S. (1999). The Zero-divisor Graph of a Commutative Ring. *J. Algebra*, 217, 434-447.
- [2] Beauregard, R. A., Suryanarayan, E. R. (2000). Parametric Representation of Primitive Pythagorean Triples, in Nelsen, R. B., *Proofs Without Words: Exercises in Visual Thinking II*. Mathematical Association of America, p. 120, ISBN 978-0-88385-721-2, OCLC 807785075.
- [3] Bernhart, F. R., Price, H. L. (2005). Heron's Formula, Descartes Circles and Pythagorean Triangles. arXiv:math/0701624.
- [4] Carmichael, R. D. (1959). *The Theory of Numbers and Diophantine Analysis*, Dover Publications.
- [5] Dawson, B. (1994). A Ring of Pythagorean Triples, *Missouri Journal of Mathematical Sciences*, 6, 72-77.
- [6] Diestel, R. (2005). *Graph Theory (Graduate Texts in Mathematics)*. Springer. ISSN 0072-5285.
- [7] Driss, B., Jilali, M. and Fouad, T. (2017). Extended Zero- Divisor Graphs of Idealizations. *Communication Korean Mathematical Society*, 32(1), 7-17.
- [8] Grytczuk, A. (1997). Note on a Pythagorean Ring. *Missouri Journal of Mathematical Sciences*, 9, 83-89.
- [9] Spiroff S. and Wickham, C. (2011). A Zero Divisor Graph Determined by Equivalence Classes of Zero Divisors. *Communications in Algebra*, 39(7), 2338-2348.
- [10] Wojtowicz, M. (2000). Algebraic Structures of Some Sets of Pythagorean Triples I. *Missouri Journal of Mathematical Sciences*, 12, 31-35.

3rd International Conference: Constructive Mathematical Analysis

An Alternative Method for Associated Curves in Modified Orthogonal Frame

Nural Yüksel¹, Nurdan Oğraş¹ and Süleyman Şenyurt²

¹ Department of Mathematics, Erciyes University, Kayseri, Türkiye

² Department of Mathematics, Ordu University, Ordu, Türkiye

yükseln@erciyes.edu.tr, nurdanogras@gmail.com, senyurtsuleyman52@gmail.com

key-words: Associated curves, modified orthogonal frame.

Abstract:

This study defines tangent, principal normal and binormal associated curves with modified orthogonal frame in Euclidean 3-space so that each of these curve's vector is in its partner's osculator, normal and rectifying plane, respectively. A new moving frame and its curvatures are generated with each associated curve. In addition the solutions of the distance functions between the curve and its associated curve are investigated and possible results were examined separately for different curves. It is obtained that involute curves belong to the tangent associated curve in general and Bertrand and Mannheim curves belong to the principal normal and binormal associated curves. Finally we take some curves and their associated curves as examples and show them graphically.

References

- [1] Choi, J. H., and Kim, Y. H. (2012). Associated curves of a Frenet curve and their applications. *Applied Mathematics and Computation*, 218(18), 9116-9124.
- [2] Şenyurt, S., Canlı, D., Ayvaci, K. H. (2022). Associated curves from a different point of view in E^3 . *Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics*, 71(3), 826-845.
- [3] Sasai, T. (1984). The fundamental theorem of analytic space curves and apparent singularities of Fuchsian differential equations. *Tohoku Mathematical Journal, Second Series*, 36(1), 17-24.
- [4] Bükçü, B.; Karacan M. K. On the modified orthogonal frame with curvature and torsion in 3-space. *Mathematical Sciences and Applications E-Notes* **2016**, 4(1), 184-188.
- [5] Lone, M. S., Es, H., Karacan, M. K., Bukcu, B. (2019). On some curves with modified orthogonal frame in Euclidean 3-space. *Iranian Journal of Science and Technology, Transactions A: Science*, 43, 1905-1916.
- [6] Elsayied, H. K., Altaha, A. A., Elsharkawy, A. (2021). On some special curves according to the modified orthogonal frame in Minkowski 3-space E_1^3 . *Kasmera*, 49(1), 2-15.
- [7] Elsayied, H. K., Altaha, A. A., Elsharkawy, A. (2022). Bertrand curves with the modified orthogonal frame in Minkowski 3-space E_1^3 . *Rev. Edu*, 392(6), 43-55.
- [8] Elsharkawy, A., Elshenhab, A. M. (2022). Mannheim curves and their partner curves in Minkowski 3-space E_1^3 . *Demonstratio Mathematica*, 55(1), 798-811.

3rd International Conference: Constructive Mathematical Analysis

Fixed Point Theorems in Extended G-Menger Spaces and Application

Ismail Tahiri¹, Lahcen Oumertou¹, Mohammed Lamarti Sefian¹ and Abdelmalek Ouahab²

¹ AMCS Research Team, Dept. Math and Comp Sci, ENS, Abdelmalek Essaadi University, Tetouan, Morocco

² Laboratory (L.A.N.O), Department of Mathematics, Faculty of Science, Mohammed I University, Oujda, Morocco

istahiri@uae.ac.ma, lahcen.oumertou@etu.uae.ac.ma, lamarti.mohammed.sefian@uae.ac.ma,

ouahab05@yahoo.fr key-words: Fixed point, t -norm of H -type, G-Menger spaces, probabilistic φ -contraction mapping.

Abstract:

This paper introduces the concept of extended G-Menger spaces, a novel framework designed to unify and generalize several contemporary metric structures. Our construction provides a comprehensive extension of G-Menger spaces, Menger probabilistic metric spaces, controlled strong b-Menger spaces, and generalized G_b -metric space, offering a more versatile setting for analysis under uncertainty. The core of our investigation lies in establishing new fixed-point theorems for probabilistic φ -contraction mappings within this generalized context. By leveraging the properties of continuous t -norms in conjunction with a newly introduced bounded control function, we rigorously demonstrate the existence and uniqueness of fixed points under well-defined conditions. Beyond the theoretical contributions, we investigate the fundamental topological properties of these spaces. To showcase the practical power and applicability of our framework, we present a direct application to solving a Volterra-type integral equation, thereby bridging abstract theory with a concrete problem in functional analysis. The results presented herein not only subsume and extend a range of existing findings but also contribute a robust new tool to the broader fields of fixed-point theory and probabilistic analysis.

References

- [1] Achoun, Y., Mbarki, A., Tahiri, I., and Sefian, M. L. (2024). A Novel Generalization of Strong Probabilistic b -Metric Spaces and Fixed Point Theorems. *Sahand Communications in Mathematical Analysis*, 21(4), 93-108.
- [2] Jachymski, J. (2010). On probabilistic φ -contractions on Menger spaces. *Nonlinear Analysis*, 73, 1131-1137.
- [3] Jain, K., and Kaur, J. (2019). A generalization of G-metric spaces and related fixed point theorems. *Mathematical Inequalities and Applications*, 22, 1145-1160.
- [4] Menger, K. (2003). Statistical metrics. *Selecta Mathematica*, 2, 433-435.
- [5] Zhou, C., Wang, S., Ćirić, L., and Alsulami, S. M. (2014). Generalized probabilistic metric spaces and fixed point theorems. *Fixed Point Theory and Applications*, 1, 1-15.

3rd International Conference: Constructive Mathematical Analysis

Fixed Point Theory in Graphical Modular Metric Spaces: A Unified Approach via Banach Kannan, and Reich-Type Contractions

Ekber Girgin¹, Elif Özbay Azap² and Mahpeyker Öztürk²

¹ Department of Engineering Fundamental Sciences, Sakarya University of Applied Sciences, Sakarya, Türkiye

² Department of Mathematics, Sakarya University, Sakarya, Türkiye

ekbergirgin@subu.edu.tr, elif.ozbay@ogr.sakarya.edu.tr, mahpeykero@sakarya.edu.tr

key-words: Graphical modular metric space, graphical modular-Banach contraction, graphical modular-Kannan contraction, graphical modular-Reich contraction.

Abstract:

This study introduces and develops the novel framework of graphical modular metric spaces, an innovative synthesis of modular metric theory and graph-based structures. Motivated by recent advances in fixed point theory and the emerging utility of graph-theoretic metrics, we define and analyze non-Archimedean graphical modular metric spaces, extending classical concepts in a structured and topologically enriched setting. Within this framework, we investigate the existence and uniqueness of fixed points for three prominent contraction types- Banach, Kannan, and Reich contractions—in the context of directed graph-preserving mappings, in which the obtained results generalize classical fixed point theorems and provide a unified theoretical foundation for fixed point analysis in modular metric spaces endowed with graph connectivity.

References

- [1] Banach, S. (1922). Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fundamenta Mathematicae*, 3, 133-181.
- [2] Jachymski, J. (2007). The Contraction Principle for Mappings on a Metric Space with a Graph. *Proceed. of the Amer. Math. Soc.*, 135(10), 3241-3248.
- [3] Chistyakov, V. V. (2010). Modular Metric Spaces, I: Basic Concept. *Nonlinear Anal.*, 72, 1-14.
- [4] Chistyakov, V. V. (2010). Modular Metric Spaces, II: Application to Superposition Operators. *Nonlinear Anal.*, 72, 15-30.
- [5] Girgin, E., Öztürk, M. (2019). Modified Suzuki-simulation Type Contractive Mapping in Non-Archimedean Quasi-modular Metric Spaces and Application to Graph Theory. *Mathematics*, 7:769.

3rd International Conference: Constructive Mathematical Analysis

Convergence of a Family of Mellin-Steklov Type Exponential Sampling Operators

Dilek Ozer¹, Tuncer Acar¹

¹Department of Mathematics, Selcuk University, Konya, Türkiye

dilekozer@yahoo.com, tunceracar@gmail.com

key-words: L^p convergence, Mellin-Steklov integrals, exponential-type sampling series.

Abstract:

In this talk, we introduce Mellin-Steklov exponential sampling operators of order r , where $r \in \mathbf{N}$, defined through appropriate Mellin-Steklov integrals. We investigate the approximation properties of these operators in the spaces of continuous and bounded functions and L^p spaces ($1 \leq p < \infty$) on $\mathbf{R}^+$. Additionally, we present a quantitative Voronovskaja- type theorem and examine the convergence behavior of these newly defined operators in logarithmically weighted function spaces.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/ 2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Acar, T., Kursun, S. and Acar, O. (2024). Approximation Properties of Exponential Sampling Series in Logarithmic Weighted Spaces. B. Iran Math. Soc., 50:36.
- [2] Aral, A., Acar, T. and Kursun, S. (2022). Generalized Kantorovich Forms of Exponential Sampling Series. Anal. and Math. Physics, 12(2).
- [3] Bardaro, C., Faina, L. and Mantellini, I.,(2017). A Generalization of the Exponential Sampling Series and Its Approximation Properties. Math. Slovaca, 67(6), 1481-1496.
- [4] Butzer, P. L., Ries, S. and Stens, R. L. (1987). Approximation of Continuous and Discontinuous Functions by Generalized Sampling Series. J. Approx. Theory, 50, 25-39.
- [5] Costarelli, D. (2024). Convergence and High Order of Approximation by Steklov Sampling Operators. Banach J. Math. Anal., 18:70.

3rd International Conference: Constructive Mathematical Analysis

A New Generalization of the Digital Contraction Principle and Image Compression Applications

Aybala Sevde Ozkapu¹, Ozlem Acar¹

¹Department of Mathematics, Selçuk University, Konya, Türkiye

aybalasevdeozkapu@yahoo.com, acarozlem@ymail.com

key-words: Digital contraction, digital metric space, image compression.

Abstract:

In 2000 Branciari [1], defined generalized metric spaces by modifying the triangle inequality condition of metric spaces in the classical meaning with the condition

$$d(x, y) \leq d(x, u) + d(u, v) + d(v, y)$$

for every $x, y \in X$ and different points $u, v \in X$ (each different from x and y). Then in 2014, Jleli and Samet [2], extended the Banach contraction principle by defining a class of Θ -functions on generalized metric spaces. In 2015, Ege and Karaca [3], introduced a new perspective to the literature by defining digital metric spaces and proving Banach's fixed point theorem on these spaces. In this talk, a fixed point result will be given using the Θ -function class in generalised digital metric spaces. Also various image compression applications will be discussed.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Branciari, A. (2000). A Fixed Point Theorem of Banach-Caccioppoli Type on a Class of Generalized Metric Spaces. Publ. Math. Debrecen, 57(1-2), 31-37.
- [2] Jleli, M., Samet, B. (2014). A New Generalization of the Banach Contraction principle. Journal of Inequalities and Applications, 2014, 1-8.
- [3] Ege, O., Karaca, I. (2015). Banach Fixed Point Theorem for Digital Images. J. Nonlinear Sci. Appl, 8(3), 237-245.

3rd International Conference: Constructive Mathematical Analysis

Convergence Properties for a New Version of the Mellin-Gauss-Weierstrass Operators

Firat Ozsarac

Department of Mathematics, Kirikkale University, Kirikkale, Türkiye

firat_ozsarac@hotmail.com

key-words: Convergence, Mellin-Gauss-Weierstrass operators, Voronovskaya-type theorem.

Abstract:

In this presentation, we give the convergence properties for a new version of the Mellin-Gauss-Weierstrass operators. Furthermore, a Voronovskaya-type theorem is expressed. This expression includes Mellin derivatives and the logarithmic moment.

Acknowledgements: The author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Bardaro, C., Mantellini, I. (2011). A Note on the Voronovskaja Theorem for Mellin-Fejer Convolution Operators. Appl. Math. Lett., 24, 2064-2067.
- [2] Bardaro, C., Mantellini, I. (2010). A Quantitative Voronovskaya Formula for Mellin Convolution Operators. Mediterr. J. Math., 7, 483-501.
- [3] Bardaro, C., Mantellini, I. (2011). Asymptotic Behaviour of Mellin-Fejer Convolution Operators. East J. Approx., 17, 181-201.
- [4] Bardaro, C., Mantellini, I. (2007). Voronovskaya-type Estimates for Mellin Convolution Operators. Res. Math., 50, 1-16.
- [5] Butzer, P. L., Jansche, S. (1997). A Direct Approach to the Mellin Transform. J. Fourier Anal. Appl., 3, 325-375.
- [6] Devore, R. A., Lorentz, G. G. (1993). Constructive Approximation, Grundlehren der Mathematischen Wissenschaften. p. 303, Springer, Berlin.
- [7] Kolbe, W., Nessel, R. J. (1972). Saturation Theory in Connection with Mellin Transform Methods. SIAM J. Math. Anal., 3, 246-262.
- [8] Mamedov, R. G. (1991). The Mellin Transform and Approximation Theory. Elm, Baku, (in Russian).
- [9] Zemanian, A. H. (1968). Generalized Integral Transformation. Interscience, New York.

3rd International Conference: Constructive Mathematical Analysis

Modular Fixed Point Framework for Seljuk Knot Motifs: A Topological Study of Karatay Madrasa and Kubadabad Palace

Gülşen Tezcan Kaya¹, Mahpeyker Öztürk², Hatice Parlatici³ and Soley Ersoy²

¹ Department of History of Arts, Sakarya University, Sakarya, Türkiye

² Department of Mathematics, Sakarya University, Sakarya, Türkiye

³ Department of Library and Documentation, Sakarya University, Sakarya, Türkiye

gtezcan@sakarya.edu.tr, mahpeykero@sakarya.edu.tr, hparlati@sakarya.edu.tr, sersoy@sakarya.edu.tr

key-words: Kubadabad Palace, Karatay Madrasa, fixed point theorem, knot theory.

Abstract:

This study proposes a novel integration of fixed point theory and Seljuk architectural design by modelling geometric tile motifs using modular function spaces. Focusing on knot-based decorations from the Karatay Madrasa and Kubadabad Palace in Konya, we explore the applicability of modular topology in capturing the recursive and invariant nature of these motifs. Using various types of contractive mappings, we show the existence of fixed points corresponding to periodic and symmetric tile patterns. A comparative analysis between metric and modular spaces is provided to justify the use of modular functionals in modelling geometric features such as rotation invariance and curve embeddings. This interdisciplinary approach bridges cultural heritage and mathematical rigor, paving the way for digital reconstruction and pattern recognition in historical architecture.

References

- [1] Kauffman, L. H. (1987). State Models and the Jones Polynomial. *Topology*, 26(3), 395-407.
- [2] Grishanov, S. A., Meshkov, V. R. and Omelchenko, A. V. (2009). A Topological Study of Textile Structures: Topological Invariants in Application to Textile Structures. *Textile Research Journal*, 79(9), 822-836.
- [3] Banach, S. (1922). Sur Les Operations Dans Les Ensembles Abstraits et Leurs Applications Aux Equations Integrales. *Fund. Math.*, 1, 133-181.
- [4] Chistyakov, V. V. (1969). A Theorem on Fixed Points in Modular Spaces and Its Application to the Theory of Differential Equations. *Mathematical Notes*, 5(2), 339-344.
- [5] Cakmakoglu Kuru, A. (2008). Orta Çağ Anadolu Türk Mimarisinde Dugum Motifi ve İkonografisi. *Erdem*, 51, 23-47.

3rd International Conference: Constructive Mathematical Analysis

On Parabolic Class Number of a Special Subgroup of the Modular Group

Seda Öztürk

¹ Department of Mathematics, Karadeniz Technical University, Trabzon, Türkiye

seda.ozturk@ktu.edu.tr

key-words: Congruence subgroups, parabolic class number, Fuchsian groups.

Abstract:

In this study, we investigate a constant of the signature of a special subgroup G_p of the modular group Γ , defined as

$$G_p = \left\{ \begin{pmatrix} a & b \\ cp^\alpha & d \end{pmatrix} \in \Gamma : a \equiv d \pmod{p} \right\}$$

where p is a prime number and α is any natural number. The main focus is the determination of the number of parabolic conjugacy classes, which correspond to the cusps of the modular surface associated with this group. Using group-theoretic and arithmetic methods, we explicitly compute the parabolic class number of G_p . The results contribute to a deeper understanding of the structural and geometric properties of modular subgroups beyond the classical congruence cases.

References

- [1] Akbaş, M. (1989). The Normalizer of Modular Subgroups. Ph. D. Thesis, Southampton University.
- [2] Akbaş, M., Singerman, D. (1992). The Signature of the Normalizer of $\Gamma_0(N)$. London Math. Soc. Lecture Note Ser., Cambridge Univ. Press, Cambridge, 165, 77-86.
- [3] Jones, G. A., Singerman, D. (1987). Complex Functions: An Algebraic and Geometric Viewpoint. Cambridge Univ. Press, Cambridge.
- [4] Jones, G. A., Singerman, D. and Wicks, K. (1991). The Modular Group and Generalized Farey Graphs. LMS Lect. Note Ser., 160, 316-338.
- [5] Keskin, R. (1998). On the Parabolic Class Number of Some Subgroups of Hecke Groups. Tr. J. of Mathematics, 22, 199-205.
- [6] Keskin, R. (1999). On the Parabolic Class Number of Some Fuchsian Groups. Note di Matematica, 19, 275-283.
- [7] Keskin, R. (2005). A Note on Some Modular Subgroups. Rocky Mountain Journal of Mathematics, 35, 573-585.
- [8] Keskin, R. (2012). Parabolic Class Number of a Subgroup of the Normalizer of $\Gamma_0(m)$. Miskolc Mathematical Notes 2, 13, 365-373.
- [9] Lang, M. L. (2001). The Signature of $\Gamma_0^+(N)$. J. Algebra, 241, 146-185.
- [10] Lehner, J. (1964). Discontinuous Groups and Automorphic Functions, Math. Surveys 8. Amer. Math. Soc.
- [11] Maclachlan, C. (1981). Groups of Units of Zero Ternary Quadratic Forms. Proc. R. Soc. Edinb., Sect. A, 88, 141-157.
- [12] Schoeneberg, B. (1974). Elliptic Modular Functions. Springer Verlag, New York.
- [13] Shimura, G. (1971). Introduction to the Arithmetic Theory of Automorphic Functions. Princeton Univ. Press.
- [14] Wilkie, H. C. (1966). On Non-euclidean Crystallographic Groups. Math. Zeit., 91, 87-102.

3rd International Conference: Constructive Mathematical Analysis

A New Proof of the Girard-Fermat Theorem

Seda Öztürk

Department of Mathematics, Karadeniz Technical University, Trabzon, Türkiye

seda.ozturk@ktu.edu.tr

key-words: Prime numbers, Fermat, two squares theorem, congruence equations.

Abstract:

In this paper, we provide a novel and straightforward proof of the well-known Girard–Fermat Theorem, which asserts that every prime $p \equiv 1 \pmod{4}$ can be written as the sum of two squares. By utilizing a result derived from Wilson’s Theorem, we offer an elegant and simplified proof, which contrasts with more complex approaches traditionally used to establish this result in number theory.

References

- [1] Euler, L. (1758). De Numerus, Qui Sunt Aggregata Quorum Quadratorum, Novi Comment. Acad. Sci. Petropolitanae, 4(1752–3), 3–40.
- [2] Euler, L. (1760). Demonstration Theormatis Fermatiani Omnem Numerum Primum Formae $4n + 1$ Esse Summam Duorum Quadratorum, Novi Comment. Acad. Sci. Petropolitanae, (1754–5), 3–13.
- [3] Gauss, C. F. (1986). Disquisitiones Arithmeticae Arthur A. Clarke, Trans. Springer-Verlag.
- [4] Stillwell, J. 1996. Introduction to Theory of Algebraic Integers by Richard Dedekind. Cambridge Mathematical Library, Cambridge University Press.
- [5] Heath-Brown, D. R. (1984). Fermat’s Two-squares Theorem. Invariant, 3–5.
- [6] Zagier, D. (1990). A One-sentence Proof that Every Prime $p \equiv 1 \pmod{4}$ is a Sum of Two Squares. American Mathematical Monthly, 97:144.
- [7] Wells, D. (1990). Are These the Most Beautiful?. Math. Intelligencer, 12, 37–41.
- [8] Elsholtz, C. (2010). A Combinatorial Approach to Sums of Two Squares and Related Problems. In: Chudnovsky, D., Chudnovsky, G. (eds) Additive Number Theory, New York, NY: Springer, 115–140.
- [9] Christopher, A. D. (2016). A Partition-theoretic Proof of Fermat’s Two Squares Theorem. Discrete Mathematics, 339, 1410–1411.
- [10] Sam, N. (2020). A Short Proof of Fermat’s Two-square Theorem, The American Mathematical Monthly, 7:127, 638–638.
- [11] Niven, I., Zuckerman, H. S. (1974). An Introduction to the Theory of Numbers. John Wiley and Sons Inc., New York.
- [12] Alaca, Ş., Williams, K. S. (2004). Introductory Algebraic Number Theory, Cambridge University Press.
- [13] Williams, K. S. (1985). Heath-Brown’s Elementary Proof of the Girard-Fermat Theorem, Carleton Coordinates, Carleton University, Ottawa, Ontario, Canada, January, 4–5.
- [14] Varouchas, Y. (1984). Une démonstration élémentaire du théorème des deux carrés. La Caverne, I.R.E.M. de Lorraine, France, Bulletin No.6, février, 31–39.

3rd International Conference: Constructive Mathematical Analysis

Mappings Contracting Triangles

Cristina Maria Păcurar¹, Ovidiu Popescu¹

¹Faculty of Mathematics and Computer Science, Transilvania University of Braşov, Braşov, Romania

cristina.pacurar@unitbv.ro

key-words: Metric space, fixed point theorem, three-valued mappings, contracting triangles.

Abstract:

The aim of the current paper is to introduce a new class of contractive mappings, which are contracting (a feature of) triangles and prove the fixed point result for such mappings. We emphasize that our main theorem encompasses many functions for which the result holds, thus being a significant improvement in the research field.

References

- [1] Banach, S. (1922). Sur les operations dans les ensembles abstraits et leur application aux equations integrales. *Fundamenta Mathematicae*, 3, 133-181.
- [2] Chatterjea, S. K. (1972). Fixed-point Theorems, *C.R. Acad. Bulgare Sci.*, 25, 727-730.
- [3] Jleli, M., Samet, B. (2014). A New Generalization of the Banach Contraction Principle. *J. Inequal. Appl.* 2014, 38.
- [4] Kannan, R. (1969). Some Results on Fixed Point — II. *Amer. Math. Monthly*, 76, 405-408.
- [5] Pata, V. (2019). *Fixed Point Theorems and Applications*. Springer, Cham.
- [6] Petrov, E. (2023). Fixed Point Theorem for Mappings Contracting Perimeters of Triangles. *J. Fixed Point Theory Appl.*, 25, 1-11.
- [7] Petrov, E., Bisht, R. K. Fixed Point Theorem for Generalized Kannan Type Mappings. *arXiv:2308.05419*.
- [8] Petrov, E. Periodic Points of Mappings Contracting Total Pairwise Distance. *arXiv:2402.02536.10*.
- [9] Popescu, O. (2021). Some Remarks on the Paper Fixed Point Theorems for Generalized Contractive Mappings in Metric Spaces. *J. Fixed Point Theory Appl.*, 23, 1-10.

3rd International Conference: Constructive Mathematical Analysis

Properties of Hyperbolic Convolution on the Unit Disc

Tumea Eisner¹, Margit Pap¹

¹ Faculty of Sciences, University of Pécs, 7624 Pécs, Hungary

eisner@ttk.pte.hu, papm@gamma.ttk.pte.hu

key-words: Hyperbolic convolution, approximate identity on the unit disc.

Abstract:

The Blaschke group was introduced by Pap and Schipp in [1, 2]. Using the representations of the Blaschke group, it was possible to generate hyperbolic wavelet transforms and multiresolution in the analytic function spaces (see [3, 4, 5, 6, 7, 8]). A subgroup of the Blaschke group induces group operation on the unit disc. Recently, Schipp introduced a new convolution operator in this group [9]. In this paper, we study further properties of the hyperbolic convolution over the unit disc, and we derive approximation identity results on the unit disc.

References

- [1] Pap, M., Schipp, F. (2006). The Voice Transform on the Blaschke Group I. PU.M.A., 17(3-4), 387-395.
- [2] Pap, M., Schipp, F. (2008). The Voice Transform on the Blaschke group II. Annales Univ. Sci. Budapest., Sect. Comput., 29, 157-173.
- [3] Pap, M., Schipp, F. (2010). The Voice Transform on the Blaschke Group III. Publ. Math. Debrecen, 75(1-2), 263-283.
- [4] Pap, M. (2010). The Voice Transform Generated by a Representation of the Blaschke Group on the Weighted Bergman Spaces. Annales Univ. Sci. Budapest., Sect. Comput., 33, 321-342.
- [5] Pap, M. (2011). Hyperbolic Wavelets and Multiresolution in $H^2(\mathbb{T})$. J. Fourier Anal. Appl., 17, 755-776.
- [6] Pap, M. (2011). Properties of the Voice Transform of the Blaschke Group and Connections with Atomic Decomposition Results in the Weighted Bergman Spaces. J. Math. Anal. Appl., 389(1), 340-350.
- [7] Pap, M. (2015). A Special Voice Transform, Analytic Wavelets, and Zernike Functions, In: Peter W Hawkes (Ed.) Advances in Imaging and Electron Physics, New York: Elsevier, Volume 188, 79-134.
- [8] Pap, M. (2019). Hyperbolic Wavelet Frames and Multiresolution in the Weighted Bergman Spaces. In: Paolo, Boggiatto; Elena, Cordero; Maurice, de Gosson; Hans, G Feichtinger; Fabio, Nicola; Alessandro, Oliaro; Anita, Tabacco - Landscapes of Time-Frequency Analysis, Basel : Birkhauser, Basel, 225-247.
- [9] Schipp, F. (2023). Convolution Operators on the Disc. Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae, Sectio Computatorica, 54, 281-293.

3rd International Conference: Constructive Mathematical Analysis

Optimal Recovery of Mappings Based on Linear Information with the help of T -splines in Banach Spaces

Vladyslav Babenko¹, Yuliya Babenko² and Nataliia Parfinovych¹

¹Oles Honchar Dnipro National University, Dnipro, Ukraine

²Kennesaw State University, Marietta, GA, USA

`babenko.vladislav@gmail.com, ybabenko@kennesaw.edu, parfinovich@mmf.dnu.edu.ua`

key-words: Optimal recovery, information operators, abstract splines.

Abstract:

Let H, X, Y be real Banach spaces, $\|\cdot\|_X$ be a norm in X , M be some subset of H , $\mathcal{L}(X, Y)$ be the space of linear bounded operators $A: X \rightarrow Y$, X^* be the space dual to X , A^* be the adjoint operator for A . By $\mathcal{R}(A)$ we denote the range of the operator A , and by $\mathcal{N}(A)$ we denote the kernel of A . The values of the functional $f \in X^*$ on $x \in X$ we denote as follows $\langle x, f \rangle_X$. For a given operator $A \in \mathcal{L}(X, Y)$ we set $\mathcal{N}(A)^\perp = \{f \in X^*: \langle x, f \rangle_X = 0 \quad \forall x \in \mathcal{N}(A)\}$. As it is well known, $\mathcal{N}(A)^\perp = \overline{\mathcal{R}(A^*)}$, where $\overline{Q}$ is the closure of the set Q .

We define the best approximation of a class $A(M)$ by a subspace F in space X by the following

$$E(A(M); F; X) := \sup_{x \in A(M)} \inf_{u \in F} \|x - u\|_X.$$

Set

$$P_F x = \{u \in F: \|x - u\|_X = E(x; F; X)\}$$

is called a metric projection of element x onto the set F . If for any $x \in X$ set $P_F x$ contains exactly one element, then the set F is called a Chebyshev set.

Let $T \in \mathcal{L}(H, Y)$ be an information operator. Elements of the manifold

$$S(T) = S_H(T) := \{s_h = h - P_{\mathcal{N}(T)} h: h \in H\}$$

are called T -splines in the space H .

We say that T -spline $s_h \in S(T)$ is T -interpolating for an element $h \in H$ if $Ts_h = Th$.

We prove that for an arbitrary $h \in H$ there exists a unique T -interpolating spline s_h . We also obtain a representation of such a spline via Th and establish a minimization property of T -interpolating splines. Suggested definition of $E(A(M); F; X)$ is aimed at studying extremal problems of approximation theory on classes $A(M)$ of elements of a Banach space X , given as an image of a set $M \subset H$ under mapping $A: H \rightarrow X$ (not necessarily linear). Many important classes of periodic and nonperiodic univariate and multivariate functions, in particular classes of solutions to various problems of mathematical physics, can be represented in such a way.

We also use T -interpolating splines to solve the problem of optimal recovery of classes $A(M)$ in space X based on available information on elements of set M , given with the help of information operator T .

3rd International Conference: Constructive Mathematical Analysis

Some Fixed Point Theorems in Modular b -Metric-Like Spaces

Nizamettin Ufuk Bostan¹, Banu Pazar Varol¹

¹Department of Mathematics, Kocaeli University, Kocaeli, Türkiye

ufuk.bostan@kocaeli.edu.tr, banupazar@kocaeli.edu.tr

key-words: b -metric-like spaces, modular b -metric-like spaces, fixed point theory, contraction principle.

Abstract:

The aim of this paper is to contribute to the fixed point theory by proving some fixed point theorems in modular b -metric-like space. The existence and uniqueness of fixed point in the modular b -metric-like space under different contraction conditions are handled.

References

- [1] Alghamdi, M. A., Hussain, N. and Salimi, P. (2013). Fixed Point and Coupled Fixed Point Theorems on b -metric-like Spaces. *J. Inequalities Appl.*, 2013, 402.
- [2] Amini-Harandi, A. (2012). Metric-like Spaces, Partial Metric Spaces and Fixed Points. *Fixed Point Theory Appl.*, 2012, 204.
- [3] Bostan, N. U., Pazar Varol, B. (2024). Fixed Point Results in Modular b -Metric-like Spaces with an Application. *Axioms*, 13(10), 726.
- [4] Chistyakov, V. V. (2010). Modular Metric Spaces, I: Basic concepts. *Nonlinear Anal.*, 72, 1-14.
- [5] Czerwik, S. (1993). Contraction Mappings in b -metric Spaces. *Acta Math. Inform. Univ. Ostrav.*, 1, 5-11.
- [6] Ege, M. E., Alaca, C. (2018). Some Results for Modular b -metric Spaces and an Application to System of Linear Equations. *Azerbaijan J. Math.*, 8, 3-14.
- [7] Girgin, E., Buyukkaya, A., Kuru, N. K. and Ozturk, M. (2024). On the Impact of Some Fixed Point Theorems on Dynamic Programming and RLC Circuit Models in R -Modular b -Metric-like Spaces. *Axioms*, 13(7), 441.
- [8] Rasham, T., Shoaib, A., Park, C., De La Sen, M., Aydi, H. and Lee, J. R. (2020). Multivalued Fixed Point Results for Two Families of Mappings in Modular-like Metric Spaces with Applications. *Complexity*, 2020, 1-10.

3rd International Conference: Constructive Mathematical Analysis

Recent Advancements on Semi-Discrete Sampling Operators in Sobolev-Orlicz Spaces

Michele Piconi¹, Gianluca Vinti¹

¹Department of Mathematics and Computer Science, University of Perugia, Perugia, Italy

michele.piconi@unipg.it, gianluca.vinti@unipg.it

key-words: Durrmeyer sampling operators, simultaneous approximation, Sobolev-Orlicz spaces, maximal operators.

Abstract:

In this talk, we present some recent approximation results for a semi-discrete family of sampling operators, called Durrmeyer Sampling Operators (DSO) [3], in Sobolev-Orlicz spaces [4]. This research is part of the theory of sampling series, which plays a key role in Approximation Theory and its applications, especially in Signal and Image Processing. DSO extend classical families such as generalized [2] and Kantorovich [1] sampling operators, being based on two different kernels: a discrete and a continuous one.

We present new convergence results for Durrmeyer-type sampling operators (DSO) in Sobolev-Orlicz spaces, focusing on the simultaneous approximation of functions and their weak derivatives [6]. These results are based on a density method and on the boundedness of maximal operators in Orlicz spaces [7, 5].

We also study a direct approach involving φ -functions that do not satisfy the Δ_2 -condition. This leads to convergence results with order of approximation in more general functional spaces, both in the sense of the Luxemburg norm and modular convergence.

Several examples are discussed, including Sobolev-Orlicz versions of Zygmund and exponential-type spaces, as well as classical Sobolev spaces.

References

- [1] Bardaro, C., Butzer, P.L., Stens, R.L. and Vinti, G. (2007). Kantorovich-Type Generalized Sampling Series in the Setting of Orlicz Spaces. *Sampling Theory in Signal and Image Processing*, 9(6), 29-52.
- [2] Butzer, P. L., Fisher, A. and Stens, R. L. (1987). Approximation of Continuous and Discontinuous Functions by Generalized Sampling Series. *Journal of Approximation Theory*, 50, 25-39.
- [3] Costarelli, D., Piconi, M. and Vinti, G. (2025). On the Regularization by Durrmeyer-sampling Type Operators in L^p -spaces via a Distributional Approach. *Journal of Fourier Analysis and Applications*, 31, Article 11.
- [4] Donaldson, T.K., Trudinger, N.S. (1971). Orlicz-Sobolev Spaces and Imbedding Theorems. *Journal of Functional Analysis*, 8, 52-75.
- [5] Harjulehto, P., Hästö, P. (2019). Orlicz Spaces and Generalized Orlicz Spaces. *Lecture Notes in Mathematics*, vol. 2236. Springer.
- [6] Piconi, M., Vinti, G. (2025). Semi-discrete Sampling in Sobolev-Orlicz Spaces. To be submitted.
- [7] Stein, E.M. (1970). *Singular Integrals and Differentiability Properties of Functions*. Princeton University Press.

3rd International Conference: Constructive Mathematical Analysis

A Study of Fixed Points in Soft S-Metric Spaces via Hybrid-Interpolative Reich-Istrecescu Type Mappings

Hande Poşul¹, Çigdem Aras Gunduz², Arzu Erdem Coşkun² and Sadi Bayramov³

¹Department of Mathematics, Kilis 7 Aralık University, Kilis, Türkiye

²Department of Mathematics, Kocaeli University, Kocaeli, Türkiye

³Department of Algebra and Geometry, Baku State University, Baku, Azerbaijan

handeposul@gmail.com, caras@kocaeli.edu.tr, aerdem@kocaeli.edu.tr, baysadi@gmail.com

Abstract:

Classical fixed point theory has long been one of the foundational pillars of mathematical analysis and topology, with significant developments emerging over the past century. Early groundbreaking contributions include Brouwer's Fixed Point Theorem (1911) and Banach's Contraction Principle (1922), which have played a critical role in shaping both theoretical and applied mathematics. In particular, Banach's theorem guarantees the existence and uniqueness of fixed points for contraction mappings on complete metric spaces, providing a robust framework widely applicable in nonlinear analysis, differential equations, and numerical methods.

Over time, the limitations of classical fixed point theorems within standard metric spaces have become apparent, motivating the development of generalized metric structures. As a result, new concepts such as G-metric, b-metric, and S-metric spaces have been introduced. Among these, S-metric spaces, initially defined by Sedghi et al.[5], extend the classical notion of metric spaces by utilizing a three-variable distance function, thereby offering a more flexible analytical environment. These spaces have opened new avenues for exploring fixed point results under weaker or more generalized conditions.

Currently, three primary approaches are predominant in the generalization of fixed point theory:

- generalizing the contraction conditions,
- extending the underlying metric structure, and
- investigating fixed points in terms of geometric constructs.

This study employs a combination of the first two approaches by both introducing a new contraction framework and working within a broader metric structure, namely soft S-metric spaces.

Soft set theory, proposed by Molodtsov in 1999 [4], emerged as a powerful mathematical tool for handling uncertainty and imprecise information. When integrated with metric concepts, it has led to the development of soft metric spaces, and more recently, soft S-metric spaces, which offer an enriched setting for analyzing fixed point problems in uncertain environments [1, 2]. These structures are particularly valuable in fields such as decision-making, artificial intelligence, and systems analysis.

In this study, we adopt the hybrid-interpolative Reich-Istrecescu type contraction, introduced by Karapınar et al. [3], and adapt it to the context of soft S-metric spaces. This generalized contraction framework, characterized by a hybrid and parameterized structure, broadens the classical Reich-Istrecescu model and facilitates the derivation of new fixed point theorems. Within this extended setting, we establish both existence and uniqueness results, further supported by illustrative examples.

Therefore, this work contributes not only to the generalization of fixed point results but also demonstrates the effective integration of soft structures into the realm of metric fixed point theory. In doing so, it expands the theoretical scope and application potential of fixed point theorems in more complex and uncertain systems.

References

- [1] Gündüz Aras Ç., Bayramov S. and Cafarli V. (2018). Fixed Point Theorems on Soft S-Metric Spaces, Communications in Mathematics and Applications, 9(4), 725-735.
- [2] Gündüz Aras Ç., Bayramov S. and Coşkun Erdem A. (2024). Fixed Point Theorems on Parametric Soft S-metric Spaces, Filomat, 38:19, 6753-6762.
- [3] Karapınar E., Fulga A., Shahzad N. and Roldon Lopez de Hierro A. F. (2022). Solving Integral Equations by means of Fixed Point Theory, Journal of Function Spaces, 2022(1); Article ID: 7667499.
- [4] Molodtsov, D. (1999). Soft Set Theory-first Results. Computers and Mathematics with Applications, 37(4-5), 19-31.
- [5] Sedghi S., Shobe N. and Aliouche A., (2012). A Generalization of Fixed Point Theorem in S-metric spaces. Matematiski Vesnik, 64, 258-266.

3rd International Conference: Constructive Mathematical Analysis

Normal Structure and Contractions which Diminish Radius in Metric and Banach Spaces

Abdelkader Dehici¹, Najah Redjel¹ and Sami Atailia²

¹Department of mathematics, University of Souk Ahras, Souk Ahras, Algeria

²Department of mathematics, University of Bouverdes, Bouverdes, Algeria

dehikader@yahoo.fr, N.Radjal@univ-soukahras.dz, s.atailia@univ-boumerdes.dz

key-words: Metric space, admissible subset, hyperconvex metric space, mappings diminishing the radius, orbitally wrt nonexpansive mapping, orbitally Kannan mapping, weak normal structure, weak quasi-normal structure, convex invariant subset, fixed point.

Abstract:

in this talk, by using admissible sets, we give some fixed point results for orbitally contractions which diminish the radius of invariant convex subsets and orbits. Furthermore, a characterization of the weak normal structure by the fixed point property associated with this class of mappings is established. .

References

- [1] Amini-Harandi, A., Fakhar, A. and Hajisharifi, H. R. (2016). Weak Fixed Point Property for Nonexpansive Mappings with respect to Orbits. J. Fixed. Point. Theory. Appl., 16, 601-607
- [2] Kirk, W. A., Shahzad, N. (2018). Normal Structure and Orbital Fixed Point Conditions. J. Math. Anal. Appl., 363, 461-476.
- [3] Wong, C. S. (1975). On Kannan Maps. Proc. Amer. Math. Soc., 47, 105-111.

3rd International Conference: Constructive Mathematical Analysis

Sampling Type Operators Versus AI in Medical Image Processing

Lucrezia Rinelli^{1,2}, Arianna Travaglini³ and Gianluca Vinti²

¹ Department of Mathematics and Computer Science “U.Dini” - DIMAI, University of Florence, Italy

² Department of Mathematics and Computer Science, University of Perugia, Italy

³ Department of Engineering, University of Basilicata, Italy

lucrezia.rinelli@unifi.it, arianna.travaglini@unibas.it, gianluca.vinti@unipg.it

key-words: Sampling Kantorovich operators, segmentation algorithm, image processing, U-Net neural network.

Abstract:

In this talk we will delve into recent outcomes highlighted in [4], focusing on the application of sampling-type operators in digital image processing. A key role in this work is undertaken by sampling Kantorovich (SK) operators ([1, 4]), widely appreciated for their accuracy in the reconstruction of signals that are not necessarily continuous. Building upon significant theoretical findings, our study employs these operators within a deterministic method grounded on a segmentation algorithm, in which the SK assume a pivotal role ([2, 5]). This mathematically-based approach is compared with an AI method that involves a U-Net neural network. Both approaches are aimed at segmenting the patent area of the aortic vessel from CT images of patients affected by abdominal aortic aneurysm, proposing innovative and less invasive alternatives to nephrotoxic contrast agents commonly employed in diagnosing this pathology. The numerical and visual assessment of the outcomes highlights the efficiency and reliability of both methods.

References

- [1] Bardaro, C., Butzer, P. L., Stens, R. L. and Vinti, G. (2007). Kantorovich-Type Generalized Sampling Series in the Setting of Orlicz Spaces. *Sampling Theory in Signal and Image Processing*, 6(1), 29-52.
- [2] Costarelli, D., Seracini, M. and Vinti, G. (2020). A Segmentation Procedure of the Pervious Area of the Aorta Artery from CT Images without Contrast Medium. *Mathematical Methods in the Applied Sciences*, 43, 114-133.
- [3] Costarelli, D., Vinti, G. (2011). Approximation by Multivariate Generalized Sampling Kantorovich Operators in the Setting of Orlicz Spaces. *Bollettino dell'unione Matematica Italiana*, 4(3), 445-468.
- [4] Rinelli, L., Travaglini, A., Vescera, N. and Vinti, G. (2024). An Approximation-based Approach Versus an AI One for the Study of CT Images of Abdominal Aorta Aneurysms, submitted, arXiv: 2406.01764.
- [5] Travaglini, A., Vinti, G., Scalera, G. B and Scialpi, M. (2023). A Large Scale Analysis for Testing a Mathematical Model for the Study of Vascular Pathologies. *Mathematics*, 11(8), 1-19.

3rd International Conference: Constructive Mathematical Analysis

Respecting Cauchy Equivalence Classes

Anthony J. Ripa

Computer Science, Stony Brook University, Stony Brook, USA

aripa@cs.stonybrook.edu

key-words: Cauchy sequences, equivalence classes, continuity, removable discontinuities.

Abstract:

In the standard construction of real numbers as Cauchy equivalence classes of rational sequences, only the basic arithmetic operations are defined to respect these classes, leaving other functions prone to discontinuities or undefined points that necessitate limits. This reliance on limits reintroduces complexity into a system initially built on convergence, undermining the elegance of the Cauchy construction. This paper proposes a unified approach where all functions respect Cauchy equivalence classes, ensuring they are well-defined at all points without redundant limits. By applying rational functions to sequence representatives and assigning the resulting equivalence class, we eliminate pathologies such as removable discontinuities. Examples, including difference quotients and Riemann sums, demonstrate the method's efficacy. Our approach offers simplicity and coherence, suggesting a principled construction of real analysis.

Notation. By default, an unbracketed 1 is a rational number, and bracketed [1] is real.

Method. A Cauchy sequence $(q_n)_{n \in \mathbb{N}}$ in $\mathbb{Q}$ satisfies $\forall \epsilon > 0, \exists N : m, n > N \implies |q_m - q_n| < \epsilon$. Two sequences (q_n) and (r_n) are equivalent if $\forall \epsilon > 0, \exists N : n > N \implies |q_n - r_n| < \epsilon$, and $\mathbb{R}$ is the quotient set.

For a rational partial function $f : \mathbb{Q}^k \rightarrow \mathbb{Q}$, we define its *Cauchy lift* ${}^c f : \mathbb{R}^k \rightarrow \mathbb{R}$ s.t. if $(x_1, \dots, x_k) \in \mathbb{R}^k$ and $Y \in \mathbb{R}$ then ${}^c f(x_1, \dots, x_k) = Y$ if for all $(q_{1,n}) \in x_1, \dots, (q_{k,n}) \in x_k$, if $f(q_{1,n}, \dots, q_{k,n})$ is defined, then $f(q_{1,n}, \dots, q_{k,n}) \in Y$.

In other words, if f is undefined for some representatives, ${}^c f$ remains well-defined as long as the defined cases consistently yield sequences in Y . This leverages the equivalence class structure to extend f naturally across $\mathbb{R}$, avoiding artificial undefinedness.

Example 1. We construct ${}^c \sqrt{X}$ from the rational $\sqrt{x}$. For $X \in \mathbb{R}$, take $x = \{c_1, c_2, c_3, \dots\} \in X$, and define $\sqrt{x} = \{\sqrt{c_1}, \sqrt{c_2}, \sqrt{c_3}, \dots\}$ when each c_i is a rational perfect square. We require a $Y \in \mathbb{R}$ such that if $\sqrt{x}$ is defined, then $\sqrt{x} \in Y$. For $X = [2]$, take $x = \{1^2, 1.4^2, 1.41^2, 1.414^2, \dots\}$, so $\sqrt{x} = \{1, 1.4, 1.41, 1.414, \dots\} \in [\sqrt{2}]$. Similarly, sequences like $\{3, 3.9, 3.99, \dots\} \in [4]$ may not have defined square roots pointwise, but whenever defined (e.g., perfect squares approximating 4), the result is in $[2]$. Thus, ${}^c \sqrt{X} = Y$, is well-defined for all $X \geq 0$.

Example 2. To find the derivative of $f(x) = x^2$ at $x = 1$, consider the rational function $g(h) = \frac{(1+h)^2 - 1^2}{h} = 2 + h$, defined for $h \neq 0$. Construct ${}^c g(H)$ for $H \in \mathbb{R}$. Take $h = \{c_1, c_2, c_3, \dots\} \in H$ with all $c_i \neq 0$, so $g(h) = \{2 + c_1, 2 + c_2, 2 + c_3, \dots\} \in [2 + H]$. For $H = 0$, use $h = \{\frac{1}{1}, \frac{1}{2}, \frac{1}{3}, \dots\}$, yielding $\{2 + 1, 2 + \frac{1}{2}, 2 + \frac{1}{3}, \dots\} \in [2]$. So, ${}^c g(H) = 2 + H$. Thus, ${}^c g(0) = 2$, the derivative $f'(1)$, directly obtained, without redundant limits.

Example 3. For $g(x) = x$ from 0 to 1, define the rational Riemann Sum function $f(h) = h \cdot \sum_{i=1}^{1/h} i h$. Construct ${}^c f(H)$ at $H = 0$ with $h = \{1, \frac{1}{2}, \frac{1}{3}, \dots\} \in 0$, where $1/h_n = n$ is an integer. This simplifies $f(h) = h^2 \cdot \frac{1}{h} \cdot \frac{(\frac{1}{h} + 1)}{2} = \frac{1+h}{2}$. Then, $f(h) = \{\frac{1+1}{2}, \frac{1+\frac{1}{2}}{2}, \frac{1+\frac{1}{3}}{2}, \dots\} = \{1, \frac{3}{4}, \frac{2}{3}, \dots\} \in [\frac{1}{2}]$. Thus, ${}^c f(0) = \frac{1}{2}$, the integral $\int_0^1 x dx$, directly obtained, again without redundant limits.

Summary. By mandating that all functions respect Cauchy equivalence classes, we redefine real analysis to be more coherent and intuitive, avoiding redundant limits and pathologies inherent in the classical approach. Examples confirm practicality for diverse functions. Future work could extend to other quotient constructions, potentially reinterpreting mathematical analysis.

3rd International Conference: Constructive Mathematical Analysis

S_{pmk}^* Index for the Burr X-Perks Distribution with Applications

Şule Sağlam¹ and Kadir Karakaya¹

¹ Department of Statistics, Faculty of Science, Selcuk University, Konya, Türkiye

sulesaglam75@gmail.com, kkarakaya@selcuk.edu.tr

key-words: Statistical process control, Burr X-Perks distribution, maximum likelihood estimation, interval estimation, process capability index, symmetric and asymmetric loss function.

Abstract:

Process capability indices are widely employed in quality control practices to quantitatively assess the extent to which manufacturing processes meet customer requirements. These indices provide comprehensive information about a process's performance and quality level by evaluating whether the process outputs fall within predefined specification limits. Selecting an appropriate process capability index based on the distributional characteristics of the observed data is crucial for ensuring an accurate assessment of the process. This study proposes a novel process capability index that incorporates both symmetric and asymmetric loss functions. The properties of the proposed index are thoroughly examined under the assumption that the process data follow a Burr X-Perks distribution. The point and interval estimators of the proposed index are developed using the maximum likelihood methodology, and their performance is evaluated through an extensive Monte Carlo simulation study. Furthermore, a real data application is presented to demonstrate the practical applicability of the proposed index.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Terna, G. I., and Adamu, A. U. A. (2025). Burr X-Perks Distribution: Properties and Applications. In Royal Statistical Society Nigeria Local Group Annual Conference Proceedings (pp. 466-488).

3rd International Conference: Constructive Mathematical Analysis

Variable-Order Hadamard-Type Fractional Calculus with respect to Function

Ibrahim Abdullahi Saleh¹, Zhivorad Tomovski²

¹Department of Mathematics, University of Ostrava, Ostrava, Czech Republic

²Department of Mathematics, University of Olomouc, Olomouc, Czech Republic

ibrahim.saleh.s01@osu.cz, Zhivorad.tomovski@upol.cz

key-words: Fractional Calculus, Ψ -Hadamard-type fractional integral of variable order (FIVO), Ψ -Hadamard-type fractional Derivative of variable order (FDVO), existence and uniqueness results.

Abstract:

In this paper, We introduce the Ψ -Hadamard-type fractional integral of variable order (FIVO). We propose two variants of the Ψ -Hadamard-type fractional derivative of variable order (FIVO) in the Riemann–Liouville sense and three variants in the Caputo sense. We study their relevant properties, established their interconnections and rigorously prove the relationship between these operators. In addition, We drive existence and uniqueness results for fractional differential equations involving the Riemann–Liouville Ψ -Hadamard-type fractional derivative operator of constant order by applying fixed-point theorems. Finally, we illustrate the theorem with examples.

References

- [1] Fahad, H. M., Fernandez, A., Rehman, M. U. and Siddiqi, M. (2021). Tempered and Hadamard-type Fractional Calculus with respect to Functions. *Mediterranean Journal*.
- [2] Almeida, R. (2024). On the Variable-order Fractional Derivatives with respect to Another Function. *Aequationes mathematicae*, 1-18.
- [3] Sousa J. V. D. C. , Machado J. T. and De Oliveira E. C. (2020). The Ψ -Hilfer Fractional Calculus of Variable Order and Its Applications. *Computational and Applied Mathematics*, 39(4), 296.
- [4] Almeida, R. (2017). Caputo–Hadamard Fractional Derivatives of Variable Order. *Numerical Functional Analysis and Optimization*, 38(1), 1-19.
- [5] Almeida, R., Tavares, D. and Torres, D. F. (2019). *The Variable-order Fractional Calculus of Variations*. Cham, Switzerland: Springer International Publishing.
- [6] Giusti, A., Colombaro, I., Garra, R., Garrappa, R. and Mentrelli, A. (2024). On Variable-order Fractional Linear Viscoelasticity. *Fractional Calculus and Applied Analysis*, 27(4), 1564-1578.
- [7] Albasheir, N. A., Alsinai, A., Niazi, A. U. K., Shafqat, R., Romana, Alhagyan, M. and Gargouri, A. (2023). A Theoretical Investigation of Caputo Variable Order Fractional Differential Equations: Existence, Uniqueness, and Stability Analysis. *Computational and Applied Mathematics*, 42(8), 367.

3rd International Conference: Constructive Mathematical Analysis

Adjoint Curve Using Conformable Derivative

Burcin Saltik Baek

Department of Mathematics, Erciyes University, Kayseri, Türkiye

burcinsaltik@erciyes.edu.tr

key-words: Conformable derivative, adjoint curve, Euclidean space, curvatures.

Abstract:

This study examines the geometric properties of curves in three-dimensional Euclidean space by conformable derivatives, which provide an improved correspondence to classical differentiation compared to conventional fractional derivatives. The emphasis is on the adjoint curve, which undergoes significant alterations when examined via conformable derivatives. We describe the conformable adjoint curve and examine its construction, the interrelations among curvatures, and the Frenet frame vectors. Furthermore, we examine the Darboux vector and the criteria under which the adjoint curve constitutes a general helix. We illustrate, via theoretical analysis and graphical representations, the influence of conformable derivatives on the shape of the adjoint curve.

References

- [1] Abdeljawad, T. (2015). On Conformable Fractional Calculus. *Journal of Computational and Applied Mathematics*, 279, 57-66.
- [2] Gozutok, U., Coban, H. and Sagioglu, Y. (2019). Frenet Frame with respect to Conformable Derivative. *Filomat*, 33(6).
- [3] Has, A., Yilmaz, B. (2022). Special Fractional Curve Pairs with Fractional Calculus. *International Electronic Journal of Geometry*, 15(1), 132-144.
- [4] Khalil, R., Al Horani, M., Yousef, A. and Sababheh, M. (2014). A New Definition of Fractional Derivative. *Journal of Computational and Applied Mathematics*, 264, 65-70.
- [5] Nurkan, S. K., Guven, O. A. and Karacan, M. K. (2019). Characterizations of Adjoint Curves in Euclidean 3-space. *Proceedings of the National Academy of Sciences, India Section A: Physical Sciences*, 89, 155-161.
- [6] Yajima, T., Oiwa, S. and Yamasaki, K. (2018). Geometry of Curves with Fractional-order Tangent Vector and Frenet-Serret Formulas. *Fractional Calculus and Applied Analysis*, 21(6), 1493-1505.

3rd International Conference: Constructive Mathematical Analysis

Structural Modeling of Covariate-Driven Misclassification Risk: A Likelihood-Based Approach

Sumeyra Sert¹, Coskun Kus¹

¹ Department of Statistics, Selcuk University, Konya, Türkiye

sumeyra.sert@selcuk.edu.tr, coskun@selcuk.edu.tr

key-words: Covariate effects, likelihood-based estimation, logit model, measurement error, misclassification probability.

Abstract:

This study presents a likelihood-based estimation framework developed to analyze how the risk of misclassification in quality control settings varies with measurement-related covariates. In industrial inspection systems, items are classified as conforming or nonconforming based on measured characteristics that are often subject to noise. When the precision of these measurements changes across operators, machines, or environmental conditions, the probability of incorrect classification becomes condition-dependent. The proposed model expresses this misclassification probability as a logistic function of covariates and incorporates it into the overall variance structure of the observed measurements via a numerically constructed inverse mapping. Simulation results demonstrate that the proposed method successfully recovers the covariate effects that drive misclassification and yields consistent parameter estimates under varying sample sizes. This approach provides a flexible and interpretable tool for identifying measurement conditions that impact classification reliability in quality assurance processes.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Doganaksoy, N. (2001). Assessment of Impact of Measurement Variability in the Presence of Multiple Sources of Product Variability. *Quality Engineering*, 13(1), 83-89.
- [2] Doganaksoy, N. (2021). A Simplified Formulation of Likelihood Ratio Confidence Intervals Using a Novel Property. *Technometrics*, 63(1), 127-135.
- [3] Montgomery, D. C. (2020). *Introduction to Statistical Quality Control*. John Wiley & Sons.

3rd International Conference: Constructive Mathematical Analysis

An Inverse Problem for Pseudoparabolic Equation with Memory Term and Damping

Shakir Aidos

Al-Farabi Kazakh National University, Almaty, Kazakhstan
Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan

ajdossakir@gmail.com

key-words: Inverse problem, nonlinear pseudoparabolic equation, memory term, solvability.

Abstract:

$$u_t - \kappa \Delta u_t - \lambda \Delta u - \int_0^t K(t-s) \Delta u(x, s) ds = \gamma |u|^{q-2} u + f(t) \cdot g(x, t), \text{ in } Q_T, \quad (1)$$

the initial condition

$$u(x, 0) = u_0(x) \text{ in } \Omega \quad (2)$$

the boundary condition

$$u(x, t) = 0 \text{ on } \Gamma_T, \quad (3)$$

and the integral overdetermination condition

$$\int_{\Omega} u(x, t) \omega(x) dx = h(t), \quad t \in [0, T]. \quad (4)$$

Here, the coefficient κ, λ are given positive numbers, γ is the coefficient of the damping term might be positive $\gamma > 0$ either negative $\gamma < 0$. The functions $g(x, t)$, $u_0(x)$, $\omega(x)$ and $h(t)$ are given. The exponent q is given positive number such that

$$1 < q < \infty. \quad (5)$$

The main goal of this work is to show existence and uniqueness of weak solutions to the inverse problem (1)-(4).

3rd International Conference: Constructive Mathematical Analysis

An Insight into Effects of Geography and Topography on the Urban Roadway Network Development of the Istanbul through Fractal Analysis

Recep Bilal Sikar¹, Niyazi Özgür Bezgin¹

¹ Department of Civil Engineering, Istanbul University-Cerrahpasa, Istanbul, Türkiye

recep.sikar@iuc.edu.tr, ozgur.bezgin@iuc.edu.tr

key-words: Fractal analysis, urban morphology, Istanbul, urban network analysis.

Abstract:

Urban roadway networks are complex systems shaped by various geographical and topographical factors. These elements influence not only the physical layout of roads but also their connectivity, accessibility, and overall efficiency. Fractal analysis, a mathematical approach to assess complex structures, has emerged as a valuable tool for understanding these networks' configurations and growth patterns. This research investigates and provides an insight through fractal analyses into how geography and topography can impact the development of urban roadway networks. Istanbul, which is a major metropolitan world city is composed of 39 districts. The study focuses on three of these districts that form the historical triad of Istanbul: Fatih, Beyoglu, and Uskudar. Separated by a unique waterway network composed of the Golden Horn, the Bosphorus Strait and the Marmara Sea, Istanbul's easterly, westerly and northerly expansion emanated from this peculiar district triad that form the historical core of Istanbul's transportation network.

The study uses Geographic Information System (GIS) tools to extract the roadway networks of the studied districts and Benoit Software to conduct fractal analyses on the roadway networks to calculate and compare fractal dimensions estimated by five different techniques referred to as 1. Box Counting (Db), 2. Perimeter Area (Dp), 3. Information (Di), 4. Mass (Dm), and 5. Ruler (Dr). The analysis will initially focus on the development of roadway networks within these three districts and subsequently extend the examination towards west from Fatih District, toward North from Beyoglu District and towards east from the Uskudar District.

This paper provides an insight into how fractal analysis can offer a strong method to represent the development of urban roadway networks in relation to urban topography and urban geography. Istanbul, as the subject of this paper, provides a valuable case study by consideration of its multitude of geographical features that are unique to this city. The study will discover ways to use fractal analyses to define connectivity and accessibility offered by urban roadway networks and discuss how such mathematical means of network representation can lead us to make informed choices in urban roadway network development.

References

- [1] Abid, R. I., Tortum, A. (2021). The Fractal Geometry of Turkey's Urban Transportation Networks. KSCE Journal of Civil Engineering, 25(4), 1455-1466.
- [2] Koprulu, B., Topcu, M. (2022). Kent Morfolojisinde Kentsel Doku Değişiminin Fraktal Analiz Yöntemi Kullanılarak Değerlendirilmesi: Konya Örneği. Türkiye Kentsel Morfoloji A-1, III. Kentsel Morfoloji Sempozyumu Bildiriler Kitabı, 949-981.

3rd International Conference: Constructive Mathematical Analysis

Controlling Periodic Limit Cycles of Math Anxiety

Sara D. Sony

Department of Mathematics and Statistics, Northwest Missouri State University, Maryville, USA

ssony@nwmissouri.edu

key-words: Limit cycles, math anxiety, epidemiological modeling.

Abstract:

Math anxiety among college students often displays cyclical fluctuations, particularly around assessment periods. In this study, we modeled the temporal dynamics of math anxiety using a compartmental framework adapted from mathematical epidemiology. By fitting this model to longitudinal data on students' self-reported anxiety levels, we characterized the emergence of periodic oscillations and identified key parameters influencing their amplitude and persistence. We then simulated targeted interventions such as supplemental academic support delivered prior to examinations to assess their impact on reducing anxiety peaks. Additionally, we derived the minimal intervention threshold necessary to disrupt sustained anxiety cycles. Given the socially transmissible nature of math anxiety, our findings suggest that strategies grounded in infectious disease modeling can inform effective approaches for its mitigation in educational settings.

3rd International Conference: Constructive Mathematical Analysis

Julia Sets-Geometric and Potential-Theoretic Insights

Przemysław Sprus

Doctoral School of Exact and Natural Sciences, Jagiellonian University, Krakow, Poland

p.sprus@doctoral.uj.edu.pl

key-words: Filled Julia sets, iterations, discs, Fatou sets, non-autonomous Julia sets, Green function.

Abstract:

We present some results concerning closed discs contained in the filled Julia sets associated with polynomials $f_c : \mathbb{C} \ni z \mapsto z^2 + c \in \mathbb{C}$ mostly for $c \in [-2, \frac{1}{4}]$ but also some other values. We investigate also a few non-autonomous Julia sets. As an application we mention that the pluricomplex Green function associated with the non-autonomous Julia set of the sequence $(f_{c_n})_{n=1}^\infty$ is Hölder continuous, provided $(c_n)_{n=1}^\infty \subset \overline{D}(0, \frac{1}{4})$.

References

- [1] Sprus, P. (2025). On Discs Contained in Filled Julia sets. Dolomites Research Notes on Approximation, 18, 27-40.

3rd International Conference: Constructive Mathematical Analysis

Evaluation of Lebesgue Constant with Admissible Meshes over Balls and Simplices

Mateusz Suder

Faculty of Mathematics and Computer Science, Jagiellonian University, Krakow, Poland

mateusz.suder@doctoral.uj.edu.pl

key-words: Lebesgue constant, admissible meshes.

Abstract:

We will present applications of admissible meshes for evaluating the Lebesgue constant and associated approximation errors for certain classes of polynomial approximation projections, including polynomial interpolation and weighted least squares. We will construct certain optimal admissible meshes over balls and simplices, derived from Chebyshev-type point sets, and give numerical comparisons with other known admissible meshes studied in recent years. The numerical experiments will be based on the covering radius with respect to the Dubiner distance. We will also discuss potential improvements to these constructions and open problems in this area.

The talk will be based on two papers [1, 2].

References

- [1] Białas-Cieź, L., Kenne, D. J., Sommariva, A. and Vianello, M. (2024). Evaluating Lebesgue Constants by Chebyshev Polynomial Meshes on Cube, Implex and Ball. *Electron. Trans. Numer. Anal.*, 60, 428-445
- [2] Białas-Cieź, L., Suder, M. (2024). Note on Admissible Meshes on Ball and Simplex via Dubiner Metric. *Dolomites Res. Notes Approx.*, 17, 89-96.

3rd International Conference: Constructive Mathematical Analysis

Exploring State-Measure Dynamics in Algebraic Logic Structures

Ibrahim Senturk¹, Abdullah Engin Calik², Huseyin Sirin² and Tahsin Oner¹

¹ Department of Mathematics, Ege University, Izmir, Türkiye

² Department of Physics, Ege University, Izmir, Türkiye

ibrahim.senturk@ege.edu.tr, engin.calik@ege.edu.tr, huseyin.sirin@ege.edu.tr, tahsin.oner@ege.edu.tr

key-words: State-Measure, algebraic Logic, mathematical modelling.

Abstract:

In this study, we investigate measures within algebraic logic frameworks utilizing the Sheffer stroke operation. We introduce and analyze various types of measures, including state-measures, measure-morphisms, and state-measure-morphisms, establishing their foundational axioms. We develop systematic algorithms to rigorously assess whether a given mapping satisfies the criteria for these measures. Our exploration extends to key algebraic characteristics, such as their antitonic nature and their role in defining deductive systems. Furthermore, we examine the relationship between measures and state functions, like Rie an and Bosbach states, showing that measures can be derived from these states and vice versa. Extremal state-measures are linked with affine homeomorphisms, providing a geometric perspective on the structural attributes of these algebraic logical structures. Theoretical insights are enriched with illustrative examples and proofs, laying a foundation for future research in fields like fuzzy logic, probabilistic reasoning, and the mathematical modeling of uncertainty.

Additionally, integrating this work into practical applications could prove valuable in areas such as uncertainty, fuzzy logic, quantum theory, and deductive physical models. Sheffer stroke algebras and measure-related mathematical structures offer a flexible and abstract tool for modeling the physical world, providing a robust foundation for analyzing complex systems. In practical terms, the next step involves exploring how these measures can be applied concretely, particularly in fields dealing with uncertainty, such as quantum mechanics and statistical physics. These mathematical structures can more accurately represent uncertainty in the physical realm.

Acknowledgments: This work was supported by TÜBİTAK (The Scientific and Technological Research Council of Türkiye) Project Number: 123F459.

References

- [1] Senturk, I. (2021). A View on State Operators in Sheffer Stroke Basic Algebras. *Soft Computing*, 25(17), 11471-11484.
- [2] Oner, T., Senturk, I. (2017). The Sheffer Stroke Operation Reducts of Basic Algebras. *Open Mathematics*, 15(1), 926-935.
- [3] Senturk, I., Oner, T. and Saeid, A. B. (2020). Congruences of Sheffer Stroke Basic Algebras. *Analele Universitatii "Ovidius" Constanta-Seria Matematica*, 28(2), 209-228.
- [4] Sheffer, H. M. (1913). A Set of Five Independent Postulates for Boolean Algebras, with Application to Logical Constants. *Transactions of the American Mathematical Society*, 14(4), 481-488.
- [5] Hajda, I. (2005). Sheffer Operation in Ortholattices. *Acta Universitatis Palackianae Olomucensis, Facultas Rerum Naturalium Mathematica*, 44, 19-23.
- [6] Dvureenskij, A. (2002). Measures on Quantum Structures. In *Handbook of Measure Theory*, pp. 827-868., North-Holland.
- [7] Pulmannova, A., Dvureenskij, A. (2000). *New Trends in Quantum Structures. Mathematics and its Applications*, 516.
- [8] Ciungu, L. C., Dvureenskij, A. (2010). Measures, States and de Finetti maps on Pseudo BCK-algebras. *Fuzzy Sets and Systems*, 160, 1099-1113.

3rd International Conference: Constructive Mathematical Analysis

Mappings Contracting Perimeter of Triangles in Perturbed Metric Spaces

Mirela Adriana Tarnoveanu

Faculty of Mathematics and Computer Science, Transilvania University of Braşov, Braşov, Romania

mirela.tarnoveanu@unitbv.ro

key-words: Fixed point, mappings contracting triangles, perturbed metric spaces.

Abstract:

In the present paper we introduce the notion of mappings contracting perimeter of triangles in perturbed metric spaces, which we call perturbed mappings contracting perimeters of triangles. We provide a fixed point result for such mappings. We illustrate that our results are more general with some examples.

References

- [1] S. Banach, Sur les operations dans les ensembles abstraits et leur application aux equations integrales, *Fundamenta Mathematicae* 3 (1922), 133-181.
- [2] RK Bisht, E Petrov, Three point analogue of Ćirić-Reich-Rus type mappings with non-unique fixed points, *The Journal of Analysis*, 1-19
- [3] S. K. Chatterjea, Fixed-point theorems, *C.R. Acad. Bulgare Sci.* 25 (1972), 727-730.
- [4] L.B. Ćirić, A generalization of Banach's contraction principle. *Proc. Am. Math. Soc.* 45 (1974), 267-273.
- [5] S. Czerwik, Contraction mappings in b-metric spaces, *Acta Math. Inform. Univ. Ostrav.*, 1993, 1(1), 5-11.
- [6] G.E. Hardy, T.D. Rogers, A generalization of a fixed point theorem of Reich. *Canad. Math. Bull.* 16 (1973), 201-206.
- [7] M. Jleli, B. Samet, M. Jleli and B. Samet, On a new generalization of metric spaces, *J. Fixed Point Theory Appl.*, 2018, 20, 128.
- [8] M. Jleli, B. Samet. On Banach's fixed point theorem in perturbed metric spaces. *Journal of Applied Analysis & Computation*, 2025, 15(2): 993-1001.
- [9] M. Jleli, C.M. Pacurar, B. Samet, New directions in fixed point theory in Gmetric spaces and applications to mappings contracting perimeters of triangles, *arXiv preprint (2024) arXiv:2405.11648*.
- [10] R. Kannan, Some results on fixed point e II, *Amer. Math. Monthly*, 76 (1969), pp. 405-408.
- [11] A. Meir, E. Keeler, A theorem on contraction mappings. *J. Math. Anal. Appl.* 28 (1969), 326-329
- [12] C.M. Pacurar, O. Popescu, Fixed points for three point generalized orbital triangular contractions, *arXiv preprint (2024) arXiv:2404.15682*.
- [13] C.M. Pacurar, O. Popescu, Fixed point theorem for generalized Chatterjea type mappings, *Acta Mathematica Hungarica* 173 (2), 500-509.
- [14] E. Petrov, Fixed point theorem for mappings contracting perimeters of triangles. *J. Fixed Point Theory Appl.* 25 (2023), 1-11.
- [15] E. Petrov, R. K. Bisht, Fixed point theorem for generalized Kannan type mappings, *arXiv:2308.05419*.
- [16] E. Petrov, Periodic points of mappings contracting total pairwise distance, *arXiv:2402.02536*. 10
- [17] A. Petrusel, G. Petrusel, G. (2022). Fixed Point Theory in Graph Metric Spaces. In: Daras, N.J., Rassias, T.M. (eds) *Approximation and Computation in Science and Engineering*. Springer Optimization and Its Applications, vol 180. Springer, Cham.
- [18] O. Popescu, Some remarks on the paper Fixed point theorems for generalized contractive mappings in metric spaces. *J. Fixed Point Theory Appl.* 23 (2021), 1-10.
- [19] O. Popescu, C.M. Pacurar, Mappings contracting triangles, *arXiv preprint (2024) arXiv:2403.19488*.
- [20] Proinov, P.D.: Fixed point theorems for generalized contractive mappings in metric spaces. *J. Fixed Point Theory Appl.* 22 (2020), 1-27.
- [21] I.A. Rus, *Principles and Applications of the Fixed Point Theory* (in Romanian), Editura Dacia, Clui-Napoca, 1979.

3rd International Conference: Constructive Mathematical Analysis

Robust numerical method for the singularly perturbed problem with a nonlocal condition

Zelal Temel

Department of Mathematics, University of Sıtkı Koçman, Muğla, Türkiye

zelaltemel65@gmail.com

key-words: Numerical Method, finite difference method, uniform mesh.

Abstract:

We study a useful numerical method for the singularly perturbed problem with a non-local condition. A small positive parameter is used to determine the coefficients of the terms containing the highest-order derivative in singular perturbation situations. A uniform mesh is used to build finite difference schemes. We examine these schemes' uniform convergence and stability in the discrete maximum norm with respect to the perturbation parameter. The provided numerical experiments validate the accuracy of the proposed method.

References

- [1] Kudu M., Amiraliyev G. M. (2015). Finite difference method for singularly perturbed differential equations with integral boundary condition. *Int J Math Comput.*, 26(3), 72-79.
- [2] Debela, H. G., Duressa, G. F. (2020). Accelerated fitted operator finite difference method for singularly perturbed delay differential equations with non-local boundary condition. *Journal of the Egyptian Mathematical Society*, 28, 1-16.
- [3] Elango, S., Tamilselvan, A., Vadivel, R., Gunasekaran, N., Zhu, H., Cao, J. and Li, X. (2021). Finite difference scheme for singularly perturbed reaction diffusion problem of partial delay differential equation with nonlocal boundary condition. *Advances in Difference Equations*, 2021, 1-20.

3rd International Conference: Constructive Mathematical Analysis

An Approximation Study with Mellin Integral Operators

Cem Topuz¹, Firat Ozsarac¹ and Ali Aral¹

¹ Department of Mathematics, Kirikkale University, Kirikkale, Türkiye

topuzcem@gmail.com, firat_ozsarac@hotmail.com, aliaral73@yahoo.com

key-words: Logarithmic moments, order of approximation, Voronovskaya-type theorem.

Abstract:

In this presentation, we give an overview of some results from the literature and introduce a modification of Mellin convolution-type operators. In this way, we obtain the rate of convergence with the modulus of the continuity of the m th-order Mellin derivative of function f , but without the derivative of the operator. We show moments of the new operators and describe the behavior of the modified operators. Then, we express the Taylor formula including Mellin derivatives with integral remainder. Later, a Voronovskaya-type theorem is proved. In the last part, we state order of approximation of the modified operators, and the obtained results are restated for the Mellin-Gauss-Weierstrass operator.

References

- [1] Aral, A., Acar, T. and Kursun, S. (2022). Generalized Kantorovich Forms of Exponential Sampling Series. Anal. Math. Phys., 12, 1-19.
- [2] Bardaro, C., Mantellini, I. (2007). Voronovskaya-type Estimates for Mellin Convolution Operators. Results Math., 1, 1-16.
- [3] Bardaro, C., Mantellini, I. (2011). A Note on the Voronovskaja Theorem for Mellin-Fejer Convolution operators. Appl. Math. Lett., 24, 2064-2067.
- [4] Bardaro, C., Mantellini, I. (2012). On Voronovskaja formula for Linear Combinations of Mellin-Gauss-Weierstrass Operators. Appl. Math. Comput., 218, 10171-10179.
- [5] Butzer, P. L., Jansche, S. (1997). A Direct Approach to the Mellin Transform. J. Fourier Anal. Appl., 3, 325-375.
- [6] Kolbe, W., Nessel, R. J. (1972). Saturation Theory in Connection with Mellin Transform Methods. SIAM J. Math. Anal., 3, 246-262.
- [7] Mamedov, R. (1991). The Mellin Transform and Approximation Theory, Elm, Baku.

3rd International Conference: Constructive Mathematical Analysis

Approximation Properties of Neural Network and Sampling Kantorovich Operators in Terms of Image Reconstruction

Ana Maria Acu¹, Mihai Ilina¹, Florin Sofonea¹, Arianna Travaglini^{1,2} and Gianluca Vinti^{1,3}

¹ Department of Mathematics and Informatics, University of Sibiu, Sibiu, Romania

² Department of Engineering, University of Basilicata, Potenza, Italy

³ Department of Mathematics and Computer Science, University of Perugia, Perugia, Italy

anamaria.acu@ulbsibiu.ro, mihai.ilina@ulbsibiu.ro, florin.sofonea@ulbsibiu.ro,
arianna.travaglini@unibas.it, gianluca.vinti@unipg.it

key-words: Neural Network operators, sampling Kantorovich operators, digital image processing.

Abstract:

In this talk, we present a comparative study between two powerful image reconstruction algorithms: the multivariate Neural Network (NN) operators, recently implemented for image reconstruction in [1], and the well known sampling Kantorovich (SK) operators (see, e.g. [4]). NN operators have recently attracted considerable attention, and their approximation properties have been thoroughly studied (see, e.g. [3, 5]). On the other hand, SK operators, known for their effectiveness in approximating not necessarily continuous signals, have proven to be successful in various applied fields (see, e.g. [2, 6]). We implement and analyze both methods on a dataset of reference images, evaluating their performances using three widely recognized similarity metrics: SSIM, PSNR, and the recently introduced FLIP index. Our results highlight the strengths and potential applications of each approach in the context of digital image processing.

References

- [1] Acu, A. M., Ilina, M., Sofonea, F., Travaglini, A. and Vinti, G. (2025). A Comparison Between Neural Network and Sampling Kantorovich Operators in terms of Image Reconstruction. *Lecture Notes in Computer Science*, (Accepted).
- [2] Asdrubali, F., Baldinelli, G., Bianchi, F., Costarelli, D., D'Alessandro, F., Scrucca, F., Seracini, M. and Vinti, G. (2021). Mathematical Models for the Improvement of Detection Techniques of Industrial Noise Sources from Acoustic Images. *Mathematical Methods in the Applied Sciences*, 44(13), 10448-10459.
- [3] Cheang, G. H. L. (2010). Approximation with Neural Networks Activated by Ramp Sigmoids. *J. Approx. Theory*, 162, 1450-1465.
- [4] Costarelli, D., Vinti, G. (2011). Approximation by Multivariate Generalized Sampling Kantorovich Operators in the Setting of Orlicz Spaces. *Bollettino dell'Unione Matematica Italiana*, 4(3), 445-468.
- [5] Costarelli, D., Vinti, G. (2017). Convergence for a Family of Neural Network Operators in Orlicz Spaces. *Math. Nachr.*, 290(2-3), 226-235.
- [6] Travaglini, A., Vinti, G., Scalera, G.B. and Scialpi, M. (2023). A Large Scale Analysis for Testing a Mathematical Model for the Study of Vascular Pathologies. *Mathematics*, 11(8), Art. Number 1831.

3rd International Conference: Constructive Mathematical Analysis

New Generalization of Sampling Type Series and Applications to Image Processing

Metin Turgay¹, Tuncer Acar¹

¹Department of Mathematics, Selçuk University, Konya, Türkiye

metinturgay@yahoo.com, tunceracar@ymail.com

key-words: Sampling type operators, weighted approximation, rate of convergence, image processing.

Abstract:

In this presentation, we introduce modified sampling-type operators that generalize the classical sampling operators by incorporating a transformation function ρ . The talk begins with essential definitions, including the introduction of the weighted modulus of continuity and its fundamental properties. The newly constructed operators are analyzed in terms of pointwise and uniform convergence within spaces of continuous functions. We explore the weighted approximation properties of this family of operators in weighted function spaces defined by ρ . Furthermore, we examine the modular convergence of these operators in Orlicz spaces $L^\eta(\mathbb{R}^2)$. Lastly, we present certain ρ -kernels and their applications in image processing, demonstrating that the newly constructed operators provide improved performance and achieve higher PSNR values for specific parameters w and function ρ .

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports. Authors have been supported by Scientific and Technological Research Council of Turkey (TUBITAK) under the Grant Number 1001-Project 123F123. The authors thank to TUBITAK for their supports.

References

- [1] C. Bardaro, P. L. Butzer, R. L. Stens, G. Vinti, Kantorovich-type generalized sampling series in the setting of Orlicz spaces, *Sampl. Theory Signal Image Process.*, 6 (1), (2007), 29–52.
- [2] D. Costarelli, M. Seracini, G. Vinti, A comparison between the sampling Kantorovich algorithm for digital image processing with some interpolation and quasi-interpolation methods, *Appl. Math. Comput.*, 374 (2020), Article ID: 125046.
- [3] M. Turgay, T. Acar, Approximation by Modified Generalized Sampling Series, *Mediterr. J. Math.*, 21 (2024), Article ID: 107.
- [4] M. Turgay, T. Acar, Modified Sampling Kantorovich Operators in Weighted Spaces of Functions, (2025), *FILOMAT*, accepted.
- [5] M. Turgay, T. Acar, Bivariate modified sampling Kantorovich operators and their applications to image processing, (2025), *Math. Methods Appl. Sci.*, accepted, DOI: 10.1002/mma.70025.

3rd International Conference: Constructive Mathematical Analysis

Geometric Evolution of Space Curves According to Positional Adapted Frame in Euclidean 3-Space

Gökhan Türk¹, Muhammed T. Sariaydin¹

¹Selcuk University, Department of Mathematics, Konya, Türkiye

gturk53@gmail.com, talatsariaydin@gmail.com

key-words: Inextensible flow, positional adapted frame, Euclidean space.

Abstract:

In this study, the inextensible flows of a spatial curve obtained using a positional adapted frame in Euclidean 3-space. Then, surfaces constructed with the help of the positional adapted frame, and the flows of these surfaces through inextensible curves examined. Finally, to verify the theoretical findings, various examples provided and visualized using drawing software.

References

- [1] Aslan, S., Yaylı, Y. (2016). Canal Surfaces with Quaternions. *Advances in Applied Clifford Algebras*, 26, 31-38.
- [2] Baş, S., Körpınar, T. (2013). Inextensible Flows of Spacelike Curves on Spacelike Surfaces According to Darboux Frame in $\mathbb{M}_1^3$. *Bol. Soc. Paran. Mat.*, 31, 9–17.
- [3] Baş, S., Körpınar, T., Asil, V. and Sariaydin, M. T. (2016). On the Quaternionic Inextensible Flows of Curves in Euclidean Space $\mathbb{E}^4$. *International Conference on Mathematics and Mathematics Education (ICMME-2016)*, Firat University, Elazığ, 12-14 May 2016.
- [4] Bizzarri, M., Lávička, M. (2013). Parameterizing Rational Offset Canal Surfaces via Rational Contour Curves. *Computer-Aided Design*, 45(2), 342-350.
- [5] Bozok, H. G. (2021). A New Approach for Inextensible Flows of Curves in Pseudo-Galilean Space G_3^1 . *Konuralp Journal of Mathematics*, 9(1), 19-23.
- [6] Bozok, H. G., Dostuoğlu, O. T. (2023). Inextensible Flows of Framed Curves in 4-Dimensional Euclidean Space. *Advanced Studies: Euro-Tbilisi Mathematical Journal*, 16(4), 67-75.
- [7] Cho, H. C., Choi, H. I., Kwon, S. H., Lee, D. S. and Wee, N. S. (2004). Clifford Algebra, Lorentzian Geometry, and Rational Parametrization of Canal Surfaces. *Computer Aided Geometric Design*, 21(4), 327–339.
- [8] Gaber, S., Al Elaiw, A. (2023). Inextensible Flows of Null Cartan Curves in Minkowski Space $\mathbb{R}^{2,1}$. *Universe*, 9(3), 125.

3rd International Conference: Constructive Mathematical Analysis

Approximation of Continuous Functions by Goodman-Sharma Type Modifications of the Bernstein Operator and the Baskakov Operator

Ivan Gadjev¹, Parvan Parvanov¹ and Rumen Uluchev¹

¹Faculty of Mathematics and Informatics, Sofia University “St. Kliment Ohridski”, Sofia, Bulgaria

gadjev@fmi.uni-sofia.bg, pparvan@fmi.uni-sofia.bg, rumenu@fmi.uni-sofia.bg

key-words: Bernstein operator, Baskakov operator, Goodman-Sharma operator, direct theorem, strong converse theorem, K-functional.

Abstract:

We suggest Goodman-Sharma type modifications of the Bernstein operator for approximation of continuous functions on $[0, 1]$ and of the Baskakov operator for approximation of continuous functions on $[0, \infty)$. For the related differential operators and the approximation operators we obtained nice commutative properties in each of the cases. Also, we proved direct and strong converse theorems with respect to the associated K-functionals, see [2, 3]. Although the operators we consider are non-positive, they have the advantage of $O(n^{-2})$ rate for the approximation error compared to $O(n^{-1})$ of the classical operators.

Particular results concerning the topic was obtained by Acu and Agrawal [1] for the case of the Bernstein operator and by Finta [4], Jabbar and Hassan [5] for the Baskakov operator, etc.

Anyway our approach is much more general, covers both cases, and the results are stronger.

In addition, numerical results of the approximation by the Baskakov-Goodman-Sharma operator are given, as well as computational aspects are discussed.

Acknowledgements: This study is financed by the European Union-NextGenerationEU, through the National Recovery and Resilience Plan of the Republic of Bulgaria, project No BG-RRP-2.004-0008.

References

- [1] Acu, A. M., Agrawal, P. (2019). Better approximation of functions by genuine Bernstein-Durrmeyer type operators. *Carpathian J. Math.*, 35(2), 125–136.
- [2] Gadjev, I., Parvanov, P., Uluchev, R. (2024). Higher order approximation of functions by modified Goodman-Sharma operators. *Constr. Math. Anal.*, 7(4), 180–195.
- [3] Gadjev, I., Parvanov, P., Uluchev, R. (2025). On a new modification of Baskakov operators with higher order of approximation. *Mathematics*, 13(1), 64.
- [4] Finta, Z. (2005). On converse approximation theorems. *J. Math. Anal. Appl.*, 312, 159–180.
- [5] Jabbar, A. F., Hassan, A. K. (2024). Better approximation properties by new modified Baskakov operators. *J. Appl. Math.*, 2024, 8693699.

3rd International Conference: Constructive Mathematical Analysis

A Generalized Approach to the Definition of Baskakov-Durrmeyer Operators

Gülsüm Ulusoy Ada

Institute of Graduate Studies, Cankiri Karatekin University, Cankiri, Türkiye

gulsumulusoy@karatekin.edu.tr

key-words: Baskakov-Durrmeyer operators, uniform convergence, weighted modulus of continuity.

Abstract:

In the current study, we introduce a novel class of Baskakov-Durrmeyer operators, expanding the existing framework of positive linear operators. We derive sufficient conditions under which these operators exhibit weighted uniform convergence on the semi-infinite interval $[0, \infty)$, and provide precise quantitative estimates within appropriately defined weighted function spaces. Additionally, by establishing a Voronovskaya-type asymptotic formula, we rigorously prove the pointwise convergence behavior of the operators, thereby gaining deeper insight into their approximation properties.

References

- [1] Acar, T., Aral A. and Raşa I. (2014). Modified Bernstein-Durrmeyer Operators. *General Mathematics*, 22 (1), 27-41.
- [2] Acar, T., Ulusoy G. (2016). Approximation by Modified Szász-Durrmeyer Operators. *Period. Math. Hungaria*, 72 (1), 64-75.
- [3] Baskakov V. (1957). An Instance of a Sequence of Linear Positive Operators in the Space of Continuous Functions. *Doklady Akademii Nauk SSSR.*, 113 (2), 249-251.
- [4] Acar, T., Gupta V. and Aral, A. (2011). Rate of Convergence for Generalized Szász Operators. *Bulletin of Mathematical Science*, 1 (1), 99-113.
- [5] Aral, A., Inoan D. and Raşa, I. (2014). On the Generalized Szász-Mirakyan Operators. *Results in Mathematics*, 65, 441-452.
- [6] Aral, A., and Gupta, V. (2011). Generalized Szász Durrmeyer Operators. *Lobachevskii J. Math.*, 32(1), 23-31.
- [7] Holhos A. (2008). Quantitative Estimates for Positive Linear Operators in Weighted Space. *General Math.*, 16(4), 99-110.

3rd International Conference: Constructive Mathematical Analysis

Approximation Properties of Szász Mirakyan Durrmeyer Operators Preserving Exponential Functions

Gülüm Ulusoy Ada

Institute of Graduate Studies, Cankiri Karatekin University, Cankiri, Türkiye

gulsumulusoy@karatekin.edu.tr

key-words: Szász Mirakyan Durrmeyer operators, weighted modulus of continuity, uniform convergence.

Abstract:

This paper serves as a natural continuation of the work presented in [8], where the Szász-Mirakyan-Durrmeyer operators that preserve exponential functions were initially introduced. As a preliminary result, we demonstrate that the sequence of norms of these operators, when acting on weighted function spaces with varying exponential weights, remains uniformly bounded. Subsequently, we derive Korovkin-type approximation theorems grounded in exponential weighted convergence. Finally, we provide a detailed characterization of the uniform approximation errors of the operators with respect to the exponential weights.

References

- [1] Acar, T. (2015). Asymptotic Formulas for Generalized Szász Mirakyan Operators. Appl. Math. Comput., 263, 233-239.
- [2] Acar, T., Aral, A. and Gonska H. (2017). On Szász Mirakyan Operators Preserving e^{2ax} , $a > 0$. Mediterr. J. Math., 14(1), 1-14.
- [3] Acar, T., Aral, A. and Rasa, I. (2016). The New Forms of Voronovskaya's Theorem in Weighted spaces. Positivity, 20(1), 25-40.
- [4] Aral, A., Acar, T., Cardenas Morales, D. and Garrancho, P. (2017). Szász Mirakyan Type Operators which Fix Exponentials. Results Math., 72(3), 1393-1404.
- [5] Aral, A., Deniz E. and Gupta, V. (2016). On the Modification of the Szász Durrmeyer Operators. Georgian Math. J., 23(3), 323-328.
- [6] Boyanov, B. D., Veselinov, V. M. (1970). A Note on the Approximation of Functions in an Infinite Interval by Linear Positive Operators. Bull. Math. Soc. Sci. Math. Roum., 14(62), 9-13.
- [7] Coşkun, T. (2003). Weighted Approximation of Unbounded Continuous Functions by Sequences of Linear Positive Operators. Indian J. Pure Appl. Math., 34(3), 477-485.
- [8] Deniz, E., Aral, A. Gupta V. (2017). Note on Szász Mirakyan Operators Preserving e^{2ax} , $a > 0$. Numerical Functional Analysis and Optimization., 39(2), 201-207.
- [9] Gadziev, A. D. (1976). Theorems of the type of P. P. Korovkin's Theorems. Math. Zametki., 20(5), 781-786.
- [10] Gupta V. (1995). Simultaneous Approximation by Szász Durrmeyer Operators. Math. Student., 64(1-4), 27-36.
- [11] Gupta, V., Malik, N. (2017). Approximation with Certain Szász Mirakyan Operators. Khayyam J.Math., 3(2), 90-97.
- [12] Gupta, V., Pant, R. P. (1999). Rate of Convergence for the Modified Szász Mirakyan Operators on Functions of Bounded Variation. J. Math. Anal. Appl., 233(2), 476-483.
- [13] Holhoş, A. (2010). The Rate of Approximation of Functions in an Infinite Interval by Positive Linear Operators. Stud. Univ. Babeş-Bolyai, MATH., 55(2), 133-142.
- [14] Mazhar, S. M., Totik, V. (1985). Approximation by Modified Szász Operators. Acta Sci. Math., 49(1-4), 257-269.

3rd International Conference: Constructive Mathematical Analysis

Development of a New Efficient Algorithm for Multiplication of Large Integers

Meltem Uzun¹, Ferhat Kuruz² and Nihal Altuntas²

¹Software Engineering, Istanbul Sabahattin Zaim University, Istanbul, Türkiye

²Computer Engineering, Istanbul Gelisim University, Istanbul, Türkiye

meltem.uzun@izu.edu.tr, fkuruz@gelisim.edu.tr, naltuntas@gelisim.edu.tr

key-words: Integer multiplication, complexity, computer arithmetic.

Abstract:

In most of the public key cryptography algorithms it is essential to develop efficient long integer multiplications. To achieve efficient computations in this area, several techniques are developed such as Toom-Cook, Karatsuba, Harvey's method, FFT, etc. In this study, we aim to construct a new effective algorithm for evaluating n th power of big integers. We will use divide and conquer approach to deal with the complicated multiplication process. Our attempts to solve the problem will begin with considering numbers with two-digit and larger numbers with base 10, then we generalize our formula to big integers and all bases s , where $2 \leq s \leq 10$. We will prove our results and support our theoretical findings by MATLAB experiments. Taking into account the complexity degree of the algorithm, implementation of this work will be a useful application for computer science.

Acknowledgements: This research is supported by Istanbul Sabahattin Zaim University grant number 2024-BAP 400-019.

References

- [1] Bodrato, M., Zanoni, A. (2012). A New Algorithm for Long Integer Cube Computation with Some Insight into Higher Powers. Lecture Notes in Computer Science, 34–46.
- [2] Ilter, M. B., Cenk, M. (2017). Efficient Big Integer Multiplication in Cryptography. Int. J. Inf. Secur., 6(4).
- [3] Fürer, M. (2009). Faster Integer Multiplication. SIAM J. Comput., 39(3).

3rd International Conference: Constructive Mathematical Analysis

On Some New Perspectives on Hardy-Amalgam Spaces with Variable Exponents

Lassane Traore¹, Cihan Unal²

¹ Département de Mathématiques-Physique-Chimie, UFR des Sciences Biologiques, Université Peleforo GON COULIBALY, BP 1328 Korhogo, République de Côte d'Ivoire

² Department of Mathematics, Sinop University, Sinop, Türkiye

traorelassaneupgc@gmail.com, cihanunal88@gmail.com

key-words: Amalgam spaces, Hardy-amalgam spaces, variable exponents analysis.

Abstract:

In this work, we present a generalization of Hardy spaces with variable exponents. We manage the functions in these spaces using atoms. Moreover, we investigate some mapping properties of the fractional integral operator. As an application, the boundedness of some operators in Fourier analysis is studied.

References

- [1] Akbulut, A., Guliyev, V. S., Noi, T. and Sawano, Y. (2017). Generalized Hardy-Morrey Spaces. *Z. Anal. Anwend.* 36(2), 129-149.
- [2] Cruz-Uribe, D. V., Fiorenza, A. (2013). *Variable Lebesgue spaces: Foundations and Harmonic Analysis*. Birkhäuser, Heidelberg.
- [3] Diening, L., Harjulehto, P., Hästö, P. and Růžička M. (2011). *Lebesgue and Sobolev Spaces with Variable Exponents*, Springer-Verlag, Berlin.
- [4] Grafakos, L. (2009). *Modern Fourier analysis*. Springer.
- [5] Holland, F. (1975). Harmonic Analysis on Amalgam of L^p and l^q . *J. London Math. Soc.*, 10(2), 295-305.
- [6] Kováčik, O., Rákosník, J. (1991). On Spaces $L^{p(x)}$ and $W^{k,p(x)}$. *Czechoslovak Math. J.*, 41(4), 592-618.
- [7] Nakai, E., Sawano, Y. (2012). Hardy Spaces with Variable Exponents and Generalized Campanato Spaces. *J. Funct. Anal.*, 262, 3665-3748.
- [8] Orlicz, W. (1931). Über konjugierte exponentenfolgen. *Studia Math.*, 3(1), 200-211.
- [9] Stein, E. M., Weiss, G. (1960). On the Theory of Harmonic Functions of Several Variables I: The theory of H^p Spaces. *Acta Math.*, 103, 25-62.
- [10] Wiener, N. (1926). On the Representation of Functions by Trigonometrical Integrals. *Math. Z.*, 24, 575-616.

3rd International Conference: Constructive Mathematical Analysis

Commutator of Anisotropic Sharp Maximal Operator on Lorentz Spaces

Ali Akbulut¹, Necati Uyar¹ and Vagif S. Guliyev^{1,2,3,4}

¹ Department of Mathematics, Kirsehir Ahi Evran University, Kirsehir, Turkiye

² Institute of Applied Mathematics, Baku State University, Baku, Azerbaijan

³ Institute of Mathematics and Mechanics, Baku, Azerbaijan

⁴ Azerbaijan University of Architecture and Construction, Baku, Azerbaijan

aakbulut@ahievran.edu.tr, necati.uyar@avievran.edu.tr, vagif@guliyev.com

key-words: Anisotropic maximal operator, commutator, Lorentz space.

Abstract:

In this talk we consider the commutator of the anisotropic sharp maximal operator $[b, M^{d,\sharp}]$ on the Lorentz spaces $L^{p,q}(\mathbb{R}^n)$. We obtain necessary and sufficient conditions for the boundedness of the commutator of the anisotropic sharp maximal operator $[b, M^{d,\sharp}]$ on the Lorentz spaces $L^{p,q}(\mathbb{R}^n)$ when b belongs to the bounded space of mean oscillations $BMO(\mathbb{R}^n)$, thereby obtaining some new characterizations for some subclasses of $BMO(\mathbb{R}^n)$, see [1, 2].

For $f \in L^1_{\text{loc}}(\mathbb{R}^n)$, the anisotropic maximal operator M^d and the commutator of the anisotropic sharp maximal operator $[b, M^{d,\sharp}]$ are defined by

$$M^{d,\sharp}f(x) = \sup_{r>0} |\mathcal{E}(x, r)|^{-1} \int_{\mathcal{E}(x, r)} |f(y) - f_{\mathcal{E}(x, r)}| dy, \quad [b, M^{d,\sharp}]f(x) = b(x)M^{d,\sharp}f(x) - M^{d,\sharp}(bf)(x),$$

where $\mathcal{E}(x, r)$ is the ellipsoid of radius r centered at $x \in \mathbb{R}^n$ and $|\mathcal{E}(x, r)|$ denotes the Lebesgue measure of $\mathcal{E}(x, r)$. Suppose that f is a measurable function on $\mathbb{R}^n$, then we define

$$f^*(t) = \inf\{s > 0 : d_f(s) \leq t\}, \quad d_f(s) := |\{x \in \mathbb{R}^n : |f(x)| > s\}|, \quad \forall s > 0.$$

The Lorentz space $L^{p,q} \equiv L^{p,q}(\mathbb{R}^n)$, $0 < p, q \leq \infty$ is the collection of all measurable functions f on $\mathbb{R}^n$ such the quantity

$$\|f\|_{L^{p,q}} := \|t^{\frac{1}{p} - \frac{1}{q}} f^*(t)\|_{L^q(0, \infty)}$$

is finite. Clearly $L^{p,p} \equiv L^p$ and $L^{p,\infty} \equiv WL^p$. The functional $\|\cdot\|_{L^{p,q}}$ is a norm if and only if either $1 \leq q \leq p$ or $p = q = \infty$.

Theorem 3 Let $p, q \in (1, \infty)$. The following assertions are equivalent:

(i) $b \in BMO(\mathbb{R}^n)$ and $b^- \in L^\infty(\mathbb{R}^n)$.

(ii) The operator $[b, M^{d,\sharp}]$ is bounded on $L^{p,q}(\mathbb{R}^n)$.

(iii) There exist a constant $C > 0$ such that $\sup_{\mathcal{E}} \frac{\|(b(\cdot) - 2M^{d,\sharp}(b)(\cdot))\chi_{\mathcal{E}}\|_{L^{p,q}(\mathbb{R}^n)}}{\|\chi_{\mathcal{E}}\|_{L^{p,q}(\mathbb{R}^n)}} \leq C$.

(iv) There exist a constant $C > 0$ such that $\sup_{\mathcal{E}} |\mathcal{E}|^{-1} \|(b(\cdot) - 2M^{d,\sharp}(b)(\cdot))\chi_{\mathcal{E}}\|_{L^1(\mathbb{R}^n)} \leq C$.

References

- [1] Akbulut, A., Isayev, F.A., Serbetci, A. (2024). Anisotropic maximal commutator and commutator of anisotropic maximal operator on Lorentz spaces. Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., Mathematics 44(4), 5-12.
- [2] Guliyev, V.S. (2024). Maximal commutator and commutator of maximal operator on Lorentz spaces. Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., Mathematics 44(4), 43-49.

3rd International Conference: Constructive Mathematical Analysis

Characteristically Near Groups in Complex Plane Vector Fields

Tane Vergili¹, James F. Peters²

¹ Department of Mathematics, Karadeniz Technical University, Trabzon, Türkiye

² Department of Electrical and Computer Engineering, University of Manitoba, WPG, Manitoba, Canada

tane.vergili@ktu.edu.tr, james.peters3@umanitoba.ca

key-words: Characteristic, complex plane, distance, group, nearness, vector field.

Abstract:

Characteristically near groups arise from variants of the Stein-Weiss characteristic function $\varphi(x) = e^{\pm ix}$, $x \in \mathbb{R}$ from [3], which defines the characteristic of a multiplicative abelian group $(\{\varphi(x)\}, \cdot)$, i.e.,

$$\varphi : \Omega \in 2^{\mathbb{C}} \rightarrow \mathbb{C} \text{ defined by } \varphi(x) = e^{2ix} : (G, \cdot) = (\{\varphi(z)\}, \cdot).$$

Characteristic functions were introduced by W.R. Hamilton in 1837 in a study of light rays [1]. The characteristic vector $\vec{\Phi}(G) = (\varphi_1(G), \varphi_2(G), \dots, \varphi_n(G))$ contains characteristics $\varphi_i(t \in \Omega) \in [0, 1]$, which reflect the structure G , e.g.,

$$\varphi_1(G) = k \in \{0 \text{ (not in unit circle in } \mathbb{C}), 1 \text{ (in unit circle in } \mathbb{C})\}.$$

$$\varphi_2(G) = k \in \{0 \text{ (not abelian), } 1 \text{ (abelian)}\}.$$

$$\varphi_3(G) = o(G) \in \mathbb{Z}^{0+} \text{ (order of } G).$$

A characteristic distance (denoted by $d^{\Phi} : 2^X \times 2^Y \rightarrow \mathbb{C}$ [2]) is used to measure of the nearness of vector field groups, defined by

$$d^{\Phi}(A, B) = \inf_{\substack{a \in A \\ b \in B}} \{|\varphi(a) - \varphi(b)|\} = \varepsilon \in [0, 1], (\varphi(a), \varphi(b)) \in \Phi(A) \times \Phi(B).$$

Vector field groups $\vec{\Phi}(G_1), \vec{\Phi}(G_2)$ are near, provided $d^{\Phi}(\vec{\Phi}(G_1), \vec{\Phi}(G_2)) = 0$.

References

- [1] Hamilton, W. R. (1837). Essay on the Theory of Rays, Trans. Royal Irish Acad., 17, 1-144.
- [2] Peters, J. F., Vergili, T., Ucan, F. and Vakeesan, D. (2025). Indefinite Descriptive Proximities Inherent in Dynamical Systems. Cornell arXiv, 2501.02585 (2025), url=<https://arxiv.org/abs/2501.02585>.
- [3] Stein, E. M., Weiss, G. (1971). Introduction to Fourier Analysis on Euclidean Spaces. Mathematical series, no. 32, Princeton U. Press, Princeton.

3rd International Conference: Constructive Mathematical Analysis

Local Fractional Hilbert-Type Inequalities

Predrag Vuković

Faculty of Teacher Education, University of Zagreb, Zagreb, Croatia

predrag.vukovic@ufzg.hr

key-words: Hilbert inequality, conjugate parameters, homogeneous function, local fractional calculus.

Abstract:

An intriguing topic related to classical inequalities is their application to fractal spaces through local fractional calculus. The principal function of local fractional calculus is to address diverse non-differentiable issues arising in intricate systems of real-world phenomena. The non-differentiability observed in science and engineering has been represented by local fractional ordinary or partial differential equations. The local fractional calculus, while originating from real-world occurrences, is also a significant instrument in pure mathematics.

Recently, some classical real inequalities have been extended to apply within specific fractal environments. A comprehensive compilation of generalisations encompassed inequalities featuring more generalised kernels, weight functions, and integration domains, as well as extensions to a multidimensional context. Special attention is given to a category of inequalities featuring a homogenous kernel. Specifically, one establishes some weak circumstances under which the constants on the right-hand sides of these local fractional inequalities are optimal. Additionally, improved and inverted relations are derived in a general multidimensional setting.

The primary aim of this paper is to investigate new local fractional Hilbert-type inequalities. We use our general results on non-homogeneous kernels. The equivalent form and several specific cases are derived as applications.

This work is supported by Croatian Science Foundation under the project HRZZ-IP-2024-05-3882

References

- [1] Hardy, G. H., Littlewood, J. E., and Pólya, G. (1967). *Inequalities*, Second Edition, Cambridge University Press, Cambridge.
- [2] Krnić, M., Pečarić, J., Perić, I. and Vuković, P. (2012). *Recent Advances in Hilbert-type Inequalities*, Element, Zagreb.
- [3] Yang, X. J. (2011). *Local Fractional Functional Analysis and Its Applications*. Asian Academic Publisher Limited, Hong Kong.
- [4] Yang, X. J. (2012). *Advanced Local Fractional Calculus and Its Applications*. Word Science Publisher, New York.
- [5] Vuković, P., Yang, W. (2025). *Discrete Local Fractional Hilbert-type Inequalities*. Accepted in Kragujevac Journal of Mathematics.

3rd International Conference: Constructive Mathematical Analysis

Convergence Types on Interpolative Metric Spaces

Ceylan Yalcin

Department of Mathematics, Cankaya University, Ankara, Türkiye

cyalcin@cankaya.edu.tr

key-words: Summability Methods, interpolative metric spaces, statistical convergence.

Abstract:

In recent years, interpolative metric spaces have been developed as a flexible expansion of classical metric spaces, supporting the study of convergence theory and fixed point theory in expanded situations [3]. This study investigates the relationship between convergence types approaches in the context of interpolative metric spaces [2], [4]. Additionally, we present illustrative examples and establish first results that may serve as a basis for further research in the field of convergence theory and fixed point theory [1]. This research contributes to a comprehensive approach aimed at unifying various convergence techniques into generalized metric frameworks.

References

- [1] Sakalli, H. (2008). Statistical Convergence in Topological Groups. *Mathematical and Computer Modelling*, 48(9-10), 1275-1281.
- [2] Fridy, J. A. (1985). On Statistical Convergence. *Analysis*, 5, 301-313
- [3] Karapınar, E. (2024). On Interpolative Metric Spaces. *Filomat*, 38(22), 7729-7734.
- [4] Kostyrko, P., Maaj, M., ‘alst, T. and Wilczyński, W. (2001). I-convergence. *Real Analysis Exchange*, 26(2), 669-685.

3rd International Conference: Constructive Mathematical Analysis

Extended Semi-Parallel Tensor Product Surfaces in $\mathbb{E}_2^4$

Aziz Yazla

Department of Mathematics, Selcuk University, Konya, Türkiye

aziz.yazla@selcuk.edu.tr

key-words: Tensor product surface, Gaussian curvature, extended semi-parallel.

Abstract:

Tensor product immersion of two immersions of a given Riemannian manifold has begun to be studied by B.Y. Chen. In the light of Chen's definition, many researchers studied the tensor product of two immersions. Then, tensor product surfaces of Euclidean plane curves were investigated by Mihai and Rouxel. Moreover, tensor product surfaces of Lorentzian plane curves were investigated by Mihai et al. Recently Bulca and Arslan studied some special semi-parallel surfaces in Euclidean spaces. Further, Yildirim and Ilarslan considered tensor product surfaces in 4-dimensional semi-Euclidean space with index 2, $\mathbb{E}_2^4$, satisfying the semi-parallelity condition. In present study, we consider and characterize tensor product surfaces in $\mathbb{E}_2^4$ satisfying the extended semi-parallelity condition.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Bulca, B., Arslan, K. (2014). Semiparallel Tensor Product Surfaces in $\mathbb{E}_2^4$. Int. Electron. J. Geom., 7(1), 36-43.
- [2] Mihai, I., Rouxel, B. (1995). Tensor Product Surfaces of Euclidean Plane Curves. Results in Mathematics, 27(3-4), 308-315.
- [3] Mihai, I., Van de Woestyne, I., Verstraelen, P. and Walrave, J. (1995). Tensor Product Surfaces of Lorentzian Planar Curves. Bull. of the Inst. of Math. Acad. Sinica, 23(4), 357-363.
- [4] Yildirim, M., Ilarslan, K. (2016). Semi-Parallel Tensor Product Surfaces in $\mathbb{E}_2^4$. Comm. Fac. of Sci. University of Ankara Series A1 Mathematics and Statistics, 65(2), 133-142.

3rd International Conference: Constructive Mathematical Analysis

Power Series Statistical Relative Uniform Convergence of a Double Sequence of Functions at a Point and Approximation Results

Sevda Yıldız

Department of Mathematics, Sinop University, Sinop, Türkiye

sevdaorhan@sinop.edu.tr

key-words: Double sequence, Korovkin theorem, power series statistical convergence.

Abstract:

In this presentation, we introduce a novel concept in statistical convergence for double sequences, named power series statistical uniform convergence at a point, defined with respect to the power series method. We establish an approximation theorem for sequences of functions under this convergence. Furthermore, we provide an illustrative example that satisfies our new theorem but fails to hold under previously studied convergence methods. Additionally, we investigate the rate of convergence, offering a quantitative analysis of the proposed method's efficiency.

References

- [1] Demirci, K., Boccuto, A., Yıldız, S. and Dirik, F. (2020). Relative Uniform Convergence of a Sequence of Functions at a Point and Korovkin-Type Approximation Theorems. *Positivity*, 24(1), 1-11.
- [2] Demirci, K., Dirik, F. and Yıldız, S. (2023). Approximation via Statistical Relative Uniform Convergence of Sequences of Functions at a Point with respect to Power Series Method. *Afrika Matematika*, 34(3), 39.
- [3] Demirci, K., Orhan, S. (2016). Statistically Relatively Uniform Convergence of Positive Linear Operators. *Results Math.*, 69, 359-367.
- [4] Dirik, F., Demirci, K. (2010). Korovkin Type Approximation Theorem for Functions of Two Variables in Statistical Sense. *Turk. J. Math.*, 34, 73-83.
- [5] Dirik, F., Demirci, K., Yıldız, S. and Acu, A. M. (2021). The Uniform Convergence of a Double Sequence of Functions at a Point and Korovkin-type Approximation Theorems. *Georgian Mathematical Journal*, 28(4), 567-573.
- [6] Gadjiev, A. D., Orhan, C. (2002). Some Approximation Theorems via Statistical Convergence. *The Rocky Mountain Journal of Mathematics*, 32(1), 129-138.
- [7] Klippert, J., Williams, G. (2002). Uniform Convergence of a Sequence of Functions at a Point. *Internat. J. Math. Ed. Sci. Tech.*, 33(1), 51-58.
- [8] Ünver, M., Orhan, C. (2019). Statistical Convergence with respect to Power Series Methods and Applications to Approximation Theory. *Numerical Functional Analysis and Optimization*, 40(5), 535-547.

3rd International Conference: Constructive Mathematical Analysis

On the Convergence of Generalized Szász-Mirakjan Operators in the Variation Seminorm

Sare Yumrucali¹, Harun Karsli² and Fatma Tasdelen Yesildal¹

¹Department of Mathematics, Ankara University, Ankara, Türkiye

²Department of Mathematics, Bolu Abant İzzet Baysal University, Bolu, Türkiye

sareyumrucali@icloud.com, karsli_h@ibu.edu.tr tasdelen@science.ankara.edu.tr.

key-words: Linear positive operators, generalized Szász-Mirakjan operators, variation seminorm, variation detracting property, rate of convergence, modulus of continuity, convergence in variation seminorm, $TV[I]$ -space, Voronovskaya-type theorem.

Abstract:

The aim of the present paper is to introduce generalized Szász-Mirakjan operators. This article studies the property of variation seminorm and on some approximation properties of generalized Szász-Mirakjan operators not only in normed spaces but also in variation seminorm. We obtain convergence properties of our operators with the help of Korovkin's theorem and the order of convergence by using a classical approach, the second modulus of continuity and Peetre's K -functional. We also give asymptotic formula and the convergence of the derivatives for these operators. We investigate the variation detracting property of generalized Szász-Mirakjan operators. We show the convergence of generalized Szász-Mirakjan operators in variation seminorm and its rate of convergence in variation seminorm.

References

- [1] Bardaro, C., Butzer, P. L., Stens, R. L. and Vinti G. (2003). Convergence in Variation and Rates of Approximation for Bernstein-Type Polynomials and Singular Convolution Integrals. *Analysis*, (23)4, 299-346.
- [2] Stein, E. M. (1957). Function of Exponential Type. *Annals of Mathematics*, 65, 582-592.
- [3] Kreyszig, E. (1978). *Introductory Functional Analysis with Applications*. John Wiley and Sons. Inc., New York, Santa Barbara, London, Sydney, Toronto.
- [4] Mirakjan, G. M. (1941). Approximation des fonctions continues au moyen de polynômes de la forme $e^{-nx} \sum_{k=0}^n C_{n,k} x^k$. *Acad. Sci. URSS(N.S)*, 31, 201-205.
- [5] Butzer, P. L., Karsli, H. (2009). Voronovskaya-type Theorems for Derivatives of the Bernstein-Chlodovsky Polynomials and Szász-Mirakjan Operator. *Comment Math.*, 49(1), 33-57.
- [6] Karsli, H. (2013). On Convergence of Chlodovsky and Chlodovsky-Kantorovich Polynomials in the Variation Seminorm. *Mediterr. J. Math.*, 10, 41-56.
- [7] Kejal, K., Vishnu Narayan, M. (2018). Generalized Szász-Mirakjan Operators Involving Brenke Type Polynomials. 324, 228-238.
- [8] Oksuzer, O., Karsli, H. and Tasdelen, F. (2016). On Convergence of Bernstein-Stancu Polynomials in the Variation Seminorm. *Numer. Funct. Anal. Optim.*, 37(4), 1-20.
- [9] Tasdelen, F., Aktas, R. (2012). Kantorovich Type of Szász Operators Including Brenke-type Polynomials. *Abstr. Appl. Anal.*, 13, 867203.
- [10] Varma, S., Sucu, S., and Icoz, G. (2012). Generalization of Szász Operators Involving Brenke Type Polynomials. *Appl. Math. Comput.*, 64, 121-127.
- [11] Walczak, Z. (2003). On Modified Szász-Mirakjan Operators. *Novi Sad J. Math.*, 33(1), 93-107.
- [12] Tasdelen, F., Cekim, B. and Aktas, R. (2013). Kantorovich-Stancu Type Generalization of Szász Operators Including Brenke Type Polynomials. *J. Funct. Spaces Appl.*, 9, Article ID 935430.
- [13] Altomare, F., Campiti, M. (1994). Korovkin-type Approximation Theory and Its Applications. in: *De Gruyter Studies in Mathematics*, Walter De Gruyter, vol. 17, Berlin, Germany,
- [14] Landau, E. (1913). Einige ungleichungen für zweimal differenzierbare funktionen. *Proc. Lond. Math. Soc.*, 13, 43-49.
- [15] Lorentz, G. G. (1953). *Bernstein Polynomials*. University of Toronto Press, Toronto; 2 ed., Chelsea Publishing Co., New York.
- [16] Gavrea, I., Raşa, I. (1993). Remarks on Some Quantitative Korovkin-type Results. *Rev. Anal. Numer. Theor. Approx.*, 22(2) 173-176.

3rd International Conference: Constructive Mathematical Analysis

A Modification of Bernstein Operators on the Interval $(-1, 1)$

Sule Yuksel Gungor¹, Mediha Orku¹

¹ Department of Mathematics, Gazi University, Ankara, Türkiye

sulegungor@gazi.edu.tr, medihaorku@gazi.edu.tr

key-words: Bernstein operators, modulus of continuity, rate of convergence.

Abstract:

In this talk, we will introduce a modification of Bernstein operators, which are defined on the interval $(-1, 1)$ and we will also discuss the approximation properties of these operators, including the rate of convergence, weighted approximation and asymptotic formulas. Lastly, the approximation will be supported with graphs.

References

- [1] Bernstein, S. N. (1912). Démonstration du théorème de Weierstrass fondée sur le calcul des probabilités. Comm. Kharkov Math. Soc., 13, 1-2.
- [2] Cilo, A. (2012). $[-1, 1]$ aralığında Bernstein Polinomlarının Yaklaşım Özellikleri ve Yaklaşım Hızı, Master thesis, Harran University.
- [3] Usta, F. (2020). On new modification of Bernstein operators: Theory and Applications. Iran J Sci Technol Trans Sci., 44, 1119-1124.

3rd International Conference: Constructive Mathematical Analysis

On Approximation by Algebraic Version of the Trigonometric Jackson Integrals $G_{s,n}$ in Generalized Weighted Integral Metric

Teodora Zapryanova

University of Economics - Varna, bul. "Knyaz Boris I", 77, Varna, Bulgaria
teodorazap@ue-varna.bg

key-words: Linear operator, direct theorem, strong converse theorem, K -functional.

Abstract:

We characterize the errors of approximation of the algebraic version of trigonometric Jackson integrals $G_{s,n}$ in weighted integral metric with weight of the form $(1-x)^{-\frac{1}{2}-\gamma_1}(1+x)^{-\frac{1}{2}-\gamma_2}$ for $\gamma_1, \gamma_2 \in [0, 1/2)$. We prove direct and strong converse theorem in terms of the weighted K -functional. The results extend the case $\gamma_1 = \gamma_2 = 0$ to weights with negative powers.

References

- [1] Adell J.A., Cardenaz - Morales P., *Strong converse inequalities for Bernstein Operators via Krawtchouk Pynomials*. ArXiv: 2309. 11967 (2023).
- [2] Bustamante J., *Baskakov- Kantorovich operators reproducing affine functions*. J. Numer. Anal. Approx. Theory, 51 (1) 2022, 67-82.
- [3] DeVore R. A. and Lorentz G. G., *Constructive Approximation*. Berlin, Springer-Verlag, 1993.
- [4] Ditzian Z. and Ivanov K., *Strong Converse Inequalities*. J. Anal. Math. **61** (1993), 61–111.
- [5] Draganov B. R., *Strong converse inequalities for the weighted simultaneous approximation by the Szász-Mirakjan operators*. Sertica Math. J., 46 (2020), 261-286.
- [6] Draganov B. R. and Gadjev I., *Approximation of functions by Szász-Mirakjan-Kantorovich operators*. Numerical Functional Analysis and Optimization, Vol. 40, 2019, 803-824.
- [7] Gadjev I., *Strong converse result for uniform approximation by Meyer-Konig and Zeller operator*. J. Math. Anal. Appl., 428 (2015), 32-42.
- [8] Ivanov K., *A Characterization Theorem for the K -functional for Kantorovich and Durrmeyer Operators*, in "Approximation theory": A volume dedicated to Borislac Bojanov, Marin Drinov Academic Publishing House, Sofia, (2004), 112-125.
- [9] Lorentz G., *Approximation of Functions*. Holt-Rinehart and Winston, New York, 1966.
- [10] Totik V., *Strong converse inequalities*. J. Approx. Theory 76, (1994), 369-375.

3rd International Conference: Constructive Mathematical Analysis

Estimating Covariance in Generalized Least Squares for Frechet Distribution

Khemissi Zahia

Department of Mathematics, University of Khaider Mohamed, Biskra, Algeria

zahia.khemissi@univ-biskra.dz

key-words: Generalized least squares, weighted least squares, covariance matrices.

Abstract:

The Frechet (extreme value type II) distribution is one of the probability distributions used to model extreme events. We consider two-parameter Frechet distribution with shape parameter α , scale parameter β .

After some algebraic manipulation and for a sample size of n and $x_{(1)} \leq x_{(2)} \leq \dots \leq x_{(n)}$ the regression model is rewritten as:

$$\ln x_{(i)} = -\ln \beta - \frac{1}{\alpha} \ln (-\ln (F(x_{(i)}))) \quad (1)$$

If we replace $\ln x_{(i)}$ with Y_i , $-\ln \beta$ with a_0 , $-\frac{1}{\alpha}$ with a_1 and $\ln (-\ln (F(x_{(i)})))$ with X_i , the regression model (1) occurs as:

$$Y_i = a_0 + a_1 X_i \quad (2)$$

In the linear regression model given in (2), if the form of the variance of $Y = (Y_1, \dots, Y_n)$ is $\sigma^2 \mathbf{V} = \Sigma$, GLS minimizes

$$\hat{\theta}_{GLS} = (X^t V X)^{-1} X^t V Y$$

where $\hat{\theta}_{GLS}$ is the vector of the GLS estimates of $\hat{\theta} = (a_0, a_1)$ and X is the matrix of ones and x_i .

Now, the problem is to estimate the $\mathbf{V}$ matrix for the considered distributions. by using Taylor expansion [2], it is possible to approximate the expectation of the observation and also variance and covariance between the observations.

the approximate formula for the $\mathbf{V}$ matrix can be expressed as follows:

$$v_{ij} = \frac{i}{(n+1-i)} \frac{1}{\frac{i}{n+1} (-\ln(\frac{i}{n+1}))} \frac{1}{\frac{j}{n+1} (-\ln(\frac{j}{n+1}))}, \quad i \prec j$$

where v_{ij} is an element of the $\mathbf{V}$ matrix.

References

- [1] Zahia. K., Brahim, B. and Benatia, F. (2022). Heavy Tail Index Estimator through Weighted Least Squares Rank Regression. Journal of Siberian Federal University Mathematics and Physics, 15(6), 797-805.
- [2] Yeliz M. K. (2015). Generalized Least Squares and Weighted Least Squares Estimation. Methods for Distributional Parameters. Revstat Statistical Journal, 13(3), 263-282.

3rd International Conference: Constructive Mathematical Analysis

Numerical Study and Convergence of a Finite Element Method for the Kazhikhov-Smagulov-Korteweg Pollution System

Gafarou Zongo¹, Yibour Corentin Bassonon¹ and Arouna Ouedraogo¹

¹Laboratoire de Mathématiques, Informatique et Applications (LaMIA), Département de Mathématiques, Université Norbert Zongo, Koudougou, Burkina Faso

gafarouzongo91@gmail.com, corentinbassonon@gmail.com, arounaoued2002@yahoo.fr

key-words: Numerical simulation, Kazhikhov-Smagulov-Korteweg equations, viscous incompressible fluid mixture.

Abstract:

The numerical modeling of pollutant transport in mixtures of viscous incompressible fluids plays a crucial role in environmental applications, particularly in wastewater treatment and pollution control. In this study, we conduct a detailed numerical analysis of a Kazhikhov-Smagulov-Korteweg-type model, which describes the behavior of such fluid mixtures with capillary effects and variable density. We first establish a variational formulation of the system that accounts for incompressibility, viscosity, capillarity, and source terms related to pollutant transport. Based on this formulation, we construct a fully discrete scheme using finite element methods for spatial discretization and a second-order Backward Differentiation Formula (BDF2) for time integration, applied to both the velocity and density variables. The proposed scheme is proven to be unconditionally stable using the G -norm framework specific to BDF2 methods. We also prove the convergence of the numerical solution to a global weak solution of the original system. To illustrate the relevance of the model, we simulate a practical scenario involving the dispersion of a chemical pollutant in treated wastewater, where differences in viscosity and local density gradients play a significant role. The results confirm the ability of the method to capture realistic pollutant dynamics and demonstrate its robustness in simulating complex environmental flows.

References

- [1] Akbas, M, Kaya, S. and Rebholz. L. (2017). On the Stability at All Times of Linearly Extrapolated BDF2 Time Stepping for Multiphysics Incompressible Flow Problems. *Numer. Methods Partial Differ. Equ.*, 33, 995-1017.
- [2] Antontsev, S. N., Kazhikhov, A.V. and Monakhov, V. N. (1990). *Boundary Value Problems in Mechanics of Nonhomogeneous Fluids*. Studies in Mathematics and Its Applications, 22, North-Holland, Publishing Co., Amsterdam.
- [3] Cabrales, R. C., Guillén-González, F. and Gutiérrez-Santacreu, J. V. (2011). Stability and Convergence for a Complete Model of Mass Diffusion. *Appl. Numer. Math.*, 61, 1161-1185.

3rd International Conference: Constructive Mathematical Analysis

On a Two-layer Conjugation Problem for the Heat-Conduction Equation in a Cylinder

Kulyash Kaliyeva and Jiayue Wang¹

¹ Al-Farabi Kazakh National University, Almaty, Kazakhstan

Kaliyeva.kulyash@kaznu.kz, susanwjy123@gmail.com

key-words: Two-layer medium, temperature field, unsteady heat conduction, integral transform method, conjugation conditions.

Abstract:

The analytical theory of unsteady heat conduction in a cylindrical surface finds wide application in solving important technical and technological problems of heat- and mass-transfer processes that express energy or material balances. This paper examines an initial-boundary-value problem for the heat-conduction equation on a cylindrical surface with a prescribed initial temperature distribution, boundary conditions, and conjugation conditions on the contact cylindrical surface. The conjugation boundary condition (of the fourth kind) corresponds either to heat exchange between a body's surface and the surrounding medium according to Fourier's law of conduction or to heat exchange between contacting solids when the temperatures of the contacting surfaces are identical (ideal, perfect thermal contact). In addition, when a solid body is washed by a fluid (or gas), the heat transfer from the fluid (gas) to the body's surface in the immediate vicinity of that surface occurs by molecular heat conduction, which likewise leads to heat exchange described by the fourth-kind boundary condition. According to the second law of thermodynamics, a spontaneous heat-transfer process in space arises when there is a temperature difference and proceeds in the direction of decreasing temperature. Thermodynamic processes are accompanied by heat exchange between structural elements within the working space of thermal installations and the surrounding environment, where heat is a quantitative measure of energy; the regularities of these heat-transfer processes are treated by the theory of heat exchange and heat transfer. Using the method of separation of variables together with integral transforms—the Laplace transform with respect to time and the zero-order Bessel transform with respect to the radial coordinate—a closed-form analytical solution to the unsteady two-layer heat-conduction problem is obtained. The investigation of initial-boundary-value problems for the heat-conduction equation and the development of analytical solution methods constitute an important and rapidly evolving area of mathematical physics. Solutions of this type can be applied to compute transient temperature fields and heat fluxes in two-layer sheet products, structures, and buildings, as well as in cylindrical specimens and apparatuses whose thermophysical parameters depend functionally on temperature and on interfacial boundaries. They can also be used to determine the conditions under which two-layer heat-exchange processes proceed, based on temperature fields and the corresponding initial and boundary conditions. The practical significance of the results lies in solving concrete initial-boundary-value problems of thermophysical processes.

References

- [1] Kaliyeva, K. (2014). Energy conservation law in the free atmosphere. *International Journal of Engineering and Innovative Technology (IJEIT)*, 3(11), 50-61.
- [2] Kaliyeva, K. (2014). Existence and Uniqueness of the Navier-Stokes Problem in Infinite Space. *IAENG International Journal of Applied Mathematics* (Proceedings of the World Congress on Engineering 2014, Vol II).
- [3] Kaliyeva, K. (2015). Energy Conservation Law for the Turbulent Motion in the Free Atmosphere. In *Handbook of Research on Computational Simulation and Modeling in Engineering* (pp. 105-138). IGI Global.

3rd International Conference: Constructive Mathematical Analysis

A Numerical Approach Based on a Dimensionality Reduction Procedure for Approximating Multiple Integrals

Mohamed Rahal¹ and Djaouida Guettal²

¹ Laboratory of Fundamental And Numerical Mathematics, Department of Mathematics, University Ferhat Abbas Setif 1, Setif, Algeria

² Laboratory of Fundamental And Numerical Mathematics, Department of Basic Education in Technology, University Ferhat Abbas Setif 1, Setif, Algeria

mrahal@univ-setif.dz, djaouida.guettal@univ-setif.dz

key-words: Multiple integral, numerical integration, α -dense curves.

Abstract:

This paper explores a dimensionality reduction procedure using parametric α -dense curves with a minimal density α for approximating multiple integrals. By utilizing specific α -dense curves $\ell_\alpha(t)$ of finite length, the multiple integral of a positive continuous function f over Ω can be approximated by a single integral that corresponds to the length of $\ell_\alpha(t)$ and the density α . Finally, we provide some numerical examples.

References

- [1] Garcia, G., Mora, G. and Redtwitz, D. A. (2017). Box-Counting Dimension Computed By α -Dense Curves. World Scientific Publishing Company, 25(5), 1750039-1750050.
- [2] Ziadi, A. (2000). Generation of α -dense curves and application to global optimization. Kybernetes, 27(1), 71–82.
- [3] Ziadi, A., Guettal. D. and Cherruault. Y. (2005). Global optimization: The Alienor mixed method with Piyavskii-Shubert technique. Kybernetes, 34(7/8), 1049-1058.

3rd International Conference: Constructive Mathematical Analysis

Approximation Properties of Modified Jakimovski-Leviatan Operators

Kadir Kanat¹, Melek Sofyalıoğlu Aksoy¹ and Selin Erdal Özkan¹

¹Ankara Hacı Bayram Veli University, Department of Mathematics, Ankara, Türkiye
kadir.kanat@hbv.edu.tr, melek.sofyalioğlu@hbv.edu.tr, selin.erdal@hbv.edu.tr

key-words: Jakimovski-Leviatan operators, hypergeometric function.

Abstract:

In this talk, we provide a generalization of the Stancu Type Jakimovski-Leviatan operators using hypergeometric functions. We look into the convergence rate and approximation properties. We illustrate theoretical findings by using numerical examples. Finally, we offer a graphic comparison between the recently constructed operators and the Stancu Type Jakimovski-Leviatan operators.

References

- [1] Özarslan, M.A, Çekim, B. Confluent Appell polynomials, Journal of Computational and Applied Mathematics 424 (2023) 114984.
- [2] Rainville, E.D. Special Functions, The Macmillan Company, New York, 1960.
- [3] Sucu, S., Varma, S. Generalization of Jakimovski-Leviatan type Szasz operators, Appl. Math. Comput. 270 (2015) 977-983.
- [4] Vasić, B. I. Voronovskaja type theorem for some nonpositive Kantorovich type operators. Carpathian Journal of Mathematics, 40(1), (2024) 187-194.



FOR DETAILS

 <https://iccma.selcuk.edu.tr/>

 <https://dergipark.org.tr/en/pub/cma>

CMA
CONSTRUCTIVE MATHEMATICAL ANALYSIS